



Simple rules govern the diversity of bacterial nicotianamine-like metallophores

Clémentine Laffont, Catherine Brutesco, Christine Hajjar, Gregorio Cullia,
Roberto Fanelli, Laurent Ouerdane, Florine Cavelier, Pascal Arnoux

► To cite this version:

Clémentine Laffont, Catherine Brutesco, Christine Hajjar, Gregorio Cullia, Roberto Fanelli, et al..
Simple rules govern the diversity of bacterial nicotianamine-like metallophores. *Biochemical Journal*,
2019, 476 (15), pp.2221-2233. 10.1042/bcj20190384 . cea-02275929

HAL Id: cea-02275929

<https://cea.hal.science/cea-02275929>

Submitted on 3 Sep 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Simple rules govern the diversity of bacterial nicotianamine-like metallophores

Clémentine Laffont¹, Catherine Brutesco¹, Christine Hajjar¹, Gregorio Cullia², Roberto Fanelli²,
Laurent Ouerdane³, Florine Cavelier², Pascal Arnoux^{1,*}

¹Aix Marseille Univ, CEA, CNRS, BIAM, Saint Paul-Lez-Durance, France F-13108.

²Institut des Biomolécules Max Mousseron, IBMM, UMR-5247, CNRS, Université Montpellier,
ENSCM, Place Eugène Bataillon, 34095 Montpellier cedex 5, France.

³CNRS-UPPA, Laboratoire de Chimie Analytique Bio-inorganique et Environnement, UMR 5254,
Hélioparc, 2, Av. Angot 64053 Pau, France.

***Correspondance:** Pascal Arnoux, Aix Marseille Univ, CEA, CNRS, BIAM, Saint Paul-Lez-Durance, France F-13108; pascal.arnoux@cea.fr; Tel. 04-42-25-35-70

Keywords: opine/opaline dehydrogenase, CntM, nicotianamine-like metallophore, bacillopaline

ABSTRACT

In metal-scarce environments, some pathogenic bacteria produce opine-type metallophores mainly to face the host's nutritional immunity. This is the case of staphylopine, pseudopaline and yersinopine, identified in *Staphylococcus aureus*, *Pseudomonas aeruginosa* and *Yersinia pestis* respectively. Depending on the species, these metallophores are synthesized by two (CntLM) or three enzymes (CntKLM), CntM catalyzing the last step of biosynthesis using diverse substrates (pyruvate or α -ketoglutarate), pathway intermediates (xNA or yNA) and cofactors (NADH or NADPH). Here, we explored substrate specificity of CntM by combining bioinformatics and structural analysis with chemical synthesis and enzymatic studies. We found that NAD(P)H selectivity was mainly due to the amino acid at position 33 (*S. aureus* numbering) which ensures a preferential binding to NADPH when it is an arginine. Moreover, whereas CntM from *P. aeruginosa* preferentially uses yNA over xNA, the staphylococcal enzyme is not stereospecific. Most importantly, selectivity towards α -ketoacids is largely governed by a single residue at position 150 of CntM (*S. aureus* numbering): an aspartate at this position ensures selectivity towards pyruvate whereas an alanine leads to the consumption of both pyruvate and α -ketoglutarate. Modifying this residue in *P. aeruginosa* led to a complete reversal of selectivity. Thus, opine-type metallophore diversity is governed by the absence/presence of a *cntK* gene encoding a histidine racemase, and the amino acid residue at position 150 of CntM. These two simple rules predict the production of a fourth metallophore by *Paenibacillus mucilaginosus*, which was confirmed *in vitro* and called bacillopaline.

INTRODUCTION

In metal-scarce environments, bacteria have to use efficient mechanisms for the uptake of metals required for their growth. This is particularly the case for pathogenic bacteria that have to confront the host's immune system. Indeed, the so-called "nutritional immunity" induces an additional metal limitation by sequestering iron, zinc or manganese to prevent bacterial growth [1–4]. To face this metal restriction, bacteria have developed metallophores to recover metals. In this context, nicotianamine-like metallophores have been identified in some bacteria as playing an important role in metal acquisition strategies. Staphylopine, pseudopaline and yersinopine are the three examples

currently known and recently identified in *S. aureus* [5], *P. aeruginosa* [6,7] and *Y. pestis* [8] respectively. Depending on the species, the biosynthesis of these nicotianamine-like metallophores occurs in two or three steps. When it is present, such as in *S. aureus*, CntK, a histidine racemase, transforms L-histidine (L-His) into D-histidine (D-His). CntL, a nicotianamine synthase-like, adds an aminobutyrate moiety coming from S-adenosyl methionine (SAM) on the amino group of its second substrate (L-His in *P. aeruginosa* and *Y. pestis* or D-His in *S. aureus*) to form a pathway intermediate (noted yNA when using L-His, xNA when using D-His). Finally, CntM, an enzyme belonging to the opine dehydrogenase family, condenses the pathway intermediate with an α -ketoacid (pyruvate in the case of *S. aureus* and *Y. pestis* or α -ketoglutarate (α KG) in the case of *P. aeruginosa*) using NAD(P)H to form an opine-type metallophore (Figure 1).

Opine dehydrogenase catalyzes the NAD(P)H-dependent reductive condensation of the amino group of an amino acid with an α -ketoacid to produce an N-(carboxyalkyl) amino acid, also known as an opine, which exhibits either (L,L) or (D,L) stereochemistry [9]. The variety of amino acids and α -ketoacids that can be used as substrates by opine dehydrogenases results in diverse products. For example, the octopine dehydrogenase catalyzes the production of octopine, lysopine or histopine *via* reductive condensation of pyruvate and L-arginine, L-lysine and L-histidine respectively [10–13]. Leucinopine, asparaginopine and glutaminopine are other examples of opines produced *via* reductive condensation of α -ketoglutarate with L-leucine, L-asparagine and L-glutamine respectively [14,15]. In addition to the diversity of substrates, opine dehydrogenases are also distinguished by their biological roles. Indeed, in some marine invertebrates, opine dehydrogenases participate in the anaerobic metabolism by insuring the last step of anaerobic glycolysis pathway therefore participating in the propelling of these animals [16,17]. In plants, diverse opines are found inside crown gall tumors (for example nopaline, agropine, octopine, mannopine or D-L and L-L-succinamopine) that are induced by plant pathogenic bacteria as *Agrobacterium tumefaciens* [18,19]. In this case, the opines serve as nutrients conferring selective growth advantages to the opine-producing and opine-utilizing microorganisms [20–22].

Opine dehydrogenases are also involved in the biosynthesis of nicotianamine-like metallophores in bacteria and their substrate specificity results in the production of diverse metallophores. The activity of these opine dehydrogenases from *S. aureus*, *P. aeruginosa* and *Y. pestis* (respectively called SaCntM, PaCntM and YpCntM) has been described: SaCntM uses NADPH and pyruvate with xNA to produce staphylopin [5], YpCntM uses pyruvate and NADPH with yNA to produce yersinopine [8], and PaCntM uses NAD(P)H and α -ketoglutarate with yNA to produce pseudopaline [6,7]. Therefore, substrate specificity (and eventually stereospecificity in the case of xNA vs yNA) of CntM leads to the production of diverse opine-type metallophores. These metallophores are involved in metal acquisition, with metal specificity depending on the growth medium. Staphylopin could transport copper, nickel, cobalt, zinc and iron [5,23,24], and participates in zinc uptake in zinc-scarce environments [23,25]. Pseudopaline is involved in nickel uptake in minimal media, whereas it is responsible for zinc uptake in zinc-scarce environment [6,26]. For both *S. aureus* and *P. aeruginosa*, literature reports a link between the production of opine-type metallophores and bacterial infection. For example, in *S. aureus*, the staphylopin receptor CntA plays an important role in the optimal functioning of the urease activity and is ultimately linked to its virulence as the deletion of the gene encoding this substrate binding protein leads to a decrease of murine bacteremia and urinary tract infections [24]. In *P. aeruginosa*, transcriptomic analyses showed that the biosynthetic genes for pseudopaline are overexpressed in burn wound infections in humans [27]. This overexpression would make it possible to bypass the metal limitations put in place during nutritional immunity. Moreover, the pseudopaline exporter CntI plays an important role in the survival and growth of *P. aeruginosa* in cystic fibrosis airway: deletion of the gene encoding this exporter resulted in an attenuation of this respiratory infection [28]. Similarly, the exporter of staphylopin was

found to be important for fitness in abscesses, even before staphylopyrrolidine discovery [29]. Concerning *Y. pestis*, no data indicates a link between yersinopyrrolidine production and virulence at this time and the discovery of this metallophore is so far restricted to *in vitro* studies [8].

In an effort to understand bacterial opine-type metallophore diversity, we studied substrate specificity of CntM in *S. aureus* and *P. aeruginosa* by combining bioinformatic and structural analyses with chemical synthesis and enzymatic studies. We found that a single amino acid residue was responsible for the preferential binding of NADPH, and a single residue was involved in the specificity towards pyruvate or α -ketoglutarate. This finding led us to establish simple rules involved in opine-type metallophore diversity and prompted us to dig into available genomes for a bacteria possessing a *cntK* homologue together with a *cntM* gene predicted to use α -ketoglutarate. This exploration ultimately led to the discovery of a new opine-type metallophore called bacillopaline in *Paenibacillus mucilaginosus*.

MATERIALS AND METHODS

Bioinformatic analyses

SaCntM (*sav2468*) protein sequence was analyzed by searching for homologues using Psi-BLAST search [30] through the NCBI databases (National Center for Biotechnology Information, Bethesda, Maryland, USA) and the Pfam database (European Bioinformatics Institute, Hinxton, England, UK). Sequence alignment was done using the Muscle program [31] with defaults criteria from Jalview (version 2.10.3) [32]. Residues were colored following the Clustal coloring scheme with intensity modified by a conservation color increment of 30%. Gene synteny were inspected using the MaGe MicroScope web interface [33] added to sequence alignment analysis.

Cloning, expression and purification of proteins

The gene encoding the SaCntM protein were cloned in pET-SUMO and pET-101 [5,34] and the one encoding the PaCntM protein were cloned in pET-TEV [6]. Similarly, the genes encoding the PmCntL and PmCntM proteins were cloned in pET-TEV from genomic DNA. The primers used for these constructions are listed in Table S1. After co-transformation with plasmid pRARE (encoding for rare codon in *E. coli*), *E. coli* BL21 strains were aerobically cultivated with horizontal shaking in LB media supplemented with appropriate antibiotics (kanamycin at 50 $\mu\text{g.mL}^{-1}$ for pET-SUMO and pET-TEV, ampicillin at 50 $\mu\text{g.mL}^{-1}$ for pET-101 and chloramphenicol at 25 $\mu\text{g.mL}^{-1}$ for pRARE). These strains were grown in diverse conditions (37°C or 16°C, with or without induction of protein expression by addition of 0.1 mM IPTG when the OD of the culture was about 0.6; Table S2). After overnight growth, cells were recovered by centrifugation at 5,000 g for 20 min at 4°C. Cells were resuspended in buffer A and disrupted using a Constant cell disruption system operating at 1.9 Kbar. Cell debris were removed by centrifugation at 8,000 g for 20 min. The supernatant was centrifuged at 100,000 g for 45 min at 4°C to remove cell wall debris and membrane proteins. The resulting soluble fraction was purified by batch using a nickel-charged affinity resin (Ni-NTA Agarose resin, ThermoFisher Scientific). The proteins were eluted stepwise with imidazole (15 mM wash, 250 mM or 500 mM elution). Collected fractions were transferred into imidazole-free buffer B (see Table S2 for details on the buffers used and Figure S8 for the monitoring of protein purifications).

Site directed mutagenesis

Site directed mutagenesis were performed according to standard protocol from the QuickChange II Site-Directed Mutagenesis kit (Agilent Technologies). The only difference was that *E. coli* DH10 β Competent Cells were used instead of XL1-Blue Supercompetent Cells for

transformations. Selection of colonies was done after spreading on LB plate supplemented with appropriate antibiotics. Plasmid pET-SUMO containing the gene encoding the SaCntM protein and plasmid pET-TEV containing the gene encoding the PaCntM protein were used as mutagenesis templates for the D150A and A153D substitutions respectively. For the R33H substitution from SaCntM, mutagenesis was performed from plasmid pET-101 containing the gene encoding the SaCntM protein as template. Primer pairs were designed for single substitutions and were then synthesized by Eurofins Genomics (Table S1).

Chemical synthesis of xNA and yNA

The chemical synthesis of xNA has recently been described [34]. The same strategy was used for the chemical synthesis of yNA, although L-His-OMe was used instead of D-His-OMe as starting material. Purified intermediates and their characterization are described in the supplementary materials.

CntM activity assay

Enzymatic reactions of CntM were performed at 28°C in microplates and with purified proteins. They were carried out in a reaction volume of 100 µL in buffer C (50 mM BisTrisPropane, 100 mM NaCl, pH = 7 for SaCntM and PmCntM or pH = 8 for PaCntM) containing 2 or 5 µg of enzyme, 0.2 mM of NADH or NADPH (Sigma-Aldrich), 0.2 mM of xNA or yNA (chemically synthesized), and 1 mM of pyruvate or α-ketoglutarate (Sigma-Aldrich), unless otherwise stated. The absorbance at 340 nm was measured using a microplate reader (Infinite 200 Pro; Tecan) to follow the oxidation of NADPH illustrating the progress of the reaction. Activities were calculated from the initial rate and the amount of enzyme used. Kinetic parameters were estimated according to the Michaelis-Menten kinetic without **(1)** or with the substrate inhibition model **(2)** using SigmaPlot. These values were used to plot a fit on the experimental data. (V_m : Maximum velocity; K_m : Michaelis constant; $[S]$: substrate concentration; K_i : Inhibition constant).

$$(1): v = \frac{v_m [S]}{K_m + [S]}$$

$$(2): v = \frac{v_m [S]}{K_m + [S](1 + \frac{[S]}{K_i})}$$

Fluorescence resonance energy transfer (FRET) studies

Fluorescence studies were performed using a Cary Eclipse spectrophotometer (Agilent). The FRET experiment was done using a protein concentration of 5 µM and an excitation wavelength at 280 nm (tryptophan excitation). The emission at 340nm was transferred to the NAD(P)H and the signal was recorded between 400 and 500 nm. Five spectra were averaged in order to increase signal to noise ratio.

Activity assay followed by TLC

The assay consisted in incubating the purified enzymes at a final concentration of 2.5 µM and using carboxyl-¹⁴C-labeled SAM (2.5 µM), NADPH or NADH (30 µM), L- or D-histidine (10 µM) and α-keto acid (pyruvate or α-ketoglutarate; 1mM). The total volume was 100 µL in buffer D (50 mM of Hepes, 1 mM of DTT, 1mM of EDTA, pH = 9). The mixtures were incubated for 30 min at 28 °C. The reactions were stopped by adding ethanol to a final concentration of 50 % (v/v) and the products were then separated by thin layer chromatography. An aliquot of 10 µL of the reaction mixtures were spotted on HPTLC (High Performance TLC) Silica Gel 60 Glass Plates (Merck KGaA), and the plates were developed with a phenol:n-butanol:formate:water (12:3:2:3 v/v) solvent system. These separation

parameters were used in the initial biochemical characterization of plant nicotianamine synthase [35] and of staphylopine and pseudopaline [5,6]. HPTLC plates were dried and exposed to a [¹⁴C]-sensitive imaging plate for one day. Imaging plates were then scanned on a Typhoon FLA 7000 phosphorimager (GE Healthcare).

Cell culture conditions of *Paenibacillus mucilaginosus*

Paenibacillus mucilaginosus, sub sp. 1480D (from DSMZ collection) was grown by inoculating stock bacteria into 20 mL sterile medium (TSB 1/10) at 30°C for 48h. Genomic DNA was extracted using the protocols and pretreatments for Gram-positive bacteria from DNeasy Blood and Tissue kit (Qiagen).

RESULTS AND DISCUSSION

Specificity towards NADH or NADPH

The structure of CntM has been solved in a binary/tertiary complex with NADPH or NADPH and xNA, revealing the residues that are involved in the complex formation and building the active site [8,34] (pdb code 6C4T and 6H3D, respectively). Focusing on the NADPH binding site of SaCntM we observed that the phosphate group of NADPH is sandwiched by the side chains of two positively charged residues (R33 and K39) that are rather conserved in the CntM family (Figure 2A). However, a sequence alignment of nine CntM homologues shows that R33 residue in SaCntM is conserved in YpCntM, but replaced by a histidine in PaCntM (Figure 2B). As it is known that SaCntM and YpCntM specifically use NADPH [5,8] while PaCntM can use NADH or NADPH [6–8], we hypothesized that the nature of the amino acid residue at this position could determine the NADH/NADPH selectivity by CntM.

We therefore sought to determine the role of this residue in NAD(P)H selectivity by replacing this arginine by a histidine in a SaCntM:R33H variant. Because the histidine could either be neutral or positively charged at pH above or below its pK_a, the binding of NAD(P)H was followed at two pHs (6.0 and 8.5). We found that the WT enzyme, whatever the pH, still preferentially binds NADPH over NADH (Figure 3). On the contrary, the R33H mutant behaves as the WT at pH 6.0, whereas it loses its preferential binding property at a higher pH. This shows that a histidine at this position could serve as a selective residue, stabilizing NADPH at acidic pH when the imidazole ring of histidine is positively charged, and favoring NADH binding when histidine is neutral at basic pH, overall explaining the difference of selectivity in the literature [6–8]. In other words, at acidic pH the histidine side chain of the R33H mutant is charged and behaves like the WT, whereas at basic pH this histidine is neutral, it could not form a salt bridge with the phosphate of NADPH and therefore could not discriminate between NADPH and NADH. Interestingly, we noted that *Fictibacillus arsenicus* possesses a histidine residue at the conserved position equivalent to K39 in *S. aureus*, suggesting the same possibility of preferential NADH binding at basic pH.

Specificity towards xNA or yNA

In vivo, CntM uses the product of CntL *i.e.* xNA or yNA depending on the species. However, using enzymatically produced xNA or yNA, McFarlane *et al.* (2018) suggested that SaCntM could use both diastereoisomers. Activities from *S. aureus* (SaCntM) and from *P. aeruginosa* (PaCntM) were therefore compared for their ability to use chemically synthesized xNA and yNA as substrate (Figure 4). The chemical synthesis of xNA was recently reported [34] and we were able to synthesize yNA by following the same approaches (see the experimental procedures and the supplementary materials). Reactions were then performed *in vitro* using purified proteins and a concentration range of xNA and yNA with a saturating concentration of other substrates: 0.2 mM of NADPH and 1 mM of pyruvate

(when evaluating SaCntM) or α -ketoglutarate (when evaluating PaCntM). Overall, we found that the activity of CntM from *S. aureus* and *P. aeruginosa* towards yNA and xNA were different. In the case of SaCntM, although xNA is used *in vivo* to produce staphylopine, its activity is higher when using yNA *in vitro*. Indeed, the k_{cat} for the reaction with yNA is ~ 2 -fold higher than the one with xNA (3.18 s^{-1} and 1.67 s^{-1} respectively). However, the K_{m} for the reaction with yNA is ~ 2 -fold higher than the one with xNA ($47 \text{ }\mu\text{M}$ and $23 \text{ }\mu\text{M}$ respectively), leading to a catalytic efficiency ($k_{\text{cat}}/K_{\text{m}}$) of the same order of magnitude for the two substrates (Table 1). In the case of PaCntM, the enzyme uses yNA to produce pseudopaline *in vivo*. In agreement with this, we found that the reaction with yNA is more efficient than the one with xNA, the reaction with yNA exhibiting a catalytic efficiency ~ 10 -fold higher than with xNA ($29\,618 \text{ M}^{-1}\text{s}^{-1}$ and $2\,918 \text{ M}^{-1}\text{s}^{-1}$ respectively).

In the past decades, several opine dehydrogenases have been studied and their biosynthetic reactions generally exhibit a substrate stereospecificity towards the amino group with most enzymes using the L-stereoisomer as substrate [9]. For example, substrate stereospecificity has been outlined for the octopine dehydrogenase from *Pecten maximus* [12] and the structure of this enzyme shows a negatively charged cavity acting as a “charge ruler”, which favors L-arginine binding. Here, we found that even if SaCntM uses xNA *in vivo*, it is also capable of using yNA *in vitro*. This trend confirms the one outlined by McFarlane *et al.* [8] using enzymes from different species. Under their experimental conditions, PaCntM exhibited a k_{cat} of 0.016 s^{-1} in the presence of xNA, which was biosynthesized with SaCntL. Accordingly, we found that PaCntM preferentially used yNA whether *in vitro* or *in vivo*. All these data therefore suggest that there is a substrate stereospecificity in the case of PaCntM, which is not found in SaCntM. We further noted that, using yNA but not xNA, a drop in enzyme activity is visible at high substrate concentrations, which suggests a mechanism of substrate inhibition (Figure 4). Indeed, the fits made with a substrate inhibition model added to the Michaelis-Menten kinetic better cover the experimental data (Figure 4; plain lines). However, this substrate inhibition is not visible when using xNA as substrate.

Specificity towards pyruvate or α -ketoglutarate

In order to find amino acid residues involved in the pyruvate/ α -ketoglutarate selectivity by CntM, we searched for residues located in the vicinity of the active site (*i.e.* the nicotinamide moiety) and conserved in species known to use pyruvate (*S. aureus* and *Y. pestis*) but differing in species known to use α -ketoglutarate (*P. aeruginosa*). This pointed to the aspartic acid residue at position 150 in SaCntM, which is also present in YpCntM but replaced by an alanine in PaCntM (corresponding to residue 153 in this enzyme). Because this residue is either an aspartic acid or an alanine, we hypothesized that the nature of the amino acid residue at this position would determine the pyruvate/ α -ketoglutarate selectivity by CntM. This postulate was also proposed by McFarlane *et al.* [8]. In order to test the role of this amino acid in pyruvate/ α -ketoglutarate selectivity, we replaced the aspartic acid by an alanine in SaCntM (D150A variant) and did the opposite mutation in PaCntM (A153D). The activities of these proteins were then compared for their ability to use of pyruvate and α -ketoglutarate as substrates (Figure 5 and Figure S8 for the monitoring of protein purifications). The reactions were performed using a concentration range of pyruvate and α -ketoglutarate with a fixed concentration of others substrates: 0.2 mM of NAD(P)H and 0.2 mM of xNA (when using SaCntM) or yNA (PaCntM). As a control, we verified that the mutation did not affect the binding of the NADPH on SaCntM (Figure S9).

With regard to SaCntM, we confirmed that the WT was only able to use pyruvate, with a k_{cat} of 1.0 s^{-1} and a K_{m} of $51 \text{ }\mu\text{M}$, resulting in a catalytic efficiency of $19\,592 \text{ M}^{-1}\text{s}^{-1}$ (Figure 5 and Table 2). On the contrary, the D150A variant of SaCntM could use both pyruvate and α -ketoglutarate as substrates. Indeed, although the maximum activity is not reached within the concentration range tested, the kinetic parameters calculated for both pyruvate and α -ketoglutarate are in the same order of magnitude. This single substitution therefore led to a decreased activity when using pyruvate but most

of all, significantly led to an increased activity when using α -ketoglutarate. Even if we take a lower limit for the k_{cat} of 1.65 s^{-1} for α -ketoglutarate, this would correspond to a more than 40-fold increased as compared to the WT SaCntM. We then investigated whether the opposite mutation in PaCntM would trigger the opposite effect on substrate selectivity. Here again, we confirmed that the WT enzyme could only use α -ketoglutarate with a k_{cat} of 0.27 s^{-1} and a K_{m} of $133 \mu\text{M}$. Strikingly, we found that the A153D substitution in PaCntM led to a complete reversal of selectivity, with the variant only being able to use pyruvate and unable to use α -ketoglutarate anymore. This mutation is indeed accompanied by a more than 20-fold increase in k_{cat} for pyruvate and ~ 30 -fold decrease for α -ketoglutarate, *i.e.* a complete switch in substrate specificity. Consequently, substrate specificity of CntM towards pyruvate or α -ketoglutarate is mainly governed by this single amino acid (position 150 in SaCntM or 153 in PaCntM): an aspartate ensures the selection of pyruvate whereas an alanine leads to α -ketoglutarate specificity. To our knowledge, this is the first example showing that substrate specificity might be tuned in opine/opaline dehydrogenases family. There are however some examples of redesigned substrate specificity in dehydrogenases such as the production of a highly active malate dehydrogenase starting from lactate dehydrogenase [36]. In this case, the Gln102Arg mutation of *Bacillus stearothermophilus* lactate dehydrogenase led to a shift in $k_{\text{cat}}/K_{\text{m}}$ with malate so that it equal that of native lactate dehydrogenase for its natural substrate. Examples of site directed mutagenesis studies in the opine/opaline family were centered on the catalytic residues and all led to decreased activities towards substrates and none explored the putative α -ketoacid specificity [12,37]. Finally, there is also an example of opine binding proteins in which a single residue is key to the recognition of pyruvate vs α KG opines (*ie* octopines vs nopaline) [38].

Identification of a novel nicotianamine-like metallophore

Having established the molecular determinant for the α -ketoacid selectivity of CntM, we propose two simple rules governing the production of nicotianamine-like metallophores: 1) the presence or absence of a *cntK* homologue, which may or may not lead to the production of D-His from L-His respectively (then used by CntL to produce xNA or yNA respectively), and 2) the presence of an aspartate or an alanine at position 150 (*S. aureus* numbering), which results in pyruvate or α -ketoglutarate incorporation, respectively. Applying these rules, we searched for a species capable of producing the missing metallophore variant, which would use xNA and incorporate α -ketoglutarate, *i.e.* a species possessing a *cntK* homologue and an alanine at position responsible for α -ketoglutarate selectivity. Digging into available genomes *in silico*, we identified *Paenibacillus mucilaginosus* as a good candidate. Indeed, this species carries an A153 in CntM (as *P. aeruginosa*) and possess a *cntK* in its *cnt* operon (as *S. aureus*) (Figure 2B-C).

To check the validity of this hypothesis, genes encoding the CntL and CntM from *P. mucilaginosus* (PmCntL and PmCntM respectively) were amplified from genomic DNA, cloned in expression vectors, purified and used to determine the substrates they consumed *in vitro* (Figure 6). Enzymes activities were first assayed using TLC separation, using carboxyl- $[^{14}\text{C}]$ -labeled SAM substrate. In this assay, $[^{14}\text{C}]$ -labeled SAM shows a characteristic profile with one strong band and two others bands of much lower intensity (Figure 6 A). The incubation of $[^{14}\text{C}]$ -labeled SAM with PmCntL led to another prominent band in the presence of D-His, which was not found using L-His. This novel band migrated below the SAM band and therefore corresponds to the xNA intermediate. This indicates that PmCntL uses D-His and not L-His, and further validates the link between the presence of a *cntK* gene and the use of D-His by CntL. We then tested the substrate specificity of PmCntM by co-incubating both PmCntL and PmCntM together with diverse substrates: pyruvate or α -ketoglutarate with NADH or NADPH. When using NADPH and D-His with either pyruvate or α -ketoglutarate, we neither detected the SAM nor xNA pattern, but found a novel band migrating just above the SAM in the presence of pyruvate (which corresponds to staphylopin) and just below the SAM in the presence of α -ketoglutarate. Surprisingly, we thus found that PmCntM could use both pyruvate and α -ketoglutarate in this TLC assay. However, TLC experiments were run using a defined incubation time

(30 min), which might hinder differences in enzyme efficiency. We therefore determined the enzymatic parameters of PmCntM using purified protein and a concentration range of pyruvate and α -ketoglutarate, with a fixed concentration of others substrates: 0.2 mM of NADPH and 0.2 mM of xNA (Figure 6 B). As a result, we found that the catalytic efficiency is 10-fold better for α -ketoglutarate than for pyruvate ($10\,367\text{ M}^{-1}\text{s}^{-1}$ and $1\,016\text{ M}^{-1}\text{s}^{-1}$ respectively; Table 3). These data therefore suggest that our hypothesis is valid: CntL and CntM from *P. mucilaginosus* produce an additional variant of opine-type metallophore with D-His, NADPH and α -ketoglutarate. This metallophore has been called bacillopaline as it belongs to the opaline family (Figure 6 C). Moreover, we found that, in addition to using both pyruvate and α -ketoglutarate, CntM from *P. mucilaginosus* is also able to use both xNA and yNA (Figure S10, Table S3).

Contrary to human pathogens like *S. aureus*, *P. aeruginosa* or *Y. pestis* in which opine-type metallophores were discovered [5–8], *P. mucilaginosus* is a soil bacteria. The production of opine-type metallophores was shown to be regulated by zinc through the zur (zinc uptake repressor) repressor [6,25], which is likely the case in *P. mucilaginosus*, with a putative zur box upstream the *cnt* operon. A possible hypothesis would be that bacillopaline production could be induced in calcareous soils, where zinc bioavailability has been shown to be low [39]. In the past decade, *P. mucilaginosus* have been studied for its capacity in wastewater treatment, but also as biofertilizer and plant growth promoting rhizobacteria [40–42]. Indeed, in addition to producing biofloculants or plant hormones, this species is able to solubilize phosphates, to fix nitrogen and to produce ammonia. Moreover, it is able to produce siderophores, which contributes to plant growth by indirectly preventing the growth of plant pathogens [41,43]. Similarly, we could hypothesize that bacillopaline could contribute in the same way to maintain plant micronutrients homeostasis.

In conclusion, studying the substrate specificity of the enzyme catalyzing the last step of nicotianamine-like metallophore biosynthesis allowed us to determine simple rules governing their production. First, the presence or absence of a *cntK* gene leads to the use of respectively D-His or L-His by CntL resulting in the incorporation of respectively xNA or yNA by CntM. Secondly, the presence of an aspartate or an alanine at position 150 on CntM (*S. aureus* numbering) results in pyruvate or α -ketoglutarate incorporation, respectively. Thanks to these simple rules, it is now possible to predict the nature of the nicotianamine-like metallophore produced by all bacteria possessing a *cnt* operon in their genome.

Acknowledgments

We thank the Agence Nationale de la Recherche (grant ANR-14-CE09-0007-02) for initial support and the association Vaincre la Mucoviscidose (VLM, grant RFI20160501495) for financial support.

Conflict of Interest

The authors declare that they have no conflicts of interest with the contents of this article.

Author contributions

C.L. and P.A. designed the experiments. C.L., C.B., C.H. and L.O. carried out the experiments. C.L., C.B. and P.A. analyzed the data. G.C., R.F synthesized xNA and yNA under the supervision of F.C. C.L. and P.A. wrote the manuscript with contributions from F.C.

References

- 1 Capdevila, D. A., Wang, J. and Giedroc, D. P. (2016) Bacterial Strategies to Maintain Zinc Metallostasis at the Host-Pathogen Interface. *J. Biol. Chem.* **291**, 20858–20868.

- 2 Hood, M. I. and Skaar, E. P. (2012) Nutritional immunity: transition metals at the pathogen–host interface. *Nat. Rev. Microbiol.* **10**, 525–537.
- 3 Weinberg, E. D. (1975) Nutritional immunity. Host’s attempt to withhold iron from microbial invaders. *JAMA* **231**, 39–41.
- 4 Zygiel, E. M. and Nolan, E. M. (2018) Transition Metal Sequestration by the Host-Defense Protein Calprotectin. *Annu. Rev. Biochem.* **87**, 621–643.
- 5 Ghssein, G., Brutesco, C., Ouerdane, L., Fojcik, C., Izaute, A., Wang, S., Hajjar, C., Lobinski, R., Lemaire, D., Richaud, P., et al. (2016) Biosynthesis of a broad-spectrum nicotianamine-like metallophore in *Staphylococcus aureus*. *Science* **352**, 1105–1109.
- 6 Lhospice, S., Gomez, N. O., Ouerdane, L., Brutesco, C., Ghssein, G., Hajjar, C., Liratni, A., Wang, S., Richaud, P., Bleves, S., et al. (2017) *Pseudomonas aeruginosa* zinc uptake in chelating environment is primarily mediated by the metallophore pseudopaline. *Sci. Rep.* **7**.
- 7 McFarlane, J. S. and Lamb, A. L. (2017) Biosynthesis of an Opine Metallophore by *Pseudomonas aeruginosa*. *Biochemistry* **56**, 5967–5971.
- 8 McFarlane, J. S., Davis, C. L. and Lamb, A. L. (2018) Staphylopine, pseudopaline, and yersinopine dehydrogenases: A structural and kinetic analysis of a new functional class of opine dehydrogenase. *J. Biol. Chem.* **293**, 8009–8019.
- 9 Thompson, J. and Donkersloot, J. A. (1992) N-(Carboxyalkyl)Amino Acids: Occurrence, Synthesis, and Functions. *Annu. Rev. Biochem.* **61**, 517–57.
- 10 Bates, H. A., Kaushal, A., Deng, P. N. and Sciaky, D. (1984) Structure and synthesis of histopine, a histidine derivative produced by crown gall tumors. *Biochemistry* **23**, 3287–3290.
- 11 Biemann, K., Lioret, C., Asselineau, J., Lederer, E. and Polonsky, J. (1960) On the structure of lysopine, a new amino acid isolated from crown gall tissue. *Biochim. Biophys. Acta* **40**, 369–370.
- 12 Smits, S. H. J., Mueller, A., Schmitt, L. and Grieshaber, M. K. (2008) A Structural Basis for Substrate Selectivity and Stereoselectivity in Octopine Dehydrogenase from *Pecten maximus*. *J. Mol. Biol.* **381**, 200–211.
- 13 Van Thoai, N., Huc, C., Pho, D. B. and Olomucki, A. (1969) Octopine déshydrogénase : Purification et Propriétés Catalytiques. *Biochim. Biophys. Acta BBA - Enzymol.* **191**, 46–57.
- 14 Chang, C.-C. and Chen, C.-M. (1983) Evidence for the presence of N^2 -(1,3-dicarboxypropyl)-L-amino acids in crown-gall tumors induced by *Agrobacterium tumefaciens* strains 181 and EU6. *FEBS Lett.* **162**, 432–435.
- 15 Chang, C.-C., Chen, C.-M., Adams, B. R. and Trost, B. M. (1983) Leucinopine, a characteristic compound of some crown-gall tumors. *Proc. Natl. Acad. Sci. USA* **80**, 3573–3576.
- 16 Gäde, G., Weeda, E. and Gabbott, P. A. (1978) Changes in the Level of Octopine during the Escape Responses of the Scallop, *Pecten maximus* (L.). *J. Comp. Physiol. B* **124**, 121–127.
- 17 Harcet, M., Perina, D. and Pleše, B. (2013) Opine Dehydrogenases in Marine Invertebrates. *Biochem. Genet.* **51**, 666–676.
- 18 Moore, L. W., Chilton, W. S. and Canfield, M. L. (1997) Diversity of Opines and Opine-Catabolizing Bacteria Isolated from Naturally Occurring Crown Gall Tumors. *Appl. Environ. Microbiol.* **63**, 201–207.
- 19 Dessaux, Y. and Faure, D. (2018) Niche Construction and Exploitation by *Agrobacterium*: How to Survive and Face Competition in Soil and Plant Habitats. In *Agrobacterium Biology: From Basic Science to Biotechnology* (Gelvin, S. B., ed.), pp 55–86, Springer International Publishing, Cham.
- 20 Tremblay, G., Gagliardo, R., Chilton, W. S. and Dion, P. (1987) Diversity among Opine-Utilizing Bacteria: Identification of Coryneform Isolates. *Appl. Environ. Microbiol.* **53**, 1519–1524.
- 21 Lang, J., Vigouroux, A., Planamente, S., Sahili, A. E., Blin, P., Aumont-Nicaise, M., Dessaux, Y., Moréra, S. and Faure, D. (2014) *Agrobacterium* Uses a Unique Ligand-Binding Mode for Trapping Opines and Acquiring A Competitive Advantage in the Niche Construction on Plant Host. *PLOS Pathog.* **10**, e1004444.
- 22 Montoya, A. L. (1977) Octopine and Nopaline Metabolism in *Agrobacterium tumefaciens* and Crown Gall Tumor Cells: Role of Plasmid Genes. *J. Bacteriol.* **129**, 101–107.

- 23 Grim, K. P., San Francisco, B., Radin, J. N., Brazel, E. B., Kelliher, J. L., Párraga Solórzano, P. K., Kim, P. C., McDevitt, C. A. and Kehl-Fie, T. E. (2017) The Metallophore Staphylopin Enables *Staphylococcus aureus* To Compete with the Host for Zinc and Overcome Nutritional Immunity. *mBio* (Torres, V. J., ed.) **8**.
- 24 Remy, L., Carrière, M., Derré-Bobillot, A., Martini, C., Sanguinetti, M. and Borezée-Durant, E. (2013) The *Staphylococcus aureus* Opp1 ABC transporter imports nickel and cobalt in zinc-depleted conditions and contributes to virulence: Nickel and cobalt uptake in *Staphylococcus aureus*. *Mol. Microbiol.* **87**, 730–743.
- 25 Fojcik, C., Arnoux, P., Ouerdane, L., Aigle, M., Alfonsi, L. and Borezée-Durant, E. (2018) Independent and cooperative regulation of staphylopin biosynthesis and trafficking by Fur and Zur: Regulation of *S. aureus* *cnt* operon by Fur and Zur. *Mol. Microbiol.* **108**, 159–177.
- 26 Mastropasqua, M. C., D’Orazio, M., Cerasi, M., Pacello, F., Gismondi, A., Canini, A., Canuti, L., Consalvo, A., Ciavardelli, D., Chirullo, B., et al. (2017) Growth of *Pseudomonas aeruginosa* in zinc poor environments is promoted by a nicotianamine-related metallophore: Metallophore-mediated zinc uptake in *Pseudomonas aeruginosa*. *Mol. Microbiol.* **106**, 543–561.
- 27 Bielecki, P., Puchałka, J., Wos-Oxley, M. L., Loessner, H., Glik, J., Kawecki, M., Nowak, M., Tümmler, B., Weiss, S. and dos Santos, V. A. P. M. (2011) In-Vivo Expression Profiling of *Pseudomonas aeruginosa* Infections Reveals Niche-Specific and Strain-Independent Transcriptional Programs. *PLoS ONE* (Brown, S. P., ed.) **6**, e24235.
- 28 Gi, M., Lee, K.-M., Kim, S. C., Yoon, J.-H., Yoon, S. S. and Choi, J. Y. (2015) A novel siderophore system is essential for the growth of *Pseudomonas aeruginosa* in airway mucus. *Sci. Rep.* **5**, 14644.
- 29 Ding, Y., Fu, Y., Lee, J. C. and Hooper, D. C. (2012) *Staphylococcus aureus* NorD, a Putative Efflux Pump Coregulated with the Opp1 Oligopeptide Permease, Contributes Selectively to Fitness *In Vivo*. *J. Bacteriol.* **194**, 6586–6593.
- 30 Altschul, S. F., Madden, T. L., Schäffer, A. A., Zhang, J., Zhang, Z., Miller, W. and Lipman, D. J. (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* **25**, 3389–3402.
- 31 Edgar, R. C. (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.* **32**, 1792–1797.
- 32 Clamp, M., Cuff, J., Searle, S. M. and Barton, G. J. (2004) The Jalview Java alignment editor. *Bioinformatics* **20**, 426–427.
- 33 Vallenet, D., Belda, E., Calteau, A., Cruveiller, S., Engelen, S., Lajus, A., Le Fèvre, F., Longin, C., Mornico, D., Roche, D., et al. (2013) MicroScope—an integrated microbial resource for the curation and comparative analysis of genomic and metabolic data. *Nucleic Acids Res.* **41**, D636–D647.
- 34 Hajjar, C., Fanelli, R., Laffont, C., Brutescio, C., Cullia, G., Tribout, M., Nurizzo, D., Borezée-Durant, E., Voulhoux, R., Pignol, D., et al. (2019) Control by Metals of Staphylopin Dehydrogenase Activity during Metallophore Biosynthesis. *J. Am. Chem. Soc.* **141**, 5555–5562.
- 35 Higuchi, K., Kanazawa, K., Nishizawa, N.-K. and Mori, S. (1996) The role of nicotianamine synthase in response to Fe nutrition status in Gramineae. *Plant Soil* **178**, 171–177.
- 36 Wilks, H. M., Hart, K. W., Feeney, R., Dunn, C. R., Muirhead, H., Chia, W. N., Barstow, D. A., Atkinson, T., Clarke, A. R. and Holbrook, J. J. (1988) A specific, highly active malate dehydrogenase by redesign of a lactate dehydrogenase framework. *Science* **242**, 1541–1544.
- 37 Müller, A., Janßen, F. and Grieshaber, M. K. (2007) Putative reaction mechanism of heterologously expressed octopine dehydrogenase from the great scallop, *Pecten maximus* (L). *FEBS J.* **274**, 6329–6339.
- 38 Vigouroux, A., El Sahili, A., Lang, J., Aumont-Nicaise, M., Dessaux, Y., Faure, D. and Moréra, S. (2017) Structural basis for high specificity of octopine binding in the plant pathogen *Agrobacterium tumefaciens*. *Sci. Rep.* **7**.
- 39 Alloway, B. J. (2009) Soil factors associated with zinc deficiency in crops and humans. *Environ. Geochem. Health* **31**, 537–548.

- 469 40 Deng, S., Bai, R., Hu, X. and Luo, Q. (2003) Characteristics of a bioflocculant produced by *Bacillus*
470 *mucilaginosus* and its use in starch wastewater treatment. Appl. Microbiol. Biotechnol. **60**, 588–
471 593.
- 472 41 Goswami, D., Parmar, S., Vaghela, H., Dhandhukia, P. and Thakker, J. N. (2015) Describing
473 *Paenibacillus mucilaginosus* strain N3 as an efficient plant growth promoting rhizobacteria
474 (PGPR). Cogent Food Agric. (Moral, M. T., ed.) **1**.
- 475 42 Liu, S., Tang, W., Yang, F., Meng, J., Chen, W. and Li, X. (2017) Influence of biochar application on
476 potassium-solubilizing *Bacillus mucilaginosus* as potential biofertilizer. Prep. Biochem.
477 Biotechnol. **47**, 32–37.
- 478 43 Haas, D. and Défago, G. (2005) Biological control of soil-borne pathogens by fluorescent
479 *pseudomonads*. Nat. Rev. Microbiol. **3**, 307–319.
- 480
- 481

Figures and figure legends




Figure 1: Differences in the staphylopine, pseudopaline, and yersinopine biosynthetic pathways. Adapted from [5,6,8].




Figure 2: CntM structure, sequence conservation around NADPH and operon diversity highlight protein selectivity. (A) Details of the NADPH binding site on SaCntM with side chains of residues located up to 4Å around the NADPH represented in stick. The protein is colored from N-terminus (blue) to C-terminus (red). The pdb code used for this figure corresponds to 6H3D and shows two alternative positions for the adenine moiety of NADPH. Dashed lines indicate hydrogen bonds and salt bridges. The nicotinamide moiety is where the reductive condensation of xNA with pyruvate occurs and is involved in a putative hydrogen bond with D150 side chain. (B) Sequence alignment of nine CntM protein sequences from bacteria. Threshold for the Clustal coloring scheme correspond to 30 % sequence conservation as defined in Jalview. Arrows point to residues involved in NADPH/NADH selectivity and pyruvate/ α -ketoglutarate selectivity. A weblogo representation of the entire unitprot family available from Pfam (318 entries) is drawn below the alignment, showing the conservation of position equivalent to R33 (R: 90%, H: 6% and K: 4%) and D150 (D: 72%, A: 25% and T: 3%) (C) Genomic organization of the biosynthetic genes of four different *cnt* operons in bacteria.




Figure 3: Titration of NADPH (black circles) and NADH (white circles) binding to SaCntM (5 μ M) at pH = 6.0 or pH = 8.5 followed by fluorescence energy transfer between tryptophan excitation (280 nm) and NAD(P)H emission (450 nm).




Figure 4: Activity profile of SaCntM and PaCntM using variable concentrations of xNA (black circle) and yNA (white circle) with fixed concentrations of other substrates: 0.2 mM of NADPH and 1 mM of pyruvate when evaluating SaCntM, or 0.2 mM of NADH and 1 mM of α -ketoglutarate when evaluating PaCntM. The data points are means of three replicates with standard deviations. The fits are made using the Michaelis-Menten model considering (continuous line) or not (dashed line) a substrate inhibition.

Protein	xNA vs yNA	K_m (μM)	K_i (mM)	k_{cat} (s^{-1})	k_{cat}/K_m ($\text{M}^{-1}\text{s}^{-1}$)
SaCntM	yNA	47 ± 9	2.3 ± 0.8	3.18 ± 0.25	67 584
SaCntM	xNA	23 ± 3	NA	1.67 ± 0.04	72 711
PaCntM	yNA	21 ± 7	2.4 ± 1.0	0.62 ± 0.06	29 618
PaCntM	xNA	54 ± 20	NA	0.16 ± 0.02	2 918

Table 1: Kinetic parameters of SaCntM and PaCntM activities established for a concentration range of xNA and yNA with fixed concentrations of other substrates: 0.2 mM of NADPH and 1 mM of pyruvate when evaluating SaCntM, and 0.2 mM of NADH and 1 mM of α -ketoglutarate when evaluating PaCntM. The data and the standard errors associated with were generated by SigmaPlot according to the Michaelis-Menten model with or without substrate inhibition. (NA: Not Applicable; V_m : Maximum velocity; K_m : Michaelis constant; K_i : Inhibition constant; k_{cat} : Catalytic constant (or turnover number); k_{cat}/K_m : Catalytic efficiency).




Figure 5: Activity profile of SaCntM (WT and D150A mutant) and PaCntM (WT and A153D mutant) using variable concentrations of pyruvate (black circle) and α -ketoglutarate (white circle) with fixed concentrations of others substrates: 0.2 mM of NADPH and xNA when evaluating SaCntM, and 0.2 mM of NADH and yNA when evaluating PaCntM. The data points are means of three replicates with standard deviations. The fits are made using the Michaelis-Menten model.

Protein	Pyruvate vs αKG	K_m (μM)	k_{cat} (s^{-1})	$k_{cat}/K_m(M^{-1}s^{-1})$
SaCntM	pyruvate	51 ± 6	1.00 ± 0.02	19 592
SaCntM	α KG	ND	0.04 ± 0.01	ND
SaCntM:D150A	pyruvate	$681 \pm 115^*$	$1.23 \pm 0.08^*$	1 807*
SaCntM:D150A	α KG	$2\,040 \pm 431^*$	$1.65 \pm 0.22^*$	809*
PaCntM	pyruvate	ND	0.02 ± 0.01	N.D.
PaCntM	α KG	133 ± 39	0.27 ± 0.02	2 058
PaCntM:A153D	pyruvate	$14\,986 \pm 2964^*$	$0.46 \pm 0.07^*$	31
PaCntM:A153D	α KG	ND	0.01 ± 0.01	ND

Table 2: Kinetic parameters of SaCntM (WT and D150A mutant) and PaCntM (WT or A153D mutant) activities established for a concentration range of pyruvate and α KG with fixed concentrations of other substrates: 0.2 mM of NADPH and xNA when evaluating SaCntM, and 0.2 mM of NADH and yNA when evaluating PaCntM. The data and the standard errors associated with were generated by SigmaPlot according to the Michaelis-Menten model. (ND: Not Determined). *Because the maximum enzyme activity is not reached, these values are not well defined and must be taken with caution.




Figure 6: Activity of CntL and CntM from *Paenibacillus mucilaginosus*. (A) TLC separation of reaction products incubating carboxyl- ^{14}C -labeled with purified enzyme (PmCntL and PmCntM) and various substrates (L-His (L) or D-His (D), pyruvate (Pyr) or α -ketoglutarate (α KG), NADH or NADPH). (B) Activity profile of PmCntM using variable concentrations of pyruvate (black circle) and α -ketoglutarate (white circle) with fixed concentrations of others substrates: 0.2 mM of NADPH and 0.2 mM of xNA. The data points are means of three replicates with standard deviations. The fits are made using the Michaelis-Menten model. (C) Biosynthetic pathway for the assembly of bacillopaline from D-His, SAM and α -ketoglutarate.

Protein	Pyruvate vs αKG	K_m (μM)	k_{cat} (s^{-1})	k_{cat}/K_m ($M^{-1}s^{-1}$)
PmCntM	pyruvate	180 ± 39	0.18 ± 0.01	1 016
PmCntM	α KG	42 ± 5	0.44 ± 0.01	10 367

556 Table 3: Kinetic parameters of PmCntM activities established for a concentration range of pyruvate
557 and α KG with fixed concentrations of others substrates: 0.2 mM of NADPH and 0.2 mM of xNA. The
558 data and the standard errors associated with were generated by SigmaPlot according to the Michaelis-
559 Menten model.