



New Catalytic Reactions using CO₂ for Radiolabeling of Tracers

Alexia Ohleier, Davide Audisio, Fabien Caillé, T. Cantat

► To cite this version:

Alexia Ohleier, Davide Audisio, Fabien Caillé, T. Cantat. New Catalytic Reactions using CO₂ for Radiolabeling of Tracers. Journées Annuelles de la Division de Chimie de Coordination, Jan 2019, Montpellier, France. cea-02328998

HAL Id: cea-02328998

<https://cea.hal.science/cea-02328998>

Submitted on 23 Oct 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.


New Catalytic Reactions using CO₂ for Radiolabeling of Tracers

Alexia Ohleier,^a Davide Audisio,^b Fabien Caillé^c, Thibault Cantat.^a

^a CEA Saclay - IRAMIS/NIMBE/LCMCE, 91191 Gif sur Yvette Cedex. ^b CEA Saclay - DRF/JOLIOT/SCBM/LMC, 91191 Gif sur Yvette Cedex. ^c Imagerie Moléculaire In Vivo UMR1023 CEA, INSERM, CNRS, Université Paris Sud, Université Paris-Saclay, Service Hospitalier Joliot, Orsay, France
e-mail : alexia.ohleier@cea.fr

CO₂, well known as a greenhouse gas responsible for global warming and considered as a cheap, non-toxic, renewable C1 building block, is by far the most accessible primary source of carbon for all radiolabeled compounds. However, its kinetic and thermodynamic stability significantly restrains potential applications. Broader availability of secondary labeled carbon sources would therefore grant access to a wide array of highly functionalized bioactive tracers. Indeed, ¹⁴C labelled compounds are commonly required during the development process of drugs whereas ¹¹C molecules are key radiotracers for positron emission tomography (PET) imaging.

In this context, we have been investigating CO₂ reduction into methoxy derivatives followed by a Suzuki cross-coupling to generate anisoles moieties. Selective borane mediated organocatalytic CO₂ reduction into OMe synthons has been previously disclosed by our group.^[1] However, very few couplings between methanol derivatives and aryl halides have been reported to date.^[2] Inspired by Novak's work,^[3] an innovative, efficient palladium catalyzed cross-coupling between methoxyboranes and aryl bromides has been developed under mild conditions. Working in a double chamber set-up allows both reactions to be compatible. This strategy is further expected to be tested on therapeutic targets and ideally transposed to ¹⁴C and ¹¹C isotopes.



Statut : Post-doctorante

- [1]. E. Blondiaux, J. Pouessel, T. Cantat, *Angew. Chem. Int. Ed.*, **2014**, 53, 12186 - 12190.
[2]. (a) K. E. Torraca, X. Huang, C. A. Parrish, S. L. Buchwald, *J. Am. Chem. Soc.*, **2001**, 123, 10770 - 10771. (b) E. J. Milton, J. A. Fuentes, M. L. Clarke, *Org. Biomol. Chem.*, **2009**, 7, 2645 - 2648. (c) S. Gowrisankar, H. Neumann, M. Beller, *Chem. Eur. J.*, **2012**, 18, 2498 - 2502.
[3]. G. L. Tolnai, B. Pethő, P. Králl, Z. Novák, *Adv. Synth. Catal.*, **2014**, 356, 125 - 129.