



HAL
open science

Sulfonylative Hiyama Cross-Coupling: Development and Mechanistic Insights

A Adenot, J Char, N von Wolff, G Lefèvre, T Cantat

► **To cite this version:**

A Adenot, J Char, N von Wolff, G Lefèvre, T Cantat. Sulfonylative Hiyama Cross-Coupling: Development and Mechanistic Insights. GECOM-CONCOORD 2019, May 2019, Erquy, France. cea-02329659

HAL Id: cea-02329659

<https://cea.hal.science/cea-02329659v1>

Submitted on 23 Oct 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Sulfonylative Hiyama Cross-Coupling: Development and Mechanistic Insights

A. Adenot,^a J. Char,^a N. von Wolff,^a G. Lefèvre,^a T. Cantat^{a,*}

^a: NIMBE, CEA, CNRS, Université Paris-Saclay, CEA Saclay, 91191 Gif-sur-Yvette cedex, France
aurelien.adenot@cea.fr

Due to distinctive structural and electronic features, sulfones have attracted a particular attention over the past few decades, making it a widespread functional group.^[1] Present in many contemporary pharmaceuticals and agrochemicals, they are also used as essential intermediates in organic synthesis. Therefore, numerous methodologies have been developed for their preparation.^[1] However, the most common methods suffer from significant limitations with harsh reaction conditions or regioselectivity issues. Recently, the insertion of a molecule of sulfur dioxide between two partners was investigated and reactions involving organomagnesium,^[2a] organozinc^[2b] and organoboron^[2c] compounds were reported. Herein we report a direct single-step palladium-catalyzed synthesis of sulfones involving organosilanes, sulfur dioxide and organohalides. Different mechanistic pathways were envisaged and discussed both from an experimental and theoretical standpoint.



[1] N.-W. Liu, S. Liang, G. Manolikakes, *Synthesis*, **2016**, 48, 1939.

[2] a) A. S. Deeming, C. J. Russel, A. J. Hennessy, M. C. Willis, *Org. Lett.*, **2014**, 16, 150; b) B. N. Rocke, K. B. Bahnck, M. Herr, S. Lavergne, V. Mascitti, C. Perreault, J. Polivkova, A. Shavnya, *Org. Lett.*, **2014**, 16, 154; c) Y. Chen, M. C. Willis, *Chem. Sci.*, **2017**, 8, 3249.

[3] N. von Wolff, J. Char, X. Frogneux, T. Cantat, *Angew. Chem. Int. Ed.* **2017**, 56, 5616.