



HAL
open science

Activity Coefficients from Vapor-Liquid Interfaces: A Molecular Dynamics Approach for Separation Chemistry

M. Bley, M. Duvail, Philippe Guilbaud, Jean-François Dufrêche

► To cite this version:

M. Bley, M. Duvail, Philippe Guilbaud, Jean-François Dufrêche. Activity Coefficients from Vapor-Liquid Interfaces: A Molecular Dynamics Approach for Separation Chemistry. Journées Scientifiques de Marcoule (JSM - 2018), Jun 2018, Bagnols Sur Ceze, France. cea-02338585

HAL Id: cea-02338585

<https://cea.hal.science/cea-02338585>


Submitted on 9 Dec 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Context


- Understanding **solvent extraction** – Recycling of rare earth elements and spent nuclear fuels



- The **equilibrium constant K^0** describes the extraction process and is given by the Mass Action Law




- A **multiscale approach** provides mesoscopic thermodynamic properties




Methods and concepts

- Measuring** activity and the vapor pressure




- Solvent flow:** Vapor pressure of the pure solvent always higher than of a mixture
- Ideal case: Solvent vapor pressure by **Raoult's law**
- Otherwise: **Aggregation increases, dissociation decreases** solvent vapor pressure and thus activity

p_1^* : Vapor pressure of pure solvent



Isoopiestic Measurements


- Simulating** liquid-vapor equilibria⁽¹⁾



- Molecular Dynamics using **explicit polarization**^(2,3) at 298.15 K for 15 ns

$$a_S = \frac{p}{p^*} = \frac{\rho}{\rho^*} = \frac{N \times V_{\text{gas}}^*}{N^* \times V_{\text{gas}}} = \frac{\lambda \times V_{\text{gas}}^*}{\lambda^* \times V_{\text{gas}}}$$

- Controlling** the error of the method



$$N = \langle \tau_x \rangle \times \lambda \quad \delta_N(t) = \frac{1}{\sqrt{\lambda \times t}}$$


λ : Evaporation rate $\langle \tau_x \rangle$: Time of flight

$\delta_N(t)$: Relative error of the method

a_S : Solvent activity


Aqueous dysprosium nitrate Dy(NO₃)₃ solutions (2)

Water activity a_W




$$a_W(b_E) = \frac{p_{\text{gas}}(b_E)}{p_{\text{gas}}^*} = \frac{\rho_{\text{gas}}(b_E)}{\rho_{\text{gas}}^*}$$

Osmotic coefficient ϕ_W



$$\phi_W(b_E) = \frac{-\ln a_W(b_E)}{\nu M_W b_E}$$

Activity coefficient γ_E




$$\ln \gamma_E(b_E) = \phi_W(b_E) - 1 + \int_{b=0}^{b=b_E} \frac{\phi_W(b) - 1}{b} db$$

Gibbs-Duhem relation

Organic phases – Binary Mixtures of DMDOHEMA and *n*-heptane



- Molecular dynamics simulation of liquid-vapor equilibria of organic solvent phases containing:



Pure solvent
+ Extractant
+ Water
+ Ionic species

Increasing computational cost

- Aggregation of DMDOHEMA in the organic phase relies on the availability of water⁽³⁾



$$a_{\alpha} = \gamma_{\alpha} \cdot x_{\alpha}$$

- Solvent activities in binary mixtures of *n*-heptane and DMDOHEMA show ideal behavior and follow Raoult's law in good agreement up to a high extractant concentration
- Attractive and repulsive interactions in the mixtures cancel out

Outlook and conclusion

- Aqueous salt solutions:** Results from molecular dynamics simulation are in good agreement with experimental data for different nitrate salts. This approach can be used for the validation of force fields for MD simulation with respect to thermodynamic properties.
- Organic Phases:** Method has been successfully applied on binary mixtures (*n*-heptane and DMDOHEMA, but also ethanol and water⁽⁴⁾). Allows accessing more complex organic solvent phases containing water and ionic species.

- Simulation of activity coefficients of **complex solutions with volatile solvents** for understanding solvent extraction

References

- Bley, M.; Duvail, M.; Guilbaud, P.; Dufrêche, J.-F., *J. Phys. Chem. B* **2017**, 121 (41), 9647-9658 (DOI: 10.1021/acs.jpcc.7b04011)
- Rard, J. A.; Spedding, F. H.; *J. Chem. Eng. Data* **1982**, 26, 391-395
- Meridiano, Y.; Berthon, L.; Crozes, X.; Dannus, P.; Antonio, M. R.; Chiarizia, R.; Zemb, T.; *Solvent Extr. Ion Exchange* **2009**, 27 (5-6), 607-637
- Bley, M.; Duvail, M.; Guilbaud, P.; Penisson, C.; Theisen, J.; Gabriel, J.-C.; Dufrêche, J.-F.; *Mol. Phys.* **2018** (DOI: 10.1080/00268976.2018.1444209)

Acknowledgements

This work was made possible thanks to the high performance computing facilities of TGCC/CCRT and the computing center of CEA Marcoule.