



HAL
open science

Understanding the catalytic carbonylation of epoxides for the valorization of carbon monoxide

Louise Ponsard, Emmanuel Nicolas, Thibault Cantat

► **To cite this version:**

Louise Ponsard, Emmanuel Nicolas, Thibault Cantat. Understanding the catalytic carbonylation of epoxides for the valorization of carbon monoxide. GECOM-CONCOORD 2019, May 2019, Erquy, France. cea-02339556

HAL Id: cea-02339556

<https://cea.hal.science/cea-02339556v1>

Submitted on 30 Oct 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Understanding the catalytic carbonylation of epoxides for the valorization of carbon monoxide

L. Ponsard, E. Nicolas, T. Cantat*

NIMBE, CEA, CNRS, Université Paris-Saclay, CEA Saclay, 91191 Gif-sur-Yvette cedex, France


thibault.cantat@cea.fr

Small molecules released as gaseous wastes by the industries, such as CO and CO₂, are attractive C1 building blocks for the production of chemicals. They can indeed reduce our dependence on fossil feedstocks. The catalytic insertion of carbon dioxide into epoxides has been developed over the last decades for the production of cyclic and polymeric carbonates, with industrial success.¹

The same strategy can be used for the production of polyester from epoxides and carbon monoxide, possibly released from the production of steel.

The catalytic carbonylation of epoxides with gaseous CO leads to β -lactones and this reaction has attracted a particular attention over the past few years. Yet, only few catalytic systems have been developed, all containing a cobalt carbonyl complex assisted by a Lewis acid.^{2,3,4,5}

Herein, we disclose a theoretical study to understand the mechanism⁶ of carbonylation of epoxides using a cobalt catalyst with Lewis acids. The influence of the Lewis acidity on the activity was investigated through various boron compounds. The DFT calculations, linked with a Lewis acidity scale, can thus be used as a guide for the development of a novel catalytic system.



^{1a} M. Aresta, *Wiley-VCH*, **2010**, *Weinheim*. ^b J. Langanke, A. Wolf, J. Hofmann, K. Böhm, M. Subhani, T. Müller, W. Leitner, and C. Gürtler, *Green Chem.*, **2014**, *16*, 1865. ^c Covestro's Cardyon® polyols made from CO₂, <https://www.covestro.com/en/cardyon/cardyon>

² J.T. Lee, P.J. Thomas, H. Alper, *JOC*, **2001**, *66*, 5424.

³ Y.D.Y.L. Getzler, V. Mahadevan, E.B. Lobkovsky, G.W. Coates, *J. Am. Chem. Soc.*, **2002**, *124*, 1174.

⁴ J.A.R. Schmidt, V. Mahadevan, Y.D.Y.L. Getzler, G.W. Coates, *Org. Lett.*, **2004**, *6*, 373.

⁵ J.W. Kramer, E.B. Lobkovsky, G.W. Coates, *Org. Lett.*; **2006**, *8*, 3709.

⁶ F. Molnar, G.A. Luinstra, M. Allmendinger, B. Rieger, *Chem. Eur. J.* **2003**, *9*, 1273.