



**HAL**  
open science

## Formic acid disproportionation to methanol and CO<sub>2</sub> reduction

Emmanuel Nicolas, Alicia Aloisi, Solene Savourey, Thibault Cantat

► **To cite this version:**

Emmanuel Nicolas, Alicia Aloisi, Solene Savourey, Thibault Cantat. Formic acid disproportionation to methanol and CO<sub>2</sub> reduction. GYSS@one-North, Jan 2018, Singapore, Singapore. cea-02339603

**HAL Id: cea-02339603**

**<https://cea.hal.science/cea-02339603>**

Submitted on 30 Oct 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# FORMIC ACID DISPROPORTIONATION TO METHANOL AND CO<sub>2</sub> REDUCTION

FROM RESEARCH TO INDUSTRY

cea


EMMANUEL NICOLAS, ALICIA ALOISI, SOLÈNE SAVOUREY, THIBAUT CANTAT

LCMCE, NIMBE, CEA, CNRS, Université Paris-Saclay, France


emmanuel.nicolas@cea.fr

CO<sub>2</sub> reduction to formic acid is a well-known process, but non-noble metal catalysts are required to perform this deed in an economically viable way. Moreover, full conversion of CO<sub>2</sub> to methanol is usually difficult for kinetic reasons (simultaneous transfer of 6 electrons and protons). We propose the use of Ru catalysts for formic acid disproportionation to methanol, and the design of new Cu-based catalysts that are capable of CO<sub>2</sub> reduction to the formate level.


## Concept




## Catalytic Studies



## Mechanism for Ru catalysts



## New Cu complexes



We demonstrated that a well chosen Ruthenium complex is able to disproportionate formic acid to methanol, with a pressure dependent selectivity. A new hydride copper complex can stoichiometrically reduce CO<sub>2</sub> to formic acid. Further studies to make the system catalytic are on the way in our laboratory.

A. J. M. Miller, D. M. Heinekey, J. M. Mayer, K. I. Goldberg, *Angew. Chemie Int. Ed.* **2013**, *52*, 3981–3984.

S. Savourey, G. Lefèvre, J.-C. Berthet, P. Thuéry, C. Genre, T. Cantat, *Angew. Chemie Int. Ed.* **2014**, *53*, 10466–10470.

A. Aloisi, E. Nicolas, J.-C. Berthet, P. Thuéry, T. Cantat, *Submitted*



European Research Council  
 ADIME

GYSS@ONE-NORTH 2018, SINGAPORE

INSTITUT DE FRANCE  
 FONDATION LOUIS D.



Agence de l'Environnement  
 et de la Meritrie de France