



**HAL**  
open science

## Phase separated structures of concentrated polymer solutions

Patrick P. Guenoun, Hervé Manzanarez, Hernan Garate, Denis Bouyer, Andre Deratani, Catherine Faur, Mathieu Jacob, Jean-Pierre Mericq, Olivier Lorain, Catherine Pochat-Bohatier

► **To cite this version:**

Patrick P. Guenoun, Hervé Manzanarez, Hernan Garate, Denis Bouyer, Andre Deratani, et al.. Phase separated structures of concentrated polymer solutions. Bordeaux Polymer Conference 2018, May 2018, Bordeaux, France. cea-02340260

**HAL Id: cea-02340260**

**<https://cea.hal.science/cea-02340260v1>**

Submitted on 30 Oct 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# PHASE SEPARATED STRUCTURES OF CONCENTRATED POLYMER SOLUTIONS

**Patrick GUENOUN<sup>1</sup>, Hervé MANZANAREZ<sup>2</sup>, Hernan GARATE<sup>1</sup>, Denis BOUYER<sup>2</sup>,  
André DERATANI<sup>2</sup>, Catherine FAUR<sup>2</sup>, Mathieu JACOB<sup>3</sup>, Jean-Pierre MERICQ<sup>2</sup>,  
Olivier LORAIN<sup>4</sup>, Catherine POCHAT-BOHATIER<sup>2</sup>**


<sup>1</sup>Université Paris-Saclay, LIONS, CEA Saclay, 91191 Gif sur Yvette cedex

<sup>2</sup>IEM (Institut Européen des Membranes) UMR 5635 (CNRS-ENSCM-UM2), Université Montpellier, Place Eugène Bataillon, F-34095 Montpellier, France


<sup>3</sup>Total, Pôle d'étude et de recherche de Lacq, Pôle économique 2, BP 47 – 64170, LACQ, France

<sup>4</sup>Polymem - 3 rue de L'industrie - Zone de VIC, 31320 Castanet-Tolosan - France


Phase separation of concentrated homopolymer solutions is both of fundamental interest (the large difference in viscosity/elasticity of the two phases can lead to unusual behaviors) and of practical importance (novel porous structures can be made by this process). I will first review how 2D phase-field simulations (Fig.1) reveal the influence of the mobility dependence with concentration for capturing features of phase separation like growth laws.[1] On the experimental side I will examine various water-soluble polymers and show how anomalous phase diagrams seem to be closely connected to unusual features of arrested-like phase separation as evidenced by light scattering and confocal microscopy. Thin films of these polymer solutions were used for making membranes (Fig. 2), avoiding the use of organic solvents. [2, 3, 4] The case of homo-polyelectrolytes is also of great interest since the theory is still controversial, as I will briefly recall. First results will be presented about phase separation upon changes in salt concentration, polymer concentration and temperature for a polyelectrolyte whose structure is also promising for membranes.



**Figure 1** – 2D Simulation of phase separation in polymer solution



**Figure 2:** Membrane made by phase separation of a PVA solution



**Figure 3:** Confocal image of a phase separating polyelectrolyte

## References

- [1] Manzanarez H. Mericq, J. P.; Guenoun, P.; Chikina, J. & Bouyer, D., *Chemical Engineering Science*, **2017**, 173, 411-427
- [2] M'Barki O. Hanafia, A.; Bouyer, D.; Faur, C.; Sescousse, R.; Delabre, U.; Blot, C.; Guenoun, P.; Deratani, A.; Quemener, D. & Pochat-Bohatier, C. al., *Journal of Membrane Science*, **2014**, 458, 225-235
- [3] Hanafia A. Faur, C.; Deratani, A.; Guenoun, P.; Garate, H.; Quemener, D.; Pochat-Bohatier, C. & Bouyer, D., *Journal of Membrane Science*, 2016, 526, 212-220
- [4] Bouyer D. M'Barki, O.; Pochat-Bohatier, C.; Faur, C.; Petit, E. & Guenoun, P *AIChE Journal*, **2017**, 63, 3035-3047