



Disproportionation of Silyl Formates, a Novel Strategy for Renewable Methanol Production

Clément Chauvier, Arnaud Imberdis, Pierre Thuéry, Thibault Cantat

► To cite this version:

Clément Chauvier, Arnaud Imberdis, Pierre Thuéry, Thibault Cantat. Disproportionation of Silyl Formates, a Novel Strategy for Renewable Methanol Production. International Conference On Phosphorus, Boron and Silicon (PBSi 2018), Dec 2018, Barcelona, Spain. cea-02340728

HAL Id: cea-02340728

<https://cea.hal.science/cea-02340728>

Submitted on 31 Oct 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Disproportionation of Silyl Formates, a Novel Strategy for Renewable Methanol Production

Clément Chauvier, Arnaud Imberdis, Pierre Thuéry and Thibault Cantat

Methanol (CH_3OH) is a useful C_1 building block employed in the chemical industry as precursor of important commodity chemicals such as formaldehyde, acetic acid or light olefins via the methanol-to-olefins (MTO) process. Nevertheless, the current methanol production methods is negatively impacted by the use of fossil resources. The production of CH_3OH would therefore strongly benefit from the advent of more sustainable processes like the six-electron reduction of CO_2 by hydrogenation, for example. An alternative strategy has recently emerged to generate CH_3OH from CO_2 via the intermediacy of formic acid (FA) as a C–H bond shuttle. At the onset, CO_2 is first reduced into FA by a relatively facile and intensively studied two-electron reduction (e.g. electroreduction). Subsequently, CH_3OH is produced by the disproportionation of FA. Under standard conditions, the competitive dehydrogenation and the targeted disproportionation of FA however display similar energy balances. To overcome this inextricable hurdle, we have thus sought a novel disproportionative route to methanol that suppresses the competing dehydrogenation pathway, while remaining attractive from the viewpoint of renewability. Towards that goal, we present a novel strategy based on the high-yielding catalytic disproportionation of silyl formates (HCO_2SiR_3) to methoxysilane (CH_3OSiR_3) and CO_2 followed by the release of free CH_3OH and the complete recycling of the silicon-containing byproducts generated throughout the process.

