



HAL
open science

Accurate determination of pH via hyperpolarized ^{129}Xe NMR

Estelle Léonce, Thierry Brotin, Jean-Pierre Dognon, Patrick Berthault

► To cite this version:

Estelle Léonce, Thierry Brotin, Jean-Pierre Dognon, Patrick Berthault. Accurate determination of pH via hyperpolarized ^{129}Xe NMR. 39th FGMR Annual Discussion Meeting, Sep 2017, Bayreuth, Germany. cea-02341010

HAL Id: cea-02341010

<https://cea.hal.science/cea-02341010v1>

Submitted on 31 Oct 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accurate determination of pH via hyperpolarized ^{129}Xe NMR

Estelle Léonce¹, Thierry Brotin², Jean-Pierre Dognon¹, Patrick Berthault¹

¹NIMBE, CEA, CNRS, Université Paris-Saclay, CEA Saclay, 91191 Gif-sur-Yvette, France

²ENS Lyon, Laboratoire de Chimie, 69364 Lyon cedex 07, France

Contact: patrick.berthault@cea.fr

Measuring non-invasively and in a local way the *in vivo* extra-cellular pH is of importance for the diagnosis of insidious pathologies such as various cancers. For one decade, the use of hyperpolarized agents has been proposed; each method possesses its own advantage and inconvenience, and their applicability in the *in vivo* domain is not always straightforward. For instance, some years ago, we proposed the use of cage-molecules bearing ionizable groups, in which laser-polarized xenon exhibits a pH-dependent chemical shift.¹ However extracting pH from measurement of the absolute chemical shift of caged xenon or even of the chemical shift difference between free and caged xenon is susceptible to be biased by external parameters such as temperature and magnetic susceptibility fluctuation.

Here we present an approach which is simple and fast to implement and which should be exempt of most of the problems encountered *in vivo*. The idea is to perform a differential measurement by using a pair of pH-sensitive ^{129}Xe NMR-based sensors. The method is rendered very pH-sensitive *via* the choice of two xenon host molecules in




Figure 1. High field region of the ^{129}Xe NMR spectrum of an aqueous solution containing the two cryptophanes **1** and **2** at 1 μM each. Evolution as a function of temperature.

(1) Berthault, P.; Desvaux, H.; Wendlinger, T.; Gyejacquot, M.; Stopin, A.; Brotin, T.; Dutasta, J.-P.; Boulard, Y. *Chem. – A Eur. J.* **2010**, *16*, 12941.

**Abstract submission form for the
39th FGMR Annual Discussion Meeting 2017
Joint Discussion Meeting of the French and German
Magnetic Resonance Societies**

Please use this form and the template on next page to write your abstract.

To submit your abstract, save this file as a doc or docx file, name it as follows: Surname_FirstName.docx (with the surname and first name of the presenting author), and email it to:

fgmr-gmrm-2017@uni-bayreuth.de

Please choose the desired presentation type:

Oral

Poster

Please choose the session for presenting your work:

- EPR methods (chair: Dieter Suter)
- Small molecules / Homogeneous catalysis (chair: Ruth Gschwind)
- Soft matter (chair: Kay Saalwächter)
- Relaxometry (chair: Ernst Rößler)
- Computational methods / Fast NMR (chair: Steffen Glaser)
- NMR crystallography / Materials (chair: Charlotte Martineau-Corcós)
- Structural biology (chair: Bruno Kieffer)
- Bio-macromolecules and dynamics (chair: Paul Rösch)
- MRI (chair: Olivier Girard)
- New methods (chair: Anne Lesage)
- Hyperpolarization methods (chair: Patrick Berthault)
- Paramagnetic systems (chair: Bruno Guigliarelli)