



**HAL**  
open science

# Complexes of tripodal phosphine ligands: synthesis, structure and catalytic activities in the hydroboration of CO<sub>2</sub>

A Aloisi, J-C Berthet, C Genre, Pierre Thuéry, T. Cantat

## ► To cite this version:

A Aloisi, J-C Berthet, C Genre, Pierre Thuéry, T. Cantat. Complexes of tripodal phosphine ligands: synthesis, structure and catalytic activities in the hydroboration of CO<sub>2</sub>. Journées de Chimie de Coordination de la SCF, Jan 2017, Grenoble, France. cea-02342086

**HAL Id: cea-02342086**

**<https://cea.hal.science/cea-02342086>**

Submitted on 31 Oct 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.


## COMPLEXES OF TRIPODAL PHOSPHINE LIGANDS: SYNTHESIS, STRUCTURE AND CATALYTIC ACTIVITIES IN THE HYDROBORATION OF CO<sub>2</sub><sup>1</sup>

A. Aloisi, J-C Berthet, C. Genre, P. Thuéry and T. Cantat

NIMBE, CEA, CNRS-UMR 3685, Université Paris-Saclay, CEA Saclay, 91191 Gif-sur-Yvette

E-mail: alicia.aloisi@cea.fr

Tripodal ligands play a major role in transition metal chemistry by increasing the stability of their complexes and enabling a fine control of the reactivity at the metal center through their steric and electronic properties and their specific coordination geometry. Among these ligands, triphos (MeC(CH<sub>2</sub>PPh<sub>2</sub>)<sub>3</sub>) has attracted considerable attention over the last years as it was successfully used for several applications in catalysis, including CO<sub>2</sub> hydrogenation, disproportionation of formic acid and reduction of amides<sup>2</sup>. Because small changes within a ligand can induce different outcomes in a catalytic reaction, variations on the triphos backbone have been reported<sup>3</sup>. In this study we report the use of PhSi(CH<sub>2</sub>PPh<sub>2</sub>)<sub>3</sub> and its phosphinite derivative PhSi(OPPh<sub>2</sub>)<sub>3</sub> in the coordination chemistry of first-row transition metals, namely iron(II), cobalt(II) and copper(I).



Scheme: Synthesis of iron, cobalt and copper complexes with PhSi(E)PPh<sub>2</sub> ligands (E = O, CH<sub>2</sub>)

Because CO<sub>2</sub> is a stable chemical waste, its selective multi-electron reduction under mild conditions remains a challenge<sup>4</sup> and an ideal platform to establish structure/activity relationships for analogous catalysts. The catalytic activity of complexes (1, 1', 2, 2', 2'', 3, 3') was thus evaluated in the hydroboration of CO<sub>2</sub> so as to determine the influence of the metal center and the electronic nature of the ligand in this model reaction. Within this range of complexes (scheme), copper catalysts (2 and 2') are both more reactive and more selective, enabling the formation of an acetal product from CO<sub>2</sub>.

**Acknowledgments :** This work was supported financially by the CEA, CNRS, CHARMMMAT Laboratory of Excellence, and the European Research Council (ERC starting grant agreement no. 336467). T.C. thanks the Foundation Louis D.-Institut de France for its support. The authors thank Dr. Emmanuel Nicolas for fruitful discussions regarding NMR spectroscopy.

**Statut :** Doctorant, 2<sup>e</sup> année

**N° adhérent SCF :** (si applicable) 28363

[1] Cantat et al., Dalton Trans., 2016, 45, 14774.

[2] Kalck et al., Coord. Chem. Rev., 2003, 236, 143; Kaska et al., Chem. Rev. 1994, 94, 1239; Leitner et al., Angew. Chem., Int. Ed., 2012, 51, 7499; Cantat et al., Angew. Chem., Int. Ed., 2014, 53, 10466.

[3] Miller et al., ACS Catal., 2015, 5, 2500; Leitner et al., J. Am. Chem. Soc., 2014, 136, 13217; Orlandini et al., Inorg. Chem., 1985, 24, 924.

[4] Beller et al., Nat. Commun., 2015, 6; Aresta, Carbon Dioxide as Chemical Feedstock, Wiley-VCH, Weinheim, 2010.