



**HAL**  
open science

## Surface Analysis of Isolated Nanoobjects by XPS

Olivier Sublemontier, S. Benkoula, Damien Aureau, C. Nicolas, Minna Patanen, Harald Kintz, X. Liu, Marc-Antoine Gaveau, Jean-Luc Le Garrec, E. Robert, et al.

### ► To cite this version:

Olivier Sublemontier, S. Benkoula, Damien Aureau, C. Nicolas, Minna Patanen, et al.. Surface Analysis of Isolated Nanoobjects by XPS. XIIIth International Conference on Nanostructured Materials (Nano2016), Aug 2016, Quebec, Canada. <cea-02346115>

**HAL Id: cea-02346115**

**<https://cea.hal.science/cea-02346115v1>**

Submitted on 4 Nov 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

## Surface Analysis of Isolated Nanoobjects by XPS

O. Sublemontier<sup>1\*</sup>, S. Benkoula<sup>2</sup>, D. Aureau<sup>3</sup>, C. Nicolas<sup>2</sup>, M. Patanen<sup>2</sup>, H. Kintz<sup>1</sup>, X. Liu<sup>2</sup>, M.A. Gaveau<sup>1</sup>, J.L. Le Garrec<sup>4</sup>, E. Robert<sup>2</sup>, F.A. Barreda<sup>1,2</sup>, A. Etcheberry<sup>3</sup>, C. Reynaud<sup>1</sup>, J.B. Mitchell<sup>4</sup> and C. Miron<sup>2</sup>


\**olvier.sublemontier@cea.fr*

<sup>1</sup>NIMBE-UMR 3685/CEA/CNRS/Université Paris-Saclay, 91191 Gif-sur-Yvette cedex, France, <sup>2</sup>Synchrotron SOLEIL, L'Orme des Merisiers, Saint-Aubin, BP 48, 91192 Gif-sur-Yvette Cedex, France, <sup>3</sup>Institut Lavoisier de Versailles, Université Versailles-St Quentin, UMR CNRS 8180, 78035 Versailles, France, <sup>4</sup>Institut de Physique de Rennes, Université Rennes 1, 35042 Rennes, France

X-ray photoelectron spectroscopy (XPS) is a very efficient and still progressing surface analysis technique. However, when applied to nanoobjects, this technique faces drawbacks due to interactions with the substrate and sample charging effects<sup>1</sup>. We present a new experimental approach to XPS<sup>2</sup> based on coupling soft x-ray synchrotron radiation with an in vacuum beam of free nanoparticles, focused by an aerodynamic lens system (Fig 1a). Two examples of experiments performed on the PLEIADES beamline at the SOLEIL Synchrotron facility are presented to illustrate the effectiveness of this approach.

In the first example, the structure of the Si/SiO<sub>2</sub> interface is probed on isolated silicon nanocrystals previously oxidized with ambient air<sup>1</sup> (Fig. 1b) or by heat treatment under air. Full characterization of the surface has been achieved for different sizes of nanocrystals between 4 and 80 nm and with different oxidizing treatments. The technique allows probing the presence of various oxidation states at the interface and to deduce therefrom a thickness of the oxide layer. For the smaller and more oxidized nanoparticles, a relatively abrupt interface including Si = O double bond is highlighted.

In the second example, the adsorption of water on the surface of TiO<sub>2</sub> nanoparticles is investigated in the gas phase<sup>3</sup>. TiO<sub>2</sub> free aerosols are exposed to a controlled pressure of water vapor before being analyzed directly by XPS. The technique allows here the observation of a predominantly dissociative adsorption of water on the surface of TiO<sub>2</sub> in its first stage, highlighting a largely covered surface by OH groups (Fig. 1c).



**Fig. 1a.** XPS of isolated nanoparticles. **Fig1b.** Si2p XPS spectrum of isolated silicon nanocrystals. **Fig 1c.** O1s XPS spectrum of hydrated TiO<sub>2</sub> nanoparticles.

### References

- [1] Baer, D. R., Gaspar, D. J., Nachimuthu, P., Techane, S. D. & Castner, D. G. *Anal. Bioanal. Chem.* 2010, **396**, 983.

[2] Sublemontier, O. et al. *J. Phys. Chem. Lett.* 2014, **5**, 3399.

[3] Benkoula, S. et al. *Sci. Rep.* 2015, **5**, 15088.

### **Acknowledgements**

The experiments were performed at the PLEIADES beamline at the SOLEIL Synchrotron (Proposals No. 20110423 and 20130466). The development of the MPSC has received funding from the Agence Nationale de la Recherche (ANR) under Grant No. ANR-07-NANO-0031.