



Photo-activation therapy with high-Z nanoparticles : modelling at a micrometer level and experimental comparison

Rachel DELORME

CEA, LIST, Laboratoire de Modélisation et Simulation des Systèmes, 91191 Gif-sur-Yvette CEDEX, France

Supervisors : Mathieu AGELOU (CEA, LIST, LM2S),

Christophe CHAMPION (LPMC,Metz),

Hélène ELLEAUME (INSERM, Grenoble)





Summary

- - Context and objectives
 - Simulation results (PENELOPE)
 - Experimental comparison
 - Code comparison with GEANT4
 - Conclusions and Prospects





digiteo

CARNOT CEA LIST

• Photo-activation therapy : combines low energy X-ray beams with high-Z elements (iodine, gold, gadolinium, platinum...).



Ratio of X-ray mass energy-absorption coefficients Gd or Au / water (NIST)



- Discovery of the radiosensitive effect due to the iodinated contrast products (ICP) in interventional radiology (Norman et al, Radiology (1978)).
- Studies continued by the team INSERM / ESRF: treatment of high grade gliomas (clinical trial of phase I)

ICP : Enhancement factor ~ 2 (for 10 mg/ml). Do not integrate cells.

 Hainfeld et al. (*Phys. Med. Biol. (2004)*): Survival response up to a factor of 4, treating cancerous mice with goldnanoparticles (AuNP) of 1.9nm (RX tube at 250kVp).

digite





Cez

- Cho et al. (*Phys. Med. Biol. (2005)*): Monte Carlo calculation of tumour dose enhancement with homogeneous gold up to a factor of 2.1 with Hainfeld's conditions.
- Such radiobiological enhancement not explained by macroscopic physical model. Modelling at the lower scale.
- Zhang et al. (Biomed Microdevices (2009)): Comparison of macroscopic dose calculations for a homogeneous goldwater model and nano-structured.
 Homogeneous case : overestimation of

the dose up to 16%.

• Modelling the nanoparticles.

digiteo



- Objectives, Monte Carlo simulations :
 - Challenge : find an appropriate physical observable to correlate with biological data.
 - Gold and gadolinium nanoparticles (AuNP & GdNP) : dose and electron spectra emitted from NP under irradiation (PENELOPE).
 - Unicellular geometry with gadolinium : comparison with experiments made at ESRF (European Synchrotron Radiation Facility) (PENELOPE).
 - Unicellular geometry with nanoparticles of gadolinium (GEANT4).





Electron range in water (Penelope tables)





- **PENELOPE** : Monte Carlo code adapted for low energy electron and photon transport (lower limit of electron transport : 50 eV). Variance reductions ++.
- First study : Dose deposition and spectra of secondary particles emitted from a single AuNP and GdNP.

Photon detector

Electron detector

GdNP / AuNP







Simulation results Spectra from GdNP

- X-ray beam energy : 52 keV. Diameter of GdNP : 10 nm.
- Gd K-edge : 50.2 keV.
- Au K-edge : 80.7 keV.
 - Relaxation cascade
 and X-ray interaction
 with shell and subshell well described in
 Penelope.

Electron spectra emitted from a 10 nm diameter AuNP and GdNP as a function of beam energy.

Quantity



- More electrons produced in Gold except • after the sharp Gadolinium K-edge.
- Mean energy lower after the Gd Kedge, provides a more "local" dose deposition.

• Dose deposition at 500nm around a AuNP of 100nm with a 85 keV X-ray beam.

In water

With a centred AuNP



AuNP : Increase of the dose up to a factor 100 with quasi-isotropic diffusion.



- In vitro experiments (ESRF, May/Sept 2011) : radiosensitivity on rat glioma cell line F98 with ultra small GdNP (*Angew. Chem. Int. Ed. 2011*).
 - Beam energies : 31, 49.5, 51, 65, 80 (synchrotron) and 1253 keV (Co60).
 - 5 experimental conditions :



» Internal Gd : 0.6 mg/mL

(cf. F. Taupin presentation on Monday (GRC))

» External Gd : 2.1 mg/mL (1.8 mg/mL for the incubated case)

• Simulation : Unicellular geometry.

Cea

Nanoparticles substituted with homogeneous water-gadolinium mixture localized into or outside the cell (experimental concentrations).



Square monochromatic photon source, 15µm



 Size of the extra-cellular medium : 15 to 110 µm. Same size for the X-Ray beam.



digiteo

cea



Probability density (int & ext Gd / water)



Experimental comparison Electron spectra incoming on nucleus

• Comparison of electron spectra incoming on the nucleus for all energies and conditions.

(ex : int&ext Gd / water)

 Experimental comparison normalized at 65 keV : Good correlation.

16



- GEANT4 : General purpose Monte Carlo code allowing to transport particles in matter for a wide range of applications (high energy physics, nuclear, space, medical...).
- Electromagnetic physical models for low energy electron and photon transport :
 - Livermore
 - Penelope
 - DNA
- Advantages compared to PENELOPE code :
 - Flexible
 - Complex geometries
 - Access to more detailed track information
- Drawbacks :
 - Lack of variance reductions for small structures.



Code comparison : GEANT4 Validation : doses, homogeneous media

- Macroscopical DEF : Cube of 1 cm.
- Microscopical DEF : Cell.



• Coherent results between PENELOPE and GEANT4: problem of statistics for the small volumes (variance reduction).

Cea



Cell geometry : consideration of nanoparticles of different sizes (R = 15 – 50 nm)



digiteo

CARNOT

Cea



 Coherent results between homogeneous / NP. Auto-absorption of low-energy electrons into the NP.



Cea

- Photon interactions : photoelectric, Compton, Rayleigh
- Electron interactions : Ionization, multiple scattering

digiteo

CARNOT



- Conclusions :
 - Nanoparticle model (*PENELOPE*): Study of secondary particles emitted under irradiation for different beam energies and comparison Gd/Au.
 - Unicellular model (*PENELOPE*) : calculation of DEF and electron spectra incoming on the nucleus for different conditions.
 - Comparison with experimental data (*PENELOPE*) : encouraging results.
 - Limitation of PENELOPE modeling large number of nanoparticles : GEANT4.
 - Validation GEANT4 / PENELOPE. Very long simulation time with NP for acceptable statistic.
- Prospects :
 - Improve GEANT4 physical model for nanostructures applications.
 - Investigation of track information.

- LM2S:
 - Bénédicte POUMAREDE, Mathieu AGELOU, Juan-Carlos GARCIA-HERNANDEZ, François TOLA.
- Supervisors :
 - Mathieu AGELOU (LM2S)
 - Christophe CHAMPION (Univ. De Lorraine, Metz)
 - Hélène ELLEAUME (INSERM / ESRF)
- Co-workers:
 - Florence TAUPIN (INSERM / ESRF)
 - Mélanie FLAENDERS (INSERM / ESRF)
 - Jean-Luc RAVANAT (LAN, CEA Grenoble)
 - Sébastien INCERTI (CENBG, Bordeaux)











Comparison of electron spectra between PENELOPE and GEANT4.
 (52 keV, NPGd 10 nm diameter). Good correlation. Lack of statistic for GEANT4.



DEF 1µm around 1 NP

- Gd properties : K-edge : 50.2 keV, Z : 64, density: 7.9 g/cm³
- Au properties : K-edge : 80.7 keV, Z : 79, density: 19.3 g/cm³

cea

CARNOT CEA LIST digiteo

• 1µm dose deposition of particles emitted from a AuNP as a function of energy and diameter:







Experimental comparison Normalized DEF

- Normalization at 65 keV to compare the tendency between the Nucleus DEF and SER_{4Gv}.
- Model and observable not well adapted for comparison with SER.
- Remarks :
 - Limited size of external volume.
 - Biological effects not limited to physical dose.

Cea