



HAL
open science

Constraining sediment dynamics in rivers using fallout radionuclides: How to move forward from the lessons learnt in catchments of Mexico, Laos and France

Olivier Evrard, David Gateuille, Elian Gourdin, Sylvain Huon, Julien Nemery, Nicolas Gratiot, Irène Lefevre, Sophie Ayrault, Philippe Bonté

► **To cite this version:**

Olivier Evrard, David Gateuille, Elian Gourdin, Sylvain Huon, Julien Nemery, et al.. Constraining sediment dynamics in rivers using fallout radionuclides: How to move forward from the lessons learnt in catchments of Mexico, Laos and France. ICCE 2014 Symposium: The International Symposium on Sediment Dynamics: From the Summit to the Sea, Dec 2014, New Orleans, United States. <cea-02669005>

HAL Id: cea-02669005

<https://cea.hal.science/cea-02669005v1>

Submitted on 31 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Constraining sediment dynamics in rivers using fallout radionuclides: How to move forward from the lessons learnt in catchments of Mexico, Laos and France

Olivier Evrard¹, David Gateuille¹, Elian Gourdin¹, Sylvain Huon², Julien Nemery³,
Nicolas Gratiot³, Irène Lefèvre¹, Sophie Ayrault¹, Philippe Bonté¹

¹Laboratoire des Sciences du Climat et de l'Environnement (CEA, CNRS, UVSQ; LSCE/IPSL), Gif-sur-Yvette, France, olivier.evrard@lsce.ipsl.fr; ²Université Pierre et Marie Curie (Bioemco), Paris, France ; ³Laboratoire d'étude des Transferts en Hydrologie et en Environnement (Université Grenoble Alpes, IRD, CNRS, LTHE), Grenoble, France.

ABSTRACT

Land use change (e.g. deforestation, overgrazing, cropping intensification) and the associated acceleration of soil erosion have led to an increase in sediment supply to rivers across the world during the last decades. This degradation leads to important on-site (e.g., decrease in soil fertility) and off-site (e.g., reservoir siltation and water pollution) impacts. In order to better understand the dynamics of sediment in catchments and rivers and to implement efficient mitigation measures, it is necessary to better constrain the pathways of sediment in time and space. Fallout radionuclides characterized by different half-lives and origins (Be-7, Cs-137, and unsupported Pb-210) have proved to provide very useful indications to constrain the transfer times of sediment at various scales (plot, hillslope, small catchment, large basins) and in a range of different environments and climates across the world. However, several limitations were raised regarding the assumptions underpinning the methods used to derive sediment residence time or transport distances. Furthermore, doubts regarding the validity of using those tracers in large rivers and catchments were also put forward in recently published papers.

In this study, we provide additional data indicating that the hypotheses underlying the method are meaningful. We also compiled radionuclide data (Be-7, Cs-137, and unsupported Pb-210) measured in both rainfall and suspended sediment during recent studies that were conducted between 2009–2013 in catchments of different sizes (3–60,000 km²) and located in different environments (Central Mexico, Northern Laos, France). The results obtained in the different catchments are compared and reinterpreted, with the specific objective of dealing with the limitation that fresh Be-7 fallout may not label river bank or subsurface material.

The results show that providing additional and independent information on the sources delivering sediment to the rivers helps constraining their transit times with the method based on Be-7 and unsupported Pb-210 measurements. Implications of the use of this technique in larger rivers and catchments are discussed. Their contribution to quantify the persistence time of particle-borne contaminants in those riverine systems is specifically addressed.