



HAL
open science

On positive functions with positive fourier transforms

B.G. Giraud, R. Peschanski

► **To cite this version:**

B.G. Giraud, R. Peschanski. On positive functions with positive fourier transforms. Acta Physica Polonica B, 2006, 37 (2), pp.331-346. cea-02889284

HAL Id: cea-02889284

<https://cea.hal.science/cea-02889284>

Submitted on 3 Jul 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON POSITIVE FUNCTIONS WITH POSITIVE FOURIER TRANSFORMS

B.G. GIRAUD[†], R. PESCHANSKI[‡]

Service de Physique Théorique, DSM
CE Saclay, F-91191 Gif/Yvette, France

(Received November 10, 2005)

Using the basis of Hermite–Fourier functions (*i.e.* the quantum oscillator eigenstates) and the Sturm theorem, we derive constraints for a function and its Fourier transform to be both real and positive. We propose a constructive method based on the algebra of Hermite polynomials. Applications are extended to the 2-dimensional case (*i.e.* Fourier–Bessel transforms and the algebra of Laguerre polynomials) and to adding constraints on derivatives, such as monotonicity or convexity.

PACS numbers: 02.30.Gp, 02.30.Mv, 02.30.Nw, 12.38.–t

1. Introduction

Positivity conditions for the Fourier transform of a function occur in various domains of physics. One often asks:

- What are the constraints for a real function $\psi(r)$ ensuring that its Fourier transform

$$\varphi(s) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} dr e^{isr} \psi(r) \quad (1)$$

be real and positive?

- Conversely, what are the properties of φ if ψ is real and positive?
- Finally, what are the constraints on Fourier partners such that *both* ψ and φ be real and positive?

[†] giraud@dsm-mail.saclay.cea.fr

[‡] pesch@dsm-mail.saclay.cea.fr

Physicists often need to work with concrete constructions. The practical construction of a basis of functions satisfying the above mentioned positivity properties remains, up to our knowledge, an open problem. Such questions are quite relevant in Physics. As practical examples, let us quote two well-known cases. A Fourier transform relates [1] two quantities, namely the cross section and the profile of a nucleus, which both ought to be positive. In particle physics, a 2-d Fourier-Bessel transform relates the color dipole distribution in transverse position space (derived from Quantum Chromodynamics) and the transverse momentum distribution of gluons probed during a deep-inelastic collision [2]. Such questions occur also in probability calculus, for the relation between probability distributions and characteristic functions [3], in crystallography and in condensed matter physics, *e.g.* for the interpretation of patterns, *etc.*

The problem is simplified if related to another one, that concerning the functions which are invariant [4] up to a phase factor¹ by Fourier transforms. Indeed, the most familiar examples of positive self-dual functions or distributions, thus trivially verifying the double positivity condition, which are at most scaled under Fourier transformation (FT), are the Gaussian and the Dirac comb.

Many special cases can be found, where positivity is conserved, such as, for instance, the continuous family of functions $\exp(-r^\nu)$, where $0 < \nu \leq 2$. Various *sufficient* conditions for positivity can be found in the literature, such as the convexity of ψ [7] but, up to our knowledge, no general constructive method has been presented.

The present note attempts to give general positivity criteria, in a constructive way, by taking advantage of a representation under which the FT is essentially “transparent”. Our method combines the advantages of self-duality properties with those allowed by an algebra of polynomials, where positivity means absence of real roots, hence reasonably simple conditions for the polynomial coefficients. For this sake, in the 1-d case, we select a basis made of convenient eigenstates of the FT, the Hermite–Fourier functions, *i.e.* the harmonic oscillator eigenstates. The method extends to the 2-d case, or Fourier–Bessel transform, by replacing Hermite by Laguerre polynomials.

There are general theorems about the characterization of Fourier transforms of positive functions [9]. Let us quote in the first place the Bochner theorem and its generalizations [10] which state that the Fourier transform of a positive function is *positive-definite*. But positive definiteness in the sense of such theorems does not imply plain positivity². Hence our problem

¹ They are called “self-Fourier” in [5] if the phase factor is 1 and “generalized self-Fourier” [6] (or “dual”) for other phases.

² Positive definiteness means that for any real numbers x_1, \dots, x_k and complex numbers ξ_1, \dots, ξ_k , one has $\sum_{k,j} \varphi(x_k - x_j) \bar{\xi}_j \xi_k \geq 0$.

actually could be rephrased [11] as “build positive-definite functions that are positive”.

Our formalism is the subject of Section 2. Numerical, illustrative examples are given in Section 3. Then Section 4 extends our algorithms to the 2-d problem. Hermite polynomials are replaced by Laguerre polynomials, but the algebra remains essentially the same. A brief discussion, conclusion and outlook are offered in Section 5.

2. Basic formalism

Consider the harmonic oscillator Hamiltonian, $\frac{1}{2}(p^2 + r^2)$, and its eigenwavefunctions

$$u_n(r) = \pi^{-1/4} e^{-1/2r^2} H_n(r). \tag{2}$$

Here, we set H_n to be a *square normalized* Hermite polynomial, $(-)^n e^{r^2} d^n/dr^n e^{-r^2} / \sqrt{2^n n!}$, with a positive coefficient for its highest power term. For the sake of clarity, we list the first polynomials as, $H_0 = 1$, $H_1 = \sqrt{2} r$, $H_2 = (2r^2 - 1)/\sqrt{2}$, $H_3 = (2r^3 - 3r)/\sqrt{3}$ and their recursion relation

$$a_{n+1}H_{n+1} = 2r a_n H_n - 2n a_{n-1}H_{n-1}, \tag{3}$$

where $a_n = \sqrt{2^n n!}$. It is known that the FT of such states brings only a phase

$$\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} dr e^{isr} u_n(r) = i^n u_n(s), \tag{4}$$

and thus such states give generalized self-dual functions with phase i^n . Let one expand ψ in the oscillator basis, $\psi(r) = \sum_{n=0}^N \psi_n u_n(r)$, with a truncation at some degree N . Then all odd components ψ_{2p+1} must vanish if φ must be real, and the even rest splits, under FT, into an invariant part and a part with its sign reversed, namely

$$\begin{aligned} \varphi(s) &= \pi^{-1/4} e^{-1/2s^2} [P_+(s) - P_-(s)], \\ P_+(s) &= \sum_{p=0}^{[N/4]} \psi_{4p} H_{4p}(s), \\ P_-(s) &= \sum_{p=0}^{[(N-2)/4]} \psi_{4p+2} H_{4p+2}(s), \end{aligned} \tag{5}$$

where the usual symbols $[N/4]$ and $[(N-2)/4]$ mean, respectively, the entire parts of $N/4$ and $(N-2)/4$.

Notice that, when all components ψ_n vanish except ψ_0 , then both ψ and φ are positive, because $H_0 = 1$. Hence, one may, starting from this special point in the functional space of functions, investigate those domains of parameters ψ_n where the polynomials $\mathcal{P} = P_+ + P_-$ and $\mathcal{Q} = P_+ - P_-$ have no real root. Notice that only even powers of r and s are involved. It will therefore be convenient to use auxiliary variables $\rho = r^2$ and $\sigma = s^2$, and the domain of interest for the parameters ψ_n will correspond to the absence of real *positive* roots for both ρ and σ .

The second ingredient of our approach is the well-known Sturm theorem [12] which gives a way to characterize and localize the real roots of any given polynomial. The *Sturm criterion* can be expressed as follows:

“Given a polynomial $\mathcal{P}(x)$, its *Sturm sequence* $\mathcal{S}(x) \equiv \{\mathcal{S}_1, \mathcal{S}_2, \dots, \mathcal{S}_m, \dots, \mathcal{S}_{j \leq N}\}$ is the set of polynomials

$$\mathcal{S}_1 = \mathcal{P}, \quad \mathcal{S}_2 = \frac{d\mathcal{P}}{dx}, \quad \dots \quad \mathcal{S}_m = -\mathcal{S}_{m-2} + \left[\frac{\mathcal{S}_{m-2}}{\mathcal{S}_{m-1}} \right] \mathcal{S}_{m-1} \quad \dots, \quad (6)$$

where $[\]$ designates the polynomial quotient³. To know the number of distinct roots between $x = a$ and $x = b$, count the number $\mathcal{N}(a)$ of sign changes in $\mathcal{S}(a)$ and, similarly, count $\mathcal{N}(b)$. Then the number of roots is $|\mathcal{N}(b) - \mathcal{N}(a)|$.”

The domain borders where the root number, $|\mathcal{N}(+\infty) - \mathcal{N}(0)|$, changes have to do with cancellations of the *resultant* \mathcal{R} between \mathcal{P} and $d\mathcal{P}/dx$. The cancellation of \mathcal{R} corresponds to collisions between conjugate complex roots becoming real roots and conversely. Because of the demanded positivity of ρ and σ , the borders have also to do with sign changes of $\mathcal{P}(0)$ and $\mathcal{Q}(0)$, meaning real roots ρ and σ going through 0. All such technicalities are taken care of by the Sturm criterion, which furthermore allows the labeling of each domain by its precise number of real roots.

This will be implemented here in an explicit way, analytically as much as possible, then numerically and graphically, for a few cases of a general illustrative value. For this, we will plot the shapes of domains labeled by the values of the Sturm criterion. Hence the combination of the self-dual properties of the quantum oscillator basis and of the Sturm theorem allows a constructive method for a systematic investigation of positivity conditions for a 1-d Fourier transform. This basis has the potential to represent any function in the Hilbert space \mathcal{L}^2 , but the present study concerns a finite set of components. An extension to infinite series of components may deserve other tools.

³ The sequence, made of polynomial remainders of the division of \mathcal{S}_{n-1} by \mathcal{S}_{n-2} , with $(-)$ signs, clearly stops at some $j \leq N$.

3. Positivity domains

In this section we will apply our method to the case of a basis formed with 3 or 4 Hermite–Fourier functions. In Sec. 3.1 we consider the basis ψ_0, ψ_4, ψ_8 , then in 3.2 the basis $\psi_0, \psi_4, \psi_8, \psi_{12}$, which both lie in the subspace with eigenvalue 1, and in 3.3 the basis ψ_0, ψ_2, ψ_4 where ψ_2 is in the subspace with eigenvalue -1 , furthermore, in 3.4 the influence of an additional constraint motivated by Physics, that of monotony for φ , and finally in 3.5 a comparison with the convexity constraint for ψ . Many other illustrations are possible, but these, Sec. 3.1–3.5, demonstrate the flexibility of our approach.

3.1. Mixture of three polynomials in the subspace with eigenvalue 1

Here we assume that ψ has only components ψ_0, ψ_4 and ψ_8 , hence \mathcal{P} reduces to P_+ in (5) and we can study

$$\mathcal{P} = \psi_0 + \frac{\psi_4}{2\sqrt{6}}(4\rho^2 - 12\rho + 3) + \frac{\psi_8}{24\sqrt{70}}(16\rho^4 - 224\rho^3 + 840\rho^2 - 840\rho + 105), \quad (7)$$

where $\rho = r^2$. One maintains $\psi_8 > 0$ and one reads $P_+(0) = \psi_0 + 3\psi_4/(2\sqrt{6}) + 105\psi_8/(24\sqrt{70})$.

The resultant \mathcal{R} between P and $dP/d\rho$ is

$$\begin{aligned} \mathcal{R} \propto & 2100 \psi_0 \psi_4^4 - 1050 \sqrt{6} \psi_4^5 - 240 \sqrt{70} \psi_0^2 \psi_4^2 \psi_8 + 400 \sqrt{105} \psi_0 \psi_4^3 \psi_8 \\ & - 165 \sqrt{70} \psi_4^4 \psi_8 + 480 \psi_0^3 \psi_8^2 + 4560 \sqrt{6} \psi_0^2 \psi_4 \psi_8^2 \\ & - 13320 \psi_0 \psi_4^2 \psi_8^2 - 1600 \sqrt{6} \psi_4^3 \psi_8^2 - 792 \sqrt{70} \psi_0^2 \psi_8^3 \\ & + 1728 \sqrt{105} \psi_0 \psi_4 \psi_8^3 - 612 \sqrt{70} \psi_4^2 \psi_8^3 - 10080 \psi_0 \psi_8^4 \\ & - 2520 \sqrt{6} \psi_4 \psi_8^4 + 1260 \sqrt{70} \psi_8^5. \end{aligned} \quad (8)$$

Because of the free scaling of \mathcal{P} we normalize the polynomial so that its coefficients lie on a half sphere of unit radius

$$\begin{aligned} \psi_0 &= \cos \alpha \cos \beta, \\ \psi_4 &= \sin \alpha \cos \beta, \\ \psi_8 &= \sin \beta, \quad -\pi < \alpha \leq \pi, \quad 0 < \beta \leq \pi/2. \end{aligned} \quad (9)$$

We thus show in figure 1 the domains where the number of roots increases from 0 to 4. The no root domain is the white triangle above $\alpha = \beta = 0$ and slightly right from this point. In this domain, $\psi (\equiv \varphi)$ is self-Fourier and positive.




Fig. 1. Mixture of H_0 , H_4 and H_8 . White, no real positive ρ root; black, 4 roots; darkening grays, increasing root number.

3.2. Mixture of four polynomials in the subspace with eigenvalue 1

Now we add to ψ a component ψ_{12} , hence

$$\begin{aligned} \mathcal{P} = & \psi_0 + \frac{\psi_4}{2\sqrt{6}}(4\rho^2 - 12\rho + 3) \\ & + \frac{\psi_8}{24\sqrt{70}}(16\rho^4 - 224\rho^3 + 840\rho^2 - 840\rho + 105) \\ & + \frac{\psi_{12}}{1440\sqrt{231}}(64\rho^6 - 2112\rho^5 + 23760\rho^4 \\ & - 110880\rho^3 + 207900\rho^2 - 124740\rho + 10395). \end{aligned} \quad (10)$$

While borders corresponding to $P_+(0) = 0$ obtain easily, the resultant \mathcal{R} to be considered for other borders is unwieldy and is skipped here. Taking advantage of scaling we set:

$$\begin{aligned} \psi_0 &= \cos \alpha \cos \beta \cos \gamma, \\ \psi_4 &= \sin \alpha \cos \beta \cos \gamma, \\ \psi_8 &= \sin \beta \cos \gamma, \\ \psi_{12} &= \sin \gamma, \quad |\alpha| \leq \frac{\pi}{2}, \quad |\beta| \leq \pi, \quad 0 \leq \gamma \leq \frac{\pi}{2}. \end{aligned} \quad (11)$$

This choice of spherical coordinates was designed to ensure the positivity of ψ_{12} , obviously, but also a dominance of ψ_0 near $\alpha = \beta = \gamma = 0$. The dominance is clearly useful for the positivity of \mathcal{P} . Then this S_3 sphere can be explored by various cuts according to fixed values of γ . The results are shown in




Fig. 2. Mixture of H_0, H_4, H_8 and H_{12} . Root number maps. Color code: 0 root, red; 1, yellow; 2, yellowish green; 3, bluish green; 4, blue; 5, dark purple; 6, pink. Left: cut of the parameter sphere S_3 when $\gamma = \pi/10^6$. Right: cut for $\gamma = \pi/15$ (uncolored edition: the 0 root domain is the dark triangle-like domain near the center).




Fig. 3. Left: Same as Fig. 2, with $\gamma = 2\pi/15$. Right: $\gamma = \pi/6$. See how the no root triangle shrinks.

figures 2–5, with $\gamma = \pi/10^6, \pi/15, 2\pi/15, \pi/6, \pi/5, 7\pi/30, 4\pi/15, 3\pi/10$, respectively. The color code for the number of roots is: 0 root, red; 1, yellow; 2, yellowish green; 3, bluish green; 4, blue; 5, dark purple; 6, pink (in the uncolored edition, the no root domain, if it exists, is that dark, small or




Fig. 4. Left: $\gamma = \pi/5$; Right: $\gamma = 7\pi/30$. Absence of no root domain.




Fig. 5. Left: $\gamma = 4\pi/15$; Right: $\gamma = 3\pi/10$. Progressive dominance of H_{12} with 6 ρ roots.

tiny triangle slightly right of the map center). The red domain shrinks at first very slowly when γ increases, then faster when $\gamma \simeq \pi/6$. Beyond such an order of magnitude for γ , there is no red domain and the map becomes invaded by bigger and bigger pink patches, representing the dominance of the 6 positive, real roots of H_{12} .

3.3. Two polynomials from subspace “1” mixed with one polynomial from subspace “-1”

If we consider a mixture of H_0 , H_2 and H_4 , the FT connects the two polynomials

$$\begin{aligned} \mathcal{P}(\rho) &= \psi_0 + \psi_2 \frac{2\rho - 1}{\sqrt{2}} + \psi_4 \frac{4\rho^2 - 12\rho + 3}{2\sqrt{6}}, \\ \mathcal{Q}(\sigma) &= \psi_0 - \psi_2 \frac{2\sigma - 1}{\sqrt{2}} + \psi_4 \frac{4\sigma^2 - 12\sigma + 3}{2\sqrt{6}}. \end{aligned} \tag{12}$$

We study the positivity of each polynomial separately, then of both. Notice that the parametrization

$$\begin{aligned} \psi_0 &= \cos \alpha \cos \beta, \\ \psi_2 &= \sin \alpha \cos \beta, \\ \psi_4 &= \sin \beta, \quad -\pi < \alpha \leq \pi, \quad 0 < \beta \leq \pi/2, \end{aligned} \tag{13}$$

reverses only the sign of ψ_2 if α becomes $-\alpha$. This parity operation is seen in figure 6, the white domains of which correspond to the positivity of \mathcal{P} and \mathcal{Q} , respectively. The domain of simultaneous positivity for both is the white intersection domain in the left part of figure 7, with the expected symmetry.

It is actually easy here to analyze analytically the resultants of interest for \mathcal{P} and \mathcal{Q}

$$\mathcal{R} \propto \sqrt{6} \psi_2^2 - 4 \psi_0 \psi_4 \mp 4 \sqrt{2} \psi_2 \psi_4 + 2 \sqrt{6} \psi_4^2 \tag{14}$$




Fig. 6. Mixture of H_0 , H_2 and H_4 . White domain, 0 root. Grey, 1 root; black, 2 roots. Left: results for \mathcal{P} ; Right: \mathcal{Q} .

together with signatures for the signs of roots, such as $\mathcal{P}(0)$, *etc.* This can be done also in the “spherical representation”. The white domains of figures 6, 7 are recovered.




Fig. 7. Positivity domains for the parity mixed case. Left: for both \mathcal{P} and \mathcal{Q} . Right: \mathcal{P} , \mathcal{Q} and monotony of φ .

3.4. Positivity with monotony

For some problems [2], it may be useful to request either ψ and/or φ to be monotonous functions in an interval such as $[0, \infty]$. We illustrate this in the case of an H_0, H_2, H_4 mixture, with the additional constraint

$$\frac{d\varphi}{d\sigma} \propto -12 \cos \alpha \cos \beta + 6\sqrt{2} \cos \beta \sin \alpha (2\sigma - 5) - \sin \beta \sqrt{6} (4\sigma^2 - 28\sigma + 27) < 0. \quad (15)$$

The result appears in the right part of figure 7. The white domain, corresponding to such three simultaneous conditions of positivity and monotonicity, is a severe restriction of the white domain seen in the left part of figure 7.

3.5. Positivity from convexity

A practical condition for the positivity of φ is the convexity of ψ [7]. This has been useful in particular for the derivation of baryon mass inequalities in QCD [8]. Convexity is a sufficient, but not necessary condition. Indeed, $\varphi \sim \int_0^\infty dr [1 - \cos(sr)]/s^2 d^2\psi/dr^2 > 0$. We illustrate this convexity condition for a mixture of H_0, H_4, H_8 .




Fig. 8. H_0, H_4, H_8 mixture. Left: zoom of the positivity domain. Right: convexity domain for $r > r_c = 1$.




Fig. 9. Same mixture as Fig. 8. Convexity domains. Left: $r_c^2 = .67$. Right: $r_c^2 = .4$.

Clearly, the presence of $\exp(-r^2/2)$ in front of a finite order polynomial, with the even parity of ψ and its derivability, are contradictory with “convexity everywhere”; a smooth, round maximum must occur at the origin. We thus study partial convexity conditions of the kind, “convexity between r_c and $+\infty$ ”. A reasonable choice for the order of magnitude of r_c is the position $r_c = 1$ of the inflexion point of $\exp(-r^2/2)$. The second derivative $d^2\psi/dr^2$ belongs to the same algebra. We adjusted its Sturm criterion to various values of r_c . Only small domains are found which ensure zero roots for $d^2\psi/dr^2$, because most mixtures of H_0, H_4, H_8 do oscillate. Figures 8 and 9 show, in white again, with the parametrization by Eqs. (9), the survivor

domain obtained if $r_c = 1$, right part of figure 8, then $r_c^2 = 2/3$, left part of figure 9, and $r_c^2 = 2/5$, right part of figure 9, respectively. The domain shrinks in a smooth way when r_c^2 decreases from 1 to 0.4 and disappears if $r_c^2 < \sim 0.4$. It does not increase much when $r_c \geq \mathcal{O}(1)$. The left part of figure 8, a zoom of the left part of figure 6, shows that such partial convexity domains are already included in the positivity domain of ψ .

4. Positivity for the 2-dimensional Fourier transform

The Fourier–Bessel transform in which we are here interested reads,

$$\varphi(s) = \int_0^\infty (r dr) J_0(sr) \psi(r). \quad (16)$$

For the 2-d radial space, a complete basis of states results from substituting r^2 for r into Laguerre polynomials

$$\int_0^\infty (r dr) 2 e^{-r^2} L_m(r^2) L_n(r^2) = \delta_{mn}. \quad (17)$$

For the sake of clarity, we list here the first four such normalized, “2-d radial” states,

$$\{v_0, v_1, v_2, v_3\} = \sqrt{2} e^{-\frac{1}{2}r^2} \left\{ 1, r^2 - 1, \frac{r^4 - 4r^2 + 2}{2}, \frac{r^6 - 9r^4 + 18r^2 - 6}{6} \right\}. \quad (18)$$

One can verify that the states v_n make eigenstates of the Fourier–Bessel transform,

$$\int_0^\infty (r dr) J_0(sr) v_n(r) = (-1)^n v_n(s). \quad (19)$$

Positivity conditions can again be implemented with the Sturm criterion. For instance a mixture of v_0, v_2 , from that subspace with eigenvalue 1, and v_1 , from that with eigenvalue -1 , defines the following two reciprocal partners:

$$\begin{aligned} \mathcal{P}(\rho) &= \psi_0 + \psi_1 (\rho - 1) + \psi_2 \frac{\rho^2 - 4\rho + 2}{2}, \\ \mathcal{Q}(\sigma) &= \psi_0 - \psi_1 (\sigma - 1) + \psi_2 \frac{\sigma^2 - 4\sigma + 2}{2}. \end{aligned} \quad (20)$$

The left part of figure 10 shows the positivity domain for φ and its right part shows the joint positivity domain for ψ and φ . Similarities between

polynomials involved in the present “mixed parity” case and those of case **C** of the previous section create topological similarities with figure 6 (right) and figure 7 (left) but numerical details do differ.




Fig. 10. Positivity domains for the “radial 2-d”, parity mixed case. Left: φ . Right: both ψ and φ .




Fig. 11. Left: Positivity domain for the “radial 2-d”, v_0, v_2, v_4 mixture. Right: third derivative negativity domain if $r_t^2 = 1.4$.

Interestingly enough, a sufficient condition [7] for the positivity of φ in this 2-d situation is that the third derivative, $d^3\psi/dr^3$, be negative. But, as in the previous section, case **E**, the presence of a truncated number of basis terms in our expansion may reduce this negativity condition for $d^3\psi/dr^3$ to a domain $r \geq r_t$ only. We show in figure 11 the result for a mixture of v_0 ,

v_2 and v_4 if $r_t^2 = 1.4$, a demanding situation resulting into the tiny domain in the right part of figure 11. The domain belongs to the positivity domain seen in the left part of the same figure.

5. Discussion and conclusion

To summarize our results, we have built a basis of functions verifying positivity together with their Fourier transform. The method is based upon algebras of Hermite polynomials (for 1-dimensional FT) or Laguerre polynomials in the variable r^2 (for 2-dimensional radial FT).

The Fourier transform has four eigenvalues $1, i, -1, -i$, and thus four highly degenerate eigensubspaces. Two of such subspaces are compatible with real functions remaining real. To span each subspace, we used a basis made of Hermite–Fourier (or “Laguerre–Fourier”) states. The orders (in r) of the polynomials have to be multiples of 4 if the eigenvalue is 1, and multiples of 4 plus 2 if the eigenvalue is -1 . At the cost of a truncation of such bases to a maximum order N , the conditions for positivity, convexity, *etc.* thus reduce to simple manipulations of polynomial coefficients based on the Sturm theorem. In each truncation case, one can find suitable domains for the parameters which mix the various basis polynomials. Such domains have been illustrated by the figures shown in this paper.

Some qualitative considerations may be drawn about the solutions we found. In the left part of figure 12, we display two typical solutions with self-Fourier properties. In the right part, we display one solution connecting two distinct partners ψ and φ . Their shapes show distinctive features. One




Fig. 12. Left: two examples of self Fourier states ($\varphi = \psi$). Right: mixed parity case ($\varphi \neq \psi$).

class of shapes, which are monotonic, seem to remain closer to the bare Gaussian, the building block of our method; it always belongs to the subsets we found. The other class, with oscillations, is different. One might ask if our truncations to a maximum order N limit the flexibility of our method to the vicinity of the Gaussian. It seems not to be so, as shown for instance by the very oscillating solution⁴ in the left part of figure 12, which exhibits an approximate periodicity in an interval; it is reminiscent of the Dirac comb, which is, of course, outside the set of functions constructed with a finite number of polynomials. However, from our numerical experience, see in particular figures 2 and 3 and Eqs. (10) and (11), we may risk the conjecture that, when we allow a mixture including $H_0, H_4, \dots H_{4N}$ and when the weight of H_{4N} is that maximum allowed by positivity, see the tiny red triangle in the right part of figure 3, then the corresponding function $\psi = \varphi$ has a Dirac comb limit when $N \rightarrow \infty$. The wavelength of this limit comb is also conjectured to be 1. Combs with different wavelengths may be obtained as limits, but they would not be self-Fourier. For each truncation at a given polynomial order $4N$, the locus of such candidates for comb limits can be defined by projecting the combs into the subspace spanned by $\exp(-r^2/2) H_0, \exp(-r^2/2) H_4, \dots \exp(-r^2/2) H_{4N}$, obviously.

Our bases are flexible enough to reconstruct any function having positivity properties, but in some cases convergence might be slow. It is not excluded that other bases exist, which might be more convenient to speed up the convergence and make easier the search for positivity domains. Another open problem is that of positivity for periodic functions. Such questions are beyond the scope of the present paper.

It is a pleasure to thank R. Balian, R. Enberg, R. Lacaze, C. Marquet, P. Moussa, G. Soyez and A. Voros for stimulating discussions. R.P. thanks T. Lafforgue, (Lycée Blaise Pascal, Orsay), for his fruitful contributions. Thanks are also due to E.H. Lieb and K. Scharnhorst for calling our attention to related works, quoted in the references.

⁴ Its equation reads

$$\begin{aligned} \psi = \exp(-r^2/2) & \left[0.566053 + 0.0488517(3 - 12r^2 + 4r^4) \right. \\ & + 0.0011871(105 - 840r^2 + 840r^4 - 224r^6 + 16r^8) \\ & + 0.0000164538 \left(10395 - 124740r^2 + 207900r^4 - 110880r^6 + 23760r^8 \right. \\ & \left. \left. - 2112r^{10} + 64r^{12} \right) \right]. \end{aligned}$$

REFERENCES

- [1] R.J. Glauber, in *High-Energy Physics and Nuclear Structure*, Proc. of the 2nd International Conference, Rehovoth, 1967, ed. G. Alexander, North-Holland, Amsterdam 1967, p. 311, and the references mentioned therein.
- [2] Y.V. Kovchegov, *Phys. Rev.* **D60**, 034008 (1999); *Phys. Rev.* **D61**, 074018 (2000).
- [3] P. Lévy: Fonctions caractéristiques positives [Positive characteristic functions]. *Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences, Série A, Sciences Mathématiques* **265**, 249 (1967), in French. Reprinted in: *Œuvres de Paul Lévy. Volume III. Éléments Aléatoires*, edited by D. Dugué in collaboration with P. Deheuvels, M. Ibéro, Gauthier-Villars Éditeur, Paris 1976, pp. 607–610.
- [4] E.C. Titchmarsh, *Introduction to the Theory of Fourier Integrals*, Clarendon Press, Oxford 1959.
- [5] M.J. Caola, *J. Phys. A* **24**, L1143 (1991).
- [6] G. Cincotti, F. Gori, M. Santarsiero, *J. Phys. A* **25**, L1191 (1992).
- [7] T. Lafforgue, private communication; for a general theorem, see R. Askey, *Harmonic Analysis on Homogeneous Spaces, Proc. of the Symposia in Pure Math.* Vol. 26, AMS, Providence 1973, pp. 335–338.
- [8] E.H. Lieb, *Phys. Rev. Lett.* **54**, 1987 (1985); Sh. Nussinov, *Phys. Rev. Lett.* **86**, 4762 (2001).
- [9] For a comprehensive list of references, see K. Scharnhorst, *J. Math. Phys.* **44**, 5415 (2003) [[math-ph/0206006](#)], in particular the references [111] to [125]; see also Ch. Pfannschmidt, PhD dissertation and H.-H. Thulke, PhD dissertation, Leipzig Univ. 1996, in German.
- [10] I.M. Gel'fand, N.Ya. Vilenkin, *Generalized Functions*, Vol. IV, Academic Press, New York and London 1968.
- [11] See, for general mathematical properties: J.L. Teugels, *Bull. Soc. Math. Belg.* **23**, 236 (1971); H.-J. Rossberg, *J. Math. Sci. (New York)* **76**, 2181 (1995). This article is part of: Problemy Ustoïchivosti Stokhasticheskikh Modelei [Problems of the Stability of Stochastic Models] — Trudy Seminara (Proceedings of the Seminar on Stochastic Problems), Moscow, 1993. *J. Math. Sci. (New York)* **76**, 2093 (1995); K. Schladitz, H.J. Engelbert, *Teor. Veroyatn. Primen.* **40**, 694 (1995) [*Theor. Probab. Appl.* **40**, 577 (1996)].
- [12] C. Sturm, “Mémoire sur la résolution des équations numériques”, *Bull. Sci. Math. Ferussac* **2**, 419 (1829), in French.