



HAL
open science

Zeros of some bi-orthogonal polynomials

Madan Lal Mehta

► **To cite this version:**

Madan Lal Mehta. Zeros of some bi-orthogonal polynomials. *Journal of Physics A: Mathematical and General* (1975 - 2006), 2002, 35 (3), pp.517-525. 10.1088/0305-4470/35/3/305 . cea-02904735

HAL Id: cea-02904735

<https://cea.hal.science/cea-02904735>

Submitted on 22 Jul 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Zeros of some bi-orthogonal polynomials.

Madan Lal Mehta*

C.E. de Saclay, F-91191 Gif-sur-Yvette Cedex, France

Abstract. Ercolani and McLaughlin have recently shown that the zeros of the bi-orthogonal polynomials with the weight $w(x, y) = \exp[-(V_1(x) + V_2(y) + 2cxy)/2]$, relevant to a model of two coupled hermitian matrices, are real and simple. We show that their argument applies to the more general case of the weight $(w_1 * w_2 * \dots * w_j)(x, y)$, a convolution of several weights of the same form. This general case is relevant to a model of several hermitian matrices coupled in a chain. Their argument also works for the weight $W(x, y) = e^{-x-y}/(x+y)$, $0 \leq x, y < \infty$, and for a convolution of several such weights.

1. Introduction. For a weight function $w(x, y)$ such that all the moments

$$M_{i,j} := \int w(x, y) x^i y^j dx dy \tag{1.1}$$

exist and

$$D_n := \det[M_{i,j}]_{i,j=0,1,\dots,n} \neq 0 \tag{1.2}$$

for all $n \geq 0$, unique monic polynomials $p_n(x)$ and $q_n(x)$ of degree n exist satisfying the bi-orthogonality relations (a polynomial is called monic when the coefficient of the highest degree is one)

$$\int w(x, y) p_n(x) q_m(y) dx dy = h_n \delta_{mn}. \tag{1.3}$$

Just like the orthogonal polynomials they can be expressed as determinants, e.g.

$$p_n(x) = \frac{1}{D_{n-1}} \det \begin{bmatrix} M_{0,0} & \dots & M_{0,n-1} & 1 \\ M_{1,0} & \dots & M_{1,n-1} & x \\ \vdots & \vdots & \vdots & \vdots \\ M_{n,0} & \dots & M_{n,n-1} & x^n \end{bmatrix} \tag{1.4}$$

and have integral representations, e.g.

$$p_n(x) \propto \int \Delta_n(\mathbf{x}) \Delta_n(\mathbf{y}) \prod_{j=1}^n (x - x_j) w(x_j, y_j) dx_j dy_j \tag{1.5}$$

$$\Delta_n(\mathbf{x}) := \prod_{1 \leq i < j \leq n} (x_j - x_i), \quad \Delta_n(\mathbf{y}) := \prod_{1 \leq i < j \leq n} (y_j - y_i). \tag{1.6}$$

* E-mail: mehta@sph.t.saclay.cea.fr

From limited numerical evidence for the weights

- (i) $w(x, y) = \sin(\pi xy)$, $0 \leq x, y \leq 1$;
- (ii) $w(x, y) = |x - y|$, $-1 \leq x, y \leq 1$;
- (iii) $w(x, y) = [1/(x + y)] \exp[-x - y]$, $0 \leq x, y < \infty$;
- (iv) $w(x, y) = \exp(-x^2 - y^2 - cxy)$, $-\infty < x, y < \infty$, $0 < c < 2$;

one might think that the zeros of the bi-orthogonal polynomials are real, simple, lie respectively in the x or y -support of $w(x, y)$, interlace for successive n, \dots

Alas, this is not true in general as seen by the following example due to P. Deligne. If one takes

$$w(x, y) = u(x, y) + v(x, y), \quad (1.7)$$

$$u(x, y) = \begin{cases} \delta(x - y), & -1 \leq x, y \leq 1, \\ 0, & \text{otherwise,} \end{cases} \quad (1.8)$$

$$v(x, y) = \frac{1}{8}[\delta(x - 1)\delta(y + 2) + \delta(x + 1)\delta(y - 2)]. \quad (1.9)$$

Then the zeros of $p_3(x)$ and $q_3(x)$ are complex.

However, N.M. Ercolani and K.T.-R. Mclaughlin have recently [1] shown that with the weight function

$$w_1(x, y) = \exp \left[-\frac{1}{2}V_1(x) - \frac{1}{2}V_2(y) - c_1xy \right] \quad (1.10)$$

($-\infty < x, y < \infty$), V_1 and V_2 polynomials of positive even degree, c a small non-zero real constant, all the zeros of the bi-orthogonal polynomials $p_n(x)$ and $q_n(x)$ are real and simple.

In this brief note we will show that their argument works for the following general case encountered for random hermitian matrices coupled in a linear chain. Let $V_j(x)$, $1 \leq j \leq p$, be polynomials of positive even degree and c_j , $1 \leq j < p$, be small real constants, none of them being zero ("small" so that all the moments $M_{i,j}$ defined below, eq.(1.13), exist.) Further let

$$w_k(x, y) := \exp \left[-\frac{1}{2}V_k(x) - \frac{1}{2}V_{k+1}(y) - c_kxy \right] \quad (1.11)$$

$$(w_{i_1} * w_{i_2} * \dots * w_{i_k})(\xi_1, \xi_{k+1}) := \int w_{i_1}(\xi_1, \xi_2)w_{i_2}(\xi_2, \xi_3)\dots w_{i_k}(\xi_k, \xi_{k+1})d\xi_2\dots d\xi_k \quad (1.12)$$

Moreover, assume that for all $i, j \geq 0$

$$M_{i,j} := \int x^i (w_1 * w_2 * \dots * w_{p-1})(x, y) y^j dx dy \quad (1.13)$$

exist.

Theorem. Then monic polynomials $p_j(x)$ and $q_j(x)$ can be uniquely defined by

$$\int p_j(x)(w_1 * w_2 * \dots * w_{p-1})(x, y)q_k(y)dx dy = h_j \delta_{jk} \quad (1.14)$$

and all the zeros of $p_j(x)$ and of $q_j(x)$ are real and simple.

The same argument works for any weight $W(x, y)$ such that $\det[W(x_i, y_j)]_{i,j=1,\dots,n} > 0$ for $x_1 < x_2 < \dots < x_n$, $y_1 < y_2 < \dots < y_n$ and moments $M_{i,j} = \int W(x, y)x^i y^j dx dy$ exist for all $i, j \geq 0$. For example, if $W(x, y) = [1/(x + y)] \exp[-x - y]$, $0 \leq x, y < \infty$, then monic polynomials $p_j(x)$ can be uniquely defined by

$$\int_0^\infty p_j(x)W(x, y)p_k(y)dx dy = h_j \delta_{jk} \quad (1.15)$$

(here $W(x, y)$ is symmetric in x and y so that $p_j(x) = q_j(x)$) and all the zeros of $p_j(x)$ are real, simple and non-negative.

2. Results and proofs. Here we essentially follow section 3 of reference [1]. With any monic polynomials $p_j(x)$ and $q_j(x)$ of degree j , let us write

$$P_{1,j}(x) := p_j(x) \quad (2.1)$$

$$\begin{aligned} P_{i,j}(x) &:= \int p_j(\xi)(w_1 * w_2 * \dots * w_{i-1})(\xi, x)d\xi \\ &:= \int p_j(\xi)U_{Li}(\xi, x)d\xi, \quad 1 < i \leq p \end{aligned} \quad (2.2)$$

$$Q_{p,j}(x) := q_j(x) \quad (2.3)$$

$$\begin{aligned} Q_{i,j}(x) &:= \int (w_i * w_{i+1} * \dots * w_{p-1})(x, \xi)q_j(\xi)d\xi \\ &:= \int U_{Ri}(x, \xi)q_j(\xi)d\xi \quad 1 \leq i < p \end{aligned} \quad (2.4)$$

Lemma 1. For $x_1 < x_2 < \dots < x_n$, $y_1 < y_2 < \dots < y_n$,

$$\det [w_i(x_j, y_k)]_{j,k=1,\dots,n} > 0. \quad (2.5)$$

This is essentially eq. (40) of reference [1]. This can also be seen as follows. Let $X = [x_i \delta_{ij}]$ and $Y = [y_i \delta_{ij}]$ be two $n \times n$ diagonal matrices with diagonal elements x_1, \dots, x_n and y_1, \dots, y_n respectively. Then the integral of $\exp[-c \operatorname{tr} UXU^{-1}Y]$ over the $n \times n$ unitary matrices U is given by [2]

$$K \frac{\det [\exp(-c x_i y_j)]_{i,j=1,\dots,n}}{\Delta_n(\mathbf{x})\Delta_n(\mathbf{y})} \quad (2.6)$$

where K is a positive constant depending on c and n . Hence

$$\exp \left[-\frac{1}{2} \sum_{j=1}^n (V_i(x_j) + V_{i+1}(y_j)) \right] \int dU e^{-c \operatorname{tr} UXU^{-1}Y} = K \frac{\det [w_i(x_j, y_k)]_{j,k=1,\dots,n}}{\Delta_n(\mathbf{x})\Delta_n(\mathbf{y})} \quad (2.7)$$

The left hand side is evidently positive while on the right hand side the denominator is positive since $x_1 < x_2 < \dots < x_n$ and $y_1 < y_2 < \dots < y_n$. From this eq. (2.5) follows.

Lemma 2. For $x_1 < x_2 < \dots < x_n$, $y_1 < y_2 < \dots < y_n$,

$$\det [(w_{i_1} * w_{i_2} * \dots * w_{i_\ell})(x_j, y_k)]_{j,k=1,\dots,n} > 0 \quad (2.8)$$

Proof. Binet-Cauchy formula tells us that [3]

$$\det [(w_{i_1} * w_{i_2})(x_j, y_k)]_{j,k=1,\dots,n}$$

is equal to

$$\int_{\xi_1 < \xi_2 < \dots < \xi_n} \det [w_{i_1}(x_j, \xi_k)]_{j,k=1,\dots,n} \cdot \det [w_{i_2}(\xi_j, y_k)]_{j,k=1,\dots,n} d\xi_1 \dots d\xi_n \quad (2.9)$$

By lemma 1 the integrand is every where positive, so lemma 2 is proved for the case $\ell = 2$. The proof is now completed by induction on ℓ , using again the Binet-Cauchy formula.

Lemma 3. For any monic polynomial $p_j(x)$ of degree j , $P_{i,j}(x)$, $1 \leq i \leq p$, has at most j distinct real zeros. Similarly, for any monic polynomial $q_j(x)$ of degree j , $Q_{i,j}(x)$, $1 \leq i \leq p$, has at most j distinct real zeros.

Proof. Let, if possible, $z_1 < z_2 < \dots < z_m$, $m > j$, be the distinct real zeros of $P_{i,j}(x)$. Since

$$P_{i,j}(x) = \sum_{k=0}^j a_k T_{i,k}(x), \quad (2.10)$$

with

$$T_{i,k}(x) := \int \xi^k U_{Li}(\xi, x) d\xi, \quad (2.11)$$

$$P_{i,j}(z_\ell) = 0, \quad \ell = 1, 2, \dots, m, \quad m > j \quad (2.12)$$

implies that

$$\begin{aligned} 0 &= \det \begin{bmatrix} T_{i,0}(z_1) & T_{i,1}(z_1) & \dots & T_{i,j}(z_1) \\ \dots & \dots & \dots & \dots \\ T_{i,0}(z_{j+1}) & T_{i,1}(z_{j+1}) & \dots & T_{i,j}(z_{j+1}) \end{bmatrix} \\ &= \int \det \begin{bmatrix} U_{Li}(\xi_1, z_1) & \xi_2 U_{Li}(\xi_2, z_1) & \dots & \xi_{j+1}^j U_{Li}(\xi_{j+1}, z_1) \\ \dots & \dots & \dots & \dots \\ U_{Li}(\xi_1, z_{j+1}) & \xi_2 U_{Li}(\xi_2, z_{j+1}) & \dots & \xi_{j+1}^j U_{Li}(\xi_{j+1}, z_{j+1}) \end{bmatrix} d\xi_1 \dots d\xi_{j+1} \\ &= \int \xi_2 \xi_3^2 \dots \xi_{j+1}^j \det [U_{Li}(\xi_k, z_\ell)]_{k,\ell=1,\dots,j+1} d\xi_1 \dots d\xi_{j+1} \end{aligned} \quad (2.13)$$

or

$$\int \det [U_{Li}(\xi_k, z_\ell)]_{k,\ell=1,\dots,j+1} \cdot \det [\xi_k^{\ell-1}]_{k,\ell=1,\dots,j+1} d\xi_1 \dots d\xi_{j+1} = 0 \quad (2.14)$$

in contradiction to lemma 2. Thus m can not be greater than j .

The proof for $Q_{i,j}(x)$ is similar.

Lemma 4. Let the real constants c_1, \dots, c_{p-1} , none of them being zero, be such that

$$M_{i,j} := \int x^i U_{Lp}(x,y) y^j dx dy \equiv \int x^i (w_1 * w_2 * \dots * w_{p-1})(x,y) y^j dx dy. \quad (2.15)$$

exist for all $i, j \geq 0$. Then

$$D_n := \det[M_{i,j}]_{i,j=1,\dots,n} \neq 0 \quad (2.16)$$

for any $n \geq 0$.

Proof. Let, if possible, $D_n = 0$ for some n . Then $\sum_{j=0}^n M_{i,j} q_j = 0$, q_j not all zero, and

$$\int x^i U_{Lp}(x,y) \sum_{j=0}^n q_j y^j dx dy = 0, \quad i = 0, 1, \dots, n \quad (2.17)$$

or

$$\int p_i(x) U_{Lp}(x,y) \sum_{j=0}^n q_j y^j dx dy = 0 \quad (2.18)$$

for any polynomial $p_i(x)$ of degree $i \leq n$. But

$$\int U_{Lp}(x,y) \sum_{j=0}^n q_j y^j dy \quad (2.19)$$

has at most n distinct real zeros (lemma 3). So one can choose $p_i(x)$ such that

$$p_i(x) \int U_{Lp}(x,y) \sum_{j=0}^n q_j y^j dy > 0 \quad (2.20)$$

in contradiction to eq. (2.18). So $D_n \neq 0$ and bi-orthogonal polynomials $p_j(x)$, $q_j(x)$ exist, see eqs. (1.4), (1.5).

Lemma 5. Let $p_j(x)$, $q_j(x)$ be the bi-orthogonal polynomials, eq. (1.14); or with the definitions (2.1)-(2.4)

$$\int P_{i,j}(x) Q_{i,k}(x) dx = h_j \delta_{jk}, \quad 1 \leq i \leq p \quad (2.21)$$

Then $P_{i,j}(x)$ has at least j real distinct zeros of odd multiplicity. So does have $Q_{i,j}(x)$.

Proof. Let, if possible, $z_1 < z_2 < \dots < z_m$, $m < j$, be the only real zeros of $P_{i,j}(x)$ of odd multiplicity. Set

$$R(x) = \det \begin{bmatrix} Q_{i,0}(x) & Q_{i,1}(x) & \dots & Q_{i,m}(x) \\ Q_{i,0}(z_1) & Q_{i,1}(z_1) & \dots & Q_{i,m}(z_1) \\ \vdots & \vdots & & \vdots \\ Q_{i,0}(z_m) & Q_{i,1}(z_m) & \dots & Q_{i,m}(z_m) \end{bmatrix} \quad (2.22)$$

$$= \int U_{Ri}(x, \xi) \sum_{k=0}^m \alpha_k \xi^k d\xi \quad (2.23)$$

with some constants α_k depending on z_1, \dots, z_m .

Since $m < j$, the bi-orthogonality gives

$$\int P_{i,j}(x)R(x)dx = 0. \quad (2.24)$$

However, $R(x)$ can also be written as

$$\begin{aligned} R(x) &= \int \det \begin{bmatrix} U_{Ri}(x, \xi_0) & U_{Ri}(x, \xi_1)\xi_1 & \dots & U_{Ri}(x, \xi_m)\xi_m^m \\ U_{Ri}(z_1, \xi_0) & U_{Ri}(z_1, \xi_1)\xi_1 & \dots & U_{Ri}(z_1, \xi_m)\xi_m^m \\ \vdots & \vdots & & \vdots \\ U_{Ri}(z_m, \xi_0) & U_{Ri}(z_m, \xi_1)\xi_1 & \dots & U_{Ri}(z_m, \xi_m)\xi_m^m \end{bmatrix} d\xi_0 d\xi_1 \dots d\xi_m \\ &= \int \det \begin{bmatrix} U_{Ri}(x, \xi_0) & U_{Ri}(x, \xi_1) & \dots & U_{Ri}(x, \xi_m) \\ U_{Ri}(z_1, \xi_0) & U_{Ri}(z_1, \xi_1) & \dots & U_{Ri}(z_1, \xi_m) \\ \vdots & \vdots & & \vdots \\ U_{Ri}(z_m, \xi_0) & U_{Ri}(z_m, \xi_1) & \dots & U_{Ri}(z_m, \xi_m) \end{bmatrix} \xi_1 \xi_2^2 \dots \xi_m^m d\xi_0 d\xi_1 \dots d\xi_m \\ &= \frac{1}{(m+1)!} \int \det \begin{bmatrix} U_{Ri}(x, \xi_0) & U_{Ri}(x, \xi_1) & \dots & U_{Ri}(x, \xi_m) \\ U_{Ri}(z_1, \xi_0) & U_{Ri}(z_1, \xi_1) & \dots & U_{Ri}(z_1, \xi_m) \\ \vdots & \vdots & & \vdots \\ U_{Ri}(z_m, \xi_0) & U_{Ri}(z_m, \xi_1) & \dots & U_{Ri}(z_m, \xi_m) \end{bmatrix} \prod_{0 \leq r < s \leq m} (\xi_s - \xi_r) d\xi_0 d\xi_1 \dots d\xi_m \\ &= \int_{\xi_0 < \xi_1 < \dots < \xi_m} \det \begin{bmatrix} U_{Ri}(x, \xi_0) & U_{Ri}(x, \xi_1) & \dots & U_{Ri}(x, \xi_m) \\ U_{Ri}(z_1, \xi_0) & U_{Ri}(z_1, \xi_1) & \dots & U_{Ri}(z_1, \xi_m) \\ \vdots & \vdots & & \vdots \\ U_{Ri}(z_m, \xi_0) & U_{Ri}(z_m, \xi_1) & \dots & U_{Ri}(z_m, \xi_m) \end{bmatrix} \prod_{0 \leq r < s \leq m} (\xi_s - \xi_r) d\xi_0 d\xi_1 \dots d\xi_m \end{aligned} \quad (2.25)$$

Thus $R(x)$ is represented as an integral whose integrand has a fixed sign determined by the relative ordering of the numbers x, z_1, z_2, \dots, z_m (lemma 2). It thus follows that $R(x)$ changes sign when x passes through any of the points $z_k, k = 1, \dots, m$ and at no other value of x . In other words, z_1, \dots, z_m are the only real zeros of $R(x)$ having an odd multiplicity. And therefore $P_{i,j}(x)R(x)$ has a constant sign, so that

$$\int P_{i,j}(x)R(x)dx \neq 0 \quad (2.26)$$

in contradiction to (2.24).

The proof for $Q_{i,j}(x)$ is similar.

As a consequence we have the integral representations of $P_{i,j}(x)$ for $i > 1$ and of $Q_{i,j}(x)$ for $i < p$ involving their respective zeros

$$P_{i,j}(x) \propto \int \det \begin{bmatrix} U_{Li}(\xi_0, x) & U_{Li}(\xi_1, x) & \dots & U_{Li}(\xi_j, x) \\ U_{Li}(\xi_0, z_1) & U_{Li}(\xi_1, z_1) & \dots & U_{Li}(\xi_j, z_1) \\ \vdots & \vdots & & \vdots \\ U_{Li}(\xi_0, z_j) & U_{Li}(\xi_1, z_j) & \dots & U_{Li}(\xi_j, z_j) \end{bmatrix} \prod_{0 \leq r < s \leq j} (\xi_s - \xi_r) d\xi_0 d\xi_1 \dots d\xi_j$$

(2.27)

$$Q_{i,j}(x) \propto \int \det \begin{bmatrix} U_{Ri}(x, \xi_0) & U_{Ri}(x, \xi_1) & \dots & U_{Ri}(x, \xi_j) \\ U_{Ri}(z_1, \xi_0) & U_{Ri}(z_1, \xi_1) & \dots & U_{Ri}(z_1, \xi_j) \\ \vdots & \vdots & & \vdots \\ U_{Ri}(z_j, \xi_0) & U_{Ri}(z_j, \xi_1) & \dots & U_{Ri}(z_j, \xi_j) \end{bmatrix} \prod_{0 \leq r < s \leq j} (\xi_s - \xi_r) d\xi_0 d\xi_1 \dots d\xi_j \quad (2.28)$$

Lemmas 3 and 5 tell us that if $p_j(x)$ and $q_j(x)$ are bi-orthogonal polynomials satisfying eq. (1.14), then $P_{i,j}(x)$ and $Q_{i,j}(x)$ each have exactly j distinct real zeros of odd multiplicity. In particular, the zeros of the bi-orthogonal polynomials $p_j(x) \equiv P_{1,j}(x)$ and $q_j(x) \equiv Q_{p,j}(x)$ are real and simple.

With a little more effort one can perhaps show that all the real zeros of $P_{i,j}(x)$ and of $Q_{i,j}(x)$ are simple. Other zeros, if any, must be complex. Whether the zeros of $p_j(x)$ ($q_j(x)$) interlace for successive j , remains an open question.

3. Bi-orthogonal polynomials with another weight.

For the weight $W(x, y) = [1/(x+y)] \exp[-x-y]$, $0 \leq x, y < \infty$, one can say as follows.

Lemma 1'. One has [4]

$$\det[W(x_j, y_k)]_{j,k=1,\dots,n} = \exp \left[- \sum_{j=1}^n (x_j + y_j) \right] \Delta_n(\mathbf{x}) \Delta_n(\mathbf{y}) \prod_{j,k=1}^n (x_j + y_k)^{-1} \quad (3.1)$$

which is evidently positive for $0 \leq x_1 < x_2 < \dots < x_n$, $0 \leq y_1 < y_2 < \dots < y_n$.

Lemma 3'. For any monic polynomial $p_j(x)$ of degree j , $P_j(x) := \int_0^\infty W(x, y) p_j(y) dy$ has at most j distinct real non-negative zeros.

In the proof of lemma 3, replace eqs. (2.10)-(2.14) by

$$P_j(x) = \sum_{k=0}^j a_k T_k(x), \quad (3.2)$$

$$T_k(x) = \int_0^\infty \xi^k W(\xi, x) d\xi \quad (3.3)$$

$$P_j(z_\ell) = 0, \quad \ell = 1, 2, \dots, m, \quad m > j \quad (3.4)$$

$$\begin{aligned} 0 &= \det \begin{bmatrix} T_0(z_1) & T_1(z_1) & \dots & T_j(z_1) \\ \dots & \dots & \dots & \dots \\ T_0(z_{j+1}) & T_1(z_{j+1}) & \dots & T_j(z_{j+1}) \end{bmatrix} \\ &= \int_0^\infty \det \begin{bmatrix} W(\xi_1, z_1) & \xi_2 W(\xi_2, z_1) & \dots & \xi_{j+1}^j W(\xi_{j+1}, z_1) \\ \dots & \dots & \dots & \dots \\ W(\xi_1, z_{j+1}) & \xi_2 W(\xi_2, z_{j+1}) & \dots & \xi_{j+1}^j W(\xi_{j+1}, z_{j+1}) \end{bmatrix} d\xi_1 \dots d\xi_{j+1} \\ &= \int \xi_2 \xi_3^2 \dots \xi_{j+1}^j \det [W(\xi_k, z_\ell)]_{k,\ell=1,\dots,j+1} d\xi_1 \dots d\xi_{j+1} \quad (3.5) \end{aligned}$$

or

$$\int \det [W(\xi_k, z_\ell)]_{k,\ell=1,\dots,j+1} \cdot \det [\xi_k^{\ell-1}]_{k,\ell=1,\dots,j+1} d\xi_1 \dots d\xi_{j+1} = 0 \quad (3.6)$$

in contradiction to lemma 2'. Thus m can not be greater than j .

Lemma 4'. With

$$M_{i,j} := \int_0^\infty x^i W(x,y) y^j dx dy, \quad (3.7)$$

$$D_n := \det [M_{i,j}]_{i,j=0,1,\dots,n} \neq 0 \quad (3.8)$$

for any $n \geq 0$.

In the proof of lemma 4 replace everywhere $\int U_{Lp}(x,y) \dots$ by $\int_0^\infty W(x,y) \dots$

Lemma 5'. Let $p_j(x)$ be the (bi-orthogonal) polynomials satisfying

$$\int_0^\infty W(x,y) p_j(x) p_k(y) dx dy = h_j \delta_{jk}. \quad (3.9)$$

Then $P_j(x) := \int_0^\infty W(x,y) p_j(y) dy$ and $p_j(x)$ each have at least j distinct real non-negative zeros of odd multiplicity.

Let, if possible, $0 \leq z_1 < z_2 < \dots < z_m$ $m < j$, be the only real non-negative zeros of $P_j(x)$ of odd multiplicity. Set $R(x) = \prod_{j=1}^m (x - z_j)$. Then as $m < j$, one has

$$\int_0^\infty P_j(x) R(x) dx = 0 \quad (3.10)$$

But $P_j(x)$ and $R(x)$ change sign simultaneously as x passes through the values z_1, \dots, z_m and at no other real positive value. So the product $P_j(x)R(x)$ never changes sign, in contradiction to (3.10). Therefore $P_j(x)$ has at least j distinct real non-negative zeros of odd multiplicity.

To prove that $p_j(x)$ has at least j distinct real non-negative zeros let if possible, $0 \leq z_1 < z_2 < \dots < z_m$, $m < j$, be the only such zeros. Set

$$\begin{aligned} R(x) &= \det \begin{bmatrix} P_0(x) & P_1(x) & \dots & P_m(x) \\ P_0(z_1) & P_1(z_1) & \dots & P_m(z_1) \\ \vdots & \vdots & & \vdots \\ P_0(z_m) & P_1(z_m) & \dots & P_m(z_m) \end{bmatrix} \\ &= \int_0^\infty W(x,\xi) \sum_{k=0}^m \alpha_k \xi^k d\xi \end{aligned} \quad (3.11)$$

with some constants α_k depending on z_1, \dots, z_m .

Since $m < j$, the bi-orthogonality gives

$$\int_0^\infty P_j(x) R(x) dx = 0. \quad (3.12)$$

But

$$R(x) \propto \int_0^\infty \det \begin{bmatrix} W(x, \xi_0) & W(x, \xi_1) & \dots & W(x, \xi_m) \\ W(z_1, \xi_0) & W(z_1, \xi_1) & \dots & W(z_1, \xi_m) \\ \vdots & \vdots & & \vdots \\ W(z_m, \xi_0) & W(z_m, \xi_1) & \dots & W(z_m, \xi_m) \end{bmatrix} \prod_{0 \leq r < s \leq m} (\xi_s - \xi_r) d\xi_0 d\xi_1 \dots d\xi_m \quad (3.13)$$

which says that z_1, \dots, z_m are the only distinct real non-negative zeros of $R(x)$ and therefore $p_j(x)R(x)$ has a constant sign, in contradiction to (3.12).

Conclusion. We have shown with the arguments of Ercolani and McLaughlin that if the weight $w(x, y)$ is such that $\det[w(x_i, y_j)]_{i,j=1,\dots,n} > 0$ for $x_1 < x_2 < \dots < x_n$, $y_1 < y_2 < \dots < y_n$ and moments $\int w(x, y)x^i y^j dx dy$ exist for all $i, j \geq 0$, then bi-orthogonal polynomials exist and their zeros are real, simple and lie in the respective supports of the weight $w(x, y)$. The same is true for a weight which is a convolution of several such weights.

Acknowledgements. I am thankful to P.M. Bleher to supply me a copy of the preprint [1].

References.

- [1] N.M. Ercolani and K.T.-R. McLaughlin, Asymptotic and integrable structures for bi-orthogonal polynomials associated to a random two matrix model (preprint, to appear)
- [2] See e.g. M.L. Mehta, *Random matrices*, Academic Press, new York (1991) Appendix A.5
- [3] See e.g. M.L. Mehta, *Matrix theory*, Les Editions de Physique, Z.I. de Courtaboeuf, 91944 Les Ulis Cedex, France, (1989) §3.7
- [4] See e.g. reference 3, §7.1.3