



TrioCFD: code & numerical schemes

Erell Jamelot

► To cite this version:

Erell Jamelot. TrioCFD: code & numerical schemes: E. Jamelot and TrioCFD team. POEMs 2019 - POlytopal Element Methods in Mathematics and Engineering, CIRM (Centre international de rencontres mathématiques), Apr 2019, Marseille, France. cea-02972698

HAL Id: cea-02972698

<https://cea.hal.science/cea-02972698>

Submitted on 20 Oct 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

DE LA RECHERCHE À L'INDUSTRIE



TrioCFD: code & numerical schemes

POEMs 2019

CIRM

E. Jamelot and TrioCFD team

Commissariat à l'Energie Atomique et
aux Energies Alternatives
Centre de Saclay

CEA/DEN/DANS/DM2S/STMF/LMSF

May 2, 2019

Outline

Industrial context

Numerical schemes

Work in progress

Numerical results

Future work

TrioCFD code (previously Trio_U code)

- Developed by the CEA/DEN since the early 1990s.
- Dedicated to unsteady, low Mach number, turbulent flows.
- C++, designed for HPC, open source since 2015.
- Cartesian and triangular/tetrahedral meshes.
- Explicit and semi-implicit time integration.
- Turbulence models: RANS, LES, Wall models.
- Participation in the FVCA 8 benchmark session.
- PhD student: MsFEM (Q. Feng, P. Omnes, G. Allaire).
- Post-docs: ALE (R. Pego, A. Puscas) and sensitivity analysis (C. Fiorini, A. Puscas).
- Website (under evolution): www-trio-u.cea.fr
Download: <https://sourceforge.net/projects/triocfd/>
- Our objectives: support polygonal/polyhedral meshes and high order methods.

Navier-Stokes equations

- Find $(\vec{u}, p) \in \vec{H}_0^1(\Omega) \times L_0^2(\Omega)$ such that:

$$\begin{cases} \partial_t \vec{u} - \nu \Delta \vec{u} + (\vec{u} \cdot \vec{\nabla}) \vec{u} + \vec{\nabla} p &= \vec{f}, \\ \vec{\nabla} \cdot \vec{u} &= 0. \end{cases}$$

- Time discretization (semi-implicit scheme):

1 Prediction step: Compute U^* the solution of

$$\delta t^{-1} M U^* + A U^* + L(U^n) U^* + {}^t B P^n = F^{n+1} + \delta t^{-1} M U^n.$$

At this step, we may have $B U^* \neq 0$.

2 Pressure computation step: Compute P' the solution of

$$B M^{-1} {}^t B P' = \delta t^{-1} B U^*, \quad P^{n+1} = P' + P^n.$$

3 Correction step: Compute U^{n+1} the solution of

$$M U^{n+1} = M U^* - \delta t {}^t B P'.$$

Stokes Problem

- Find $(\vec{u}, p) \in \vec{H}_0^1(\Omega) \times L_0^2(\Omega)$ such that $\begin{cases} -\nu \Delta \vec{u} + \vec{\nabla} p &= \vec{f}, \\ \vec{\nabla} \cdot \vec{u} &= 0. \end{cases}$
- Find $(\vec{u}, p) \in \vec{H}_0^1(\Omega) \times L_0^2(\Omega)$ such that

$$\begin{cases} \forall \vec{v} \in \vec{H}_0^1(\Omega), & a(\vec{u}, \vec{v}) + b(\vec{v}, p) = \langle \vec{f}, \vec{v} \rangle, \\ \forall q \in L_0^2(\Omega), & b(\vec{u}, q) = 0. \end{cases}$$

where $a(\vec{u}, \vec{v}) = (\nu \vec{\nabla} \vec{u}, \vec{\nabla} \vec{v})_0$ and $b(\vec{v}, p) = -(q, \vec{\nabla} \cdot \vec{v})_0$.

Well-posedness: a and b continuous, a coercive on $\vec{H}_0^1(\Omega)$ and the *inf-sup condition* holds

$$\forall q \in L_0^2(\Omega), \quad \sup_{\vec{v} \in \vec{H}_0^1(\Omega)} \frac{b(\vec{v}, q)}{|\vec{v}|_1} \geq \beta \|q\|_0.$$

Stokes Problem

$$(\vec{u}, p) \in \vec{H}_0^1(\Omega) \times L_0^2(\Omega) \mid \begin{cases} \forall \vec{v} \in \vec{H}_0^1(\Omega), & a(\vec{u}, \vec{v}) + b(\vec{v}, p) = \langle \vec{f}, \vec{v} \rangle, \\ \forall q \in L_0^2(\Omega), & b(\vec{u}, q) = 0. \end{cases}$$

Well-posedness: a and b continuous, a coercive on $\vec{H}_0^1(\Omega)$ and the *inf-sup condition* holds

$$\forall q \in L_0^2(\Omega), \quad \sup_{\vec{v} \in \vec{H}_0^1(\Omega)} \frac{b(\vec{v}, q)}{|\vec{v}|_1} \geq \beta \|q\|_0.$$

$$\vec{H}_0^1(\Omega) = \vec{V} \oplus \vec{V}^\perp, \quad \begin{cases} \vec{V} &:= \{\vec{v} \in \vec{H}_0^1(\Omega); \quad \vec{\nabla} \cdot \vec{v} = 0\}, \\ \vec{V}^\perp &= \{(-\Delta)^{-1} \vec{\nabla} q; \quad q \in L^2(\Omega)\}. \end{cases}$$

Abstract tools: [Girault-Raviart'86](#), chap I, cor. 2.4.

1°) The operator $\vec{\nabla}$ is an isomorphism of $L_0^2(\Omega)$ onto \vec{V}^0 such that

$$\vec{V}^0 := \left\{ \vec{y} \in \vec{H}^{-1}(\Omega); \quad \langle \vec{y}, \vec{\phi} \rangle = 0 \quad \forall \vec{\phi} \in \vec{V} \right\}.$$

2°) The operator $\vec{\nabla} \cdot$ is an isomorphism of \vec{V}^\perp onto $L_0^2(\Omega)$.

Space discretizations

- Let \mathcal{T}_h a conforming triangulation of Ω , let \mathcal{F}_h be the set of facets.
- Conforming discretization: $\vec{X}_h \subset \vec{H}_0^1(\Omega)$.

Find $(\vec{u}_h, p_h) \in \vec{X}_h \times M_h$ such that

$$\begin{cases} \forall \vec{v}_h \in \vec{X}_h, & a(\vec{u}_h, \vec{v}_h) + b(\vec{v}_h, p_h) = (\vec{f}, \vec{v}_h)_0, \\ \forall q_h \in M_h, & b(\vec{u}_h, q_h) = 0. \end{cases}$$

Nodal finite element method: Taylor-Hood, conforming Crouzeix-Raviart...

Advantage: Well-posedness of the discrete pb (check ISC).

Drawback: Couplings between the vertices.

- Non-conforming discretization: $\vec{X}_h \not\subset \vec{H}_0^1(\Omega)$.

Find $(\vec{u}_h, p_h) \in \vec{X}_h \times M_h$ such that

$$\begin{cases} \forall \vec{v}_h \in \vec{X}_h, & a_h(\vec{u}_h, \vec{v}_h) + b_h(\vec{v}_h, p_h) = (\vec{f}, \vec{v}_h)_0, \\ \forall q_h \in M_h, & b_h(\vec{u}_h, q_h) = 0. \end{cases}$$

Advantage: Couplings between the facets.

Drawback: Check continuity of a_h , b_h , ellipticity of a_h and ISC.

Space discretizations

- Find $(\vec{u}_h, p_h) \in \vec{X}_h \times M_h$ such that

$$\begin{cases} \forall \vec{v}_h \in \vec{X}_h, & a_{(h)}(\vec{u}_h, \vec{v}_h) + b_{(h)}(\vec{v}_h, p_h) = (\vec{f}, \vec{v}_h)_0, \\ \forall q_h \in M_h, & b_{(h)}(\vec{u}_h, q_h) = 0. \end{cases}$$

- In both cases, the main technical difficulty is that $\vec{V}_h \not\subset \vec{V}$ where:

$$\vec{V}_h := \{\vec{v}_h \in \vec{X}_h \mid \forall q_h \in M_h, \quad b_{(h)}(\vec{v}_h, q_h) = 0\}.$$

- Conforming discretization, a priori error estimates:

$$\begin{cases} \|\vec{u} - \vec{u}_h\|_1 & \lesssim \inf_{\vec{v}_h \in \vec{X}_h} \|\vec{u} - \vec{v}_h\|_1 + \inf_{q_h \in M_h} \|p - q_h\|_0, \\ \|p - p_h\|_0 & \lesssim \inf_{\vec{v}_h \in \vec{X}_h} \|\vec{u} - \vec{v}_h\|_1 + \inf_{q_h \in M_h} \|p - q_h\|_0. \end{cases}$$

- Non-conforming discretization, a priori error estimates:

$$\begin{cases} \|\vec{u} - \vec{u}_h\|_{\vec{X}_h} & \lesssim \inf_{\vec{v}_h \in \vec{X}_h} \|\vec{u} - \vec{v}_h\|_{\vec{X}_h} + \inf_{q_h \in M_h} \|p - q_h\|_0, \\ \|p - p_h\|_{M_h} & \lesssim \inf_{\vec{v}_h \in \vec{X}_h} \|\vec{u} - \vec{v}_h\|_{\vec{X}_h} + \inf_{q_h \in M_h} \|p - q_h\|_0 + \|r_h(u, p)\|_h^*. \end{cases}$$



Non-conforming space discretization

- (\vec{P}_1^{NC}, P_0) proposed in Crouzeix-Raviart'73 $\|\vec{u} - \vec{u}_h\|_0 \lesssim h (|\vec{u}|_1 + \|p\|_0)$.
 $\vec{X}_h := \vec{H}_0^1(\Omega) \cap \vec{P}_1^{NC}$, $M_h := L_0^2(\Omega) \cap P_0$:

$$\begin{aligned}\vec{P}_1^{NC} &:= \{\vec{v}_h \mid \forall T \in \mathcal{T}_h \vec{v}_{h|T} \in (P_1(T))^d \text{ and } \forall f \in \mathcal{F}_h [\vec{v}_h](\vec{x}_f) = 0\}, \\ P_0 &:= \{q_h \mid \forall T \in \mathcal{T}_h, q_h \in \mathbb{R}\}.\end{aligned}$$

$$a_h(\vec{u}_h, \vec{v}_h) = \sum_T (\nu \vec{\nabla} \vec{u}_h, \vec{\nabla} \vec{v}_h)_{0,T}, \quad b_h(\vec{v}_h, q_h) = - \sum_T (\vec{\nabla} \cdot \vec{v}_h, q_h)_{0,T}.$$

Drawback: Spurious discrete-divergence free vectors.



- $p_h \in L_0^2(\Omega) \cap P_0$
- $\vec{u}_h \in \vec{H}_0^1(\Omega) \cap \vec{P}_1^{NC}$

$$\vec{u}_h \cdot \vec{n}|_{\partial T} = 0 \Rightarrow \int_T \vec{\nabla} \cdot \vec{u}_h dT = 0.$$

Non-conforming space discretization

- Less degrees of freedom in \vec{X}_h Hecht'81.

- More degrees of freedom in M_h . In 2D:

- $(\vec{P}_1^{NC}, P_1 + P_b)$ Bernardi-Hecht'00 $M_h = L_0^2(\Omega) \cap P_1 + L_0^2(\Omega) \cap P_b$

$$(\vec{u}, p) \in \vec{H}^{s+1}(\Omega) \times H^s(\Omega) \quad \|\vec{u} - \vec{u}_h\|_{\vec{X}_h} \lesssim h^s (\|\vec{u}\|_{s+1} + \|p\|_s)$$

Drawback: Convergence of the discrete pressure ?

- (\vec{P}_1^{NC}, P_1) Heib'03 $M_h = L_0^2(\Omega) \cap P_1$

$$(\vec{u}, p) \in \vec{H}^{s+1}(\Omega) \times H^s(\Omega) \quad \|\vec{u} - \vec{u}_h\|_{\vec{X}_h} + \|p - p_h\|_0 \lesssim h^s (\|\vec{u}\|_{s+1} + \|p\|_s)$$


Drawback: Spurious discrete divergence-free vectors.

- $(\vec{P}_1^{NC}, P_0 + P_1)$ Heib'03 $M_h = L_0^2(\Omega) \cap P_0 + L_0^2(\Omega) \cap P_1$

$$(\vec{u}, p) \in \vec{H}^2(\Omega) \times H^1(\Omega) \quad \|\vec{u} - \vec{u}_h\|_{\vec{X}_h} + \|p - p_h\|_0 \lesssim h (\|\vec{u}\|_2 + \|p\|_1).$$

($\vec{P}_1^{NC}, P_0 + P_1$) discretization

- $M_h = L_0^2(\Omega) \cap P_0 + L_0^2(\Omega) \cap P_1, \quad P_1 := \{q_h \in \mathcal{C}^0(\Omega) \mid \forall T \in \mathcal{T}_h, q_h \in P_1(T)\}$
- $b_h(\vec{v}_h, q_{h,0} + q_{h,1}) = - \sum_T \left(\vec{\nabla} \cdot \vec{v}_h, q_{h,0} \right)_{0,T} + \sum_T \left(\vec{v}_h, \vec{\nabla} q_{h,1} \right)_{0,T}.$



$\forall \vec{v}_h \in \vec{V}_h :$

$$\vec{\nabla} \cdot \vec{v}_{h|T} = 0,$$

$$\sum_T \left(\vec{\nabla} q_{h,2}, \vec{v}_h \right)_{0,T} = 0 \quad \forall q_{h,2} \in P_2,$$


$$\sum_T \int_{\partial T} q_{h,2} \vec{v}_h \cdot \vec{n} d\sigma = 0 \quad \forall q_{h,2} \in P_2.$$

$$P_2 := \{q_{h,2} \in \mathcal{C}^0(\Omega) \mid \forall T \in \mathcal{T}_h, q_{h,2} \in P_2(T)\}$$

- $p_h \in L_0^2(\Omega) \cap P_0 + L_0^2(\Omega) \cap P_1$
- $\vec{u}_h \in \vec{H}_0^1(\Omega) \cap \vec{P}_1^{NC}$

($\vec{P}_1^{NC}, P_0 + P_1$) discretization

- $M_h = L_0^2(\Omega) \cap P_0 + L_0^2(\Omega) \cap P_1, \quad P_1 := \{q_h \in \mathcal{C}^0(\Omega) \mid \forall T \in \mathcal{T}_h, q_h \in P_1(T)\}$
- $b_h(\vec{v}_h, q_{h,0} + q_{h,1}) = - \sum_T \left(\vec{\nabla} \cdot \vec{v}_h, q_{h,0} \right)_{0,T} + \sum_T \left(\vec{v}_h, \vec{\nabla} q_{h,1} \right)_{0,T}.$



Proof based on Simpson's rule,
exact for order 3 polynomials on a segment:


$$\int_e q_{h,3} d\sigma = \frac{|e|}{6} (q_{h,3}(S_{1,e}) + 4q_{h,3}(M_e) + q_{h,3}(S_{2,e}))$$

- $p_h \in L_0^2(\Omega) \cap P_0 + L_0^2(\Omega) \cap P_1$
- $\vec{u}_h \in \vec{H}_0^1(\Omega) \cap \vec{P}_1^{NC}$

Non-conforming space discretization

- More degrees of freedom in M_h . In 3D:

▷ $(\vec{P}_1^{NC}, (P_0 + P_1 + P_e))$ T. Fortin'06.



▷ Numerical integration scheme on a triangle, exact for order 3 polynomials:


$$\int_T q_{h,3} dT = \frac{|T|}{5} \left(\sum_{i=1}^3 \frac{1}{4} q_{h,3}(S_{i,T}) + \frac{2}{3} \sum_{i=1}^3 q_{h,3}(M_{i,T}) + \frac{3}{4} q_{h,3}(G_T) \right)$$

▷ Numerous degrees of freedom.

Non-conforming space discretization

- More degrees of freedom in M_h . In 3D:

▷ $(\vec{P}_1^{NC}, (P_0 + P_1 + P_e))$ T. Fortin'06.



▷ $M_h^e := \{q_h \in \mathcal{C}^0(\Omega) \mid \forall T \in \mathcal{T}_h, q_h \in P_e(T)\}.$

For $e = [S_i S_j]$, $\gamma_{e|T} := 4 \lambda_{i,T} \lambda_{j,T}.$


▷ $M_h = ((M_h^e/M_h^{1,p} + M_h^1) \cap L_0^2(\Omega) + M_h^0 \cap L_0^2(\Omega))$

▷ For Stokes, if $\vec{f} = \vec{\nabla} p$, where $p \in H^4(\Omega) \cap L_0^2(\Omega)$ and $|p|_4 \neq 0$, then

$$\|\vec{u}_h\|_{\vec{X}_h} \lesssim h^5 \quad \text{and} \quad \|p_h - p\|_0 \lesssim h^2.$$

TrioCFD: a *finite volume-element* code.

- Control volumes of the FVM. [Emonot'92](#)



- FVM: the mass matrix is diagonal in 2D and 3D.

FEM: the mass matrix is diagonal in 2D only.

In TrioCFD code, the mass matrix is diagonal in 2D and 3D as in FVM.

- FVM and FEM: Diffusion matrix is the same.
- FVM and FEM: Coupling matrix is the same.
- FVM or FEM: Convection schemes (centered, upwind, muscl, stab)
- FVM and FEM: Source term is different ($\approx h^2$).

In TrioCFD code, the discrete source term is coded as in FEM.

Helmholtz decomposition

- $\vec{L}^2(\Omega)$ -Helmholtz decomposition and $\vec{L}^2(\Omega)$ -Helmholtz projector (Ω connected):

$$\forall \vec{f} \in \vec{L}^2(\Omega), \quad \exists (\psi, \vec{w}) \in H^1(\Omega) \times \vec{H}_0(\text{div}0, \Omega) \mid \vec{f} = \vec{\nabla} \psi + \vec{w},$$

$$\vec{L}^2(\Omega) = \vec{H}^\perp \oplus \vec{H} : \quad (\vec{\nabla} \psi, \vec{w})_0 = -(\psi, \vec{\nabla} \cdot \vec{w})_0 = 0.$$

$$\mathbb{P} : \vec{L}^2(\Omega) \rightarrow \vec{L}^2(\Omega), \quad \mathbb{P}(\vec{f}) = \vec{w}.$$

- Navier-Stokes variational formulation:

Find $(\vec{u}, p) \in \vec{H}_0^1(\Omega) \times L_0^2(\Omega)$ s.t. $\forall (\vec{v}, q) \in \vec{H}_0^1(\Omega) \times L_0^2(\Omega)$

$$\begin{aligned} \frac{d}{dt}(\vec{u}, \vec{v})_0 + (\nu \vec{\nabla} \vec{u}, \vec{\nabla} \vec{v})_0 + \left((\vec{u} \cdot \vec{\nabla}) \vec{u}, \vec{v} \right)_0 - (p, \vec{\nabla} \cdot \vec{v})_0 &= (\vec{f}, \vec{v})_0, \\ (q, \vec{\nabla} \cdot \vec{u})_0 &= 0. \end{aligned}$$

- $\vec{u} = \mathbb{P}(\vec{u}), \quad \mathbb{P}(\vec{\nabla} \vec{u}) = 0.$

Helmholtz decomposition

- $\vec{L}^2(\Omega)$ -Helmholtz decomposition and $\vec{L}^2(\Omega)$ -Helmholtz projector (Ω connected):

$$\forall \vec{f} \in \vec{L}^2(\Omega), \quad \exists (\psi, \vec{w}) \in H^1(\Omega) \times \vec{H}_0(\text{div}0, \Omega) \mid \vec{f} = \vec{\nabla} \psi + \mathbb{P}(\vec{f}),$$

$$\vec{L}^2(\Omega) = \vec{H}^\perp \oplus \vec{H} : \quad \left(\vec{\nabla} \psi, \mathbb{P}(\vec{f}) \right)_0 = - \left(\psi, \mathbb{P}(\vec{f}) \right)_0 = 0.$$

- Divergence-free momentum balance: $\forall \vec{v} \in \vec{V}$,

$$\begin{aligned} \frac{d}{dt} (\vec{u}, \vec{v})_0 + \left(\nu \vec{\nabla} \vec{u}, \vec{\nabla} \vec{v} \right)_0 + \left(\mathbb{P}((\vec{\nabla} \cdot \vec{u}) \vec{u}), \vec{v} \right)_0 &= \left(\mathbb{P}(\vec{f}), \vec{v} \right)_0, \\ &= \left(\mathbb{P}(\vec{f} + \vec{\nabla} \phi), \vec{v} \right)_0, \end{aligned}$$

- *Irrational* momentum balance: $\forall \vec{v}^\perp \in \vec{V}^\perp$,

$$\frac{d}{dt} (\vec{u}, \vec{v}^\perp)_0 - \left(p, \vec{\nabla} \cdot \vec{v}^\perp \right)_0 + \left((\vec{\nabla} \cdot \vec{u}) \vec{u}, \vec{v}^\perp \right)_0 = \left(\vec{f}, \vec{v}^\perp \right)_0.$$

- Possible discrete biorthogonalization.

Raviart-Thomas FE

- $\vec{H}(\text{div}, \Omega)$ -conforming FE Raviart-Thomas'77:

$$\vec{X}_h^{RT_0} := \{\vec{v}_h \in \vec{H}(\text{div}, \Omega) : \forall T \in \mathcal{T}_h, \quad \vec{v}_{h|T}(\vec{x}) = \vec{a}_T + b_T \vec{x}\}$$

- Raviart-Thomas projection operator (lowest order):

$$\Pi^{RT_0}(\vec{v}_{h|T})(\overrightarrow{OM}_i) \cdot \vec{n}_i = |F_i|^{-1} \int_{F_i} \vec{v}_h \cdot \vec{n}_i \, d\vec{x}, \quad \forall i \in \{1, \dots, d+1\}.$$

- $\forall \vec{v}_h \in \vec{X}_h, \quad \vec{\nabla} \cdot \vec{v}_{h|T} = \vec{\nabla} \cdot (\Pi^{RT_0} \vec{v}_{h|T}).$

- Let $(\vec{\phi}_i^\alpha)_{i,\alpha}$ be the basis of \vec{X}_h :

For $i \in \{1, \dots, N_f\}$, $\alpha \in \{1, \dots, d\}$, $\vec{\phi}_{i|T}^\alpha := (1 - d\lambda_{i|T})\vec{e}_\alpha$. One can show that:

$$\Pi^{RT_0} \left(\vec{\phi}_{i|T}^\alpha \right) = \frac{\vec{S}_{i|T} \cdot \vec{e}_\alpha}{d|T|} \left(\vec{x} - \overrightarrow{OS}_{i|T} \right).$$

Raviart-Thomas projection

- Towards a better consistency [Linke et al](#): Use Raviart-Thomas projection.
- Source term:

$$\forall \vec{v}_h \in \vec{P}_1^{NC} : \begin{cases} (\vec{f}, \vec{v}_h)_0 \neq - \sum_T (\psi, \vec{\nabla} \cdot \vec{v}_h)_{0,T} + (\vec{w}, \vec{v}_h)_0, \\ (\vec{f}, \Pi^{RT_0} \vec{v}_h)_0 = - \sum_T (\psi, \vec{\nabla} \cdot \vec{v}_h)_{0,T} + (\vec{w}, \Pi^{RT_0} \vec{v}_h)_0. \end{cases}$$

- Convection term: $(\vec{u} \cdot \vec{\nabla}) \vec{u} = (\vec{\nabla} \times \vec{u}) \times \vec{u} + \frac{1}{2} \vec{\nabla} (|\vec{u}|^2)$

$$((\vec{u}^{n-1} \cdot \vec{\nabla}) \vec{u}^n, \vec{v})_0 = (\vec{\nabla} \times \vec{u}^n, \vec{u}^{n-1} \times \vec{v})_0 - \frac{1}{2} (\vec{u}^n \cdot \vec{u}^{n-1}, \vec{\nabla} \cdot \vec{v})_0.$$

$$\sum_T \vec{\nabla} \times \vec{u}_{h|T}^n \cdot \int_T \vec{u}_h^{n-1} \times \vec{v}_h dT \rightarrow \sum_T \vec{\nabla} \times \vec{u}_{h|T}^n \cdot \int_T \Pi^{RT_0} \vec{u}_h^{n-1} \times \Pi^{RT_0} \vec{v}_h dT$$

- Mass term: $(\vec{u}_h, \vec{v}_h)_0 \rightarrow (\Pi^{RT_0} \vec{u}_h, \Pi^{RT_0} \vec{v}_h)_0.$

Numerical results in 2D: Stokes equation, no flow

- $\Omega = (0, 1)^2$, $\nu = 1$, $\vec{f} = {}^t(0, Ra(1 - y + 3y^2))$, $Ra > 0$.

$$\vec{u} = \vec{0}, \quad p = Ra \left(y^3 - \frac{y^2}{2} + y - \frac{7}{12} \right)$$

- Values of $\|\vec{u}_h\|_0$ for $Ra = 1$

$1/h$	$\vec{P}_1^{NC} - P_0$	$\vec{P}_1^{NC} - P_0 + \Pi^{RT_0}$	$\vec{P}_1^{NC} - P_1$	$\vec{P}_1^{NC} - P_0 P_1$
20	$2.74 e - 04$	$9.01 e - 13$	$1.21 e - 04$	$1.12 e - 08$
40	$6.53 e - 05$	$1.30 e - 13$	$1.39 e - 05$	$6.97 e - 10$
80	$1.66 e - 05$	$4.44 e - 13$	$1.77 e - 06$	$4.56 e - 11$
160	$4.13 e - 06$	$4.41 e - 13$	$2.21 e - 07$	$2.82 e - 12$
τ	2	—	3	4

- Values of $\|\vec{u}_h\|_0$ for $Ra = 100$

$1/h$	$\vec{P}_1^{NC} - P_0$	$\vec{P}_1^{NC} - P_0 + \Pi^{RT_0}$	$\vec{P}_1^{NC} - P_1$	$\vec{P}_1^{NC} - P_0 P_1$
20	$2.74 e - 02$	$9.01 e - 13$	$1.21 e - 06$	$1.12 e - 06$
40	$6.53 e - 03$	$1.30 e - 13$	$1.39 e - 07$	$6.97 e - 08$
80	$1.66 e - 03$	$4.44 e - 13$	$1.77 e - 08$	$4.56 e - 09$
160	$4.13 e - 04$	$4.41 e - 13$	$2.21 e - 09$	$2.82 e - 10$
τ	2	—	3	4

Numerical results in 2D: Navier-Stokes equation, stationary vortex

- $\Omega = (-1, 1)^2, \nu = 1, \vec{f} = (Re - 1)^t(x, y), Re > 0.$

$$\vec{u} = {}^t(-y, x), \quad p = Re \left(\frac{x^2 + y^2}{2} - \frac{4}{3} \right)$$

- Values of $\|\vec{u} - \vec{u}_h\|_0$ for $Re = 1$

$1/h$	$\vec{P}_1^{NC} - P_0$	$\vec{P}_1^{NC} - P_0 + \Pi^{RT_0}$	$\vec{P}_1^{NC} - P_0 P_1$
20	$1.32 e - 04$	$8.80 e - 12$	$3.36 e - 05$
40	$3.00 e - 05$	$1.27 e - 11$	$7.52 e - 06$
80	$7.86 e - 06$	$3.70 e - 11$	$2.17 e - 06$
160	$1.92 e - 06$	$1.25 e - 11$	$5.04 e - 07$
τ	2	—	4

- Values of $\|\vec{u} - \vec{u}_h\|_0$ for $Re = 10$

$1/h$	$\vec{P}_1^{NC} - P_0$	$\vec{P}_1^{NC} - P_0 + \Pi^{RT_0}$	$\vec{P}_1^{NC} - P_0 P_1$
20	$8.97 e - 04$	$8.77 e - 11$	$3.36 e - 05$
40	$2.01 e - 04$	$1.26 e - 10$	$7.52 e - 06$
80	$5.29 e - 05$	$3.70 e - 10$	$2.17 e - 06$
160	$1.12 e - 05$	$1.25 e - 10$	$5.04 e - 07$
τ	2	—	4

Numerical results in 3D: Navier-Stokes equation, stationary vortex

- $\Omega = (0, 1)^3$, $\nu = 1$, $\vec{f} = (Re - 1)^t(0, 2y - 1, 2z - 1)$.

$$\vec{u} = {}^t(0, -2z + 1, 2y - 1), \quad p = Re \left(\frac{(2y - 1)^2 + (2z - 1)^2}{2} - \frac{1}{3} \right)$$

- Values of $\|\vec{u} - \vec{u}_h\|_0$ for $Re = 1$

# DoFs	$\vec{P}_1^{NC} - P_0$	$\vec{P}_1^{NC} - P_0 + \Pi^{RT_0}$	$\vec{P}_1^{NC} - P_0 P_1$
$7.71e + 03$	$1.00 e - 03$	$3.22 e - 05$	$6.08 e - 03$
$1.53e + 04$	$5.62 e - 03$	$1.77 e - 05$	$3.80 e - 03$
$3.05e + 04$	$3.16 e - 04$	$8.77 e - 06$	$2.45 e - 03$
$6.11e + 04$	$1.78 e - 04$	$4.42 e - 06$	$1.72 e - 03$
τ	2.5	2.8	1.7

- Values of $\|\vec{u} - \vec{u}_h\|_0$ for $Re = 100$

# DoFs	$\vec{P}_1^{NC} - P_0$	$\vec{P}_1^{NC} - P_0 + \Pi^{RT_0}$	$\vec{P}_1^{NC} - P_0 P_1$
$7.71e + 03$	$1.02 e - 01$	$3.19 e - 05$	$6.10 e - 03$
$1.53e + 04$	$5.58 e - 01$	$1.75 e - 05$	$3.83 e - 03$
$3.05e + 04$	$3.15 e - 02$	$8.75 e - 06$	$2.43 e - 03$
$6.11e + 04$	$1.76 e - 02$	$4.39 e - 06$	$1.74 e - 03$
τ	2.5	2.8	1.7

Fortin-Soulie with Π^{RT_1}

- Matlab tests (Stokes, 2D) and Fortin-Soulie'86 with M. Rihani (M2 internship last year).
- No flow in $\Omega = (0, 1)^2$ with $\vec{f} = \vec{\nabla}(x^3)$ ($h = 0.05$, $N_{\vec{u}} = 6486$, $N_p = 3152$):

	$\ \vec{u} - \vec{u} - h\ _0$	$\ \vec{u} - \vec{u}_h\ _h$	$\ p - p_h\ _h$
$F - S$	$2.6e-07$	$3.7e-05$	$2.9e-03$
$F - S + \Pi^{RT_1}$	$9.7e-19$	$3.3e-17$	$2.9e-03$

- Vortex in $\Omega = (0, 1)^2$ with varying ν ($h = 0.1$, $N_{\vec{u}} = 1534$, $N_p = 726$):

$$\vec{u} = \overrightarrow{\operatorname{curl}}(x^2(1-x)^2y^2(1-y^2)), \quad p = x^3 + y^3 - 1/2, \quad \vec{f} = -\nu \Delta \vec{u} + \vec{\nabla} p.$$

	ν	$\ \vec{u} - \vec{u} - h\ _0$	$\ \vec{u} - \vec{u}_h\ _h$	$\ p - p_h\ _h$
$F - S$	$1.0e-0$	$1.5e-5$	$9.2e-4$	$2.1e-3$
	$1.0e-2$	$2.8e-4$	$2.0e-2$	$1.9e-3$
	$1.0e-4$	$9.4e-2$	$1.9e-0$	$1.9e-3$
$F - S + \Pi^{RT_1}$	ν	$\ \vec{u} - \vec{u} - h\ _0$	$\ \vec{u} - \vec{u}_h\ _h$	$\ p - p_h\ _h$
	$1.0e-0$	$1.7e-5$	$1.1e-4$	$2.8e-3$
	\vdots	\vdots	\vdots	\vdots
	$1.0e-8$	$1.7e-5$	$1.1e-4$	$2.8e-3$

Outlook

- **Ongoing work:**

Using Raviart-Thomas projection:

- ▷ Convective form: $\left((\Pi^{RT_0}(\vec{u}_h^{n+1}) \cdot \vec{\nabla}) \vec{u}_h^n, \Pi^{RT_0}(\vec{v}_h) \right)_0$.
- ▷ Unsteady state with $\Pi^{RT_0}(M)$.

Splines for INS using GeoPDEs with L. Dray, M2 student.

- **Future work:**

Handle polyhedral meshes in particular prism or hexahedra and tetrahedra.

- ▷ PolyMAC code (A. Gershfeld et al).

Order 2 method.

- ▷ Preliminary work with M. Rihani, former M2 student.

Matlab tests for Stokes in 2D and **Fortin-Soulie'86** with Π^{RT_1} .

Thank you for your attention!