



Lithium-ion capacitor safety assessment under electrical abuse tests based on ultrasound characterization and cell opening

L. Oca, Nicolas Guillet, R. Tessard, U. Iraola

► To cite this version:

L. Oca, Nicolas Guillet, R. Tessard, U. Iraola. Lithium-ion capacitor safety assessment under electrical abuse tests based on ultrasound characterization and cell opening. *Journal of Energy Storage*, 2019, 23, pp.29-36. 10.1016/j.est.2019.02.033 . cea-03214997

HAL Id: cea-03214997

<https://cea.hal.science/cea-03214997>

Submitted on 21 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

The sexual dimorphism of anticardiolipin autoantibodies in acute Q fever patients

Cléa Melenotte^{*1}, Laetitia Gay^{*1}, Soraya Mezouar¹, Nathalie Bardin², Didier Raoult¹, Jean-

Louis Mège^{1,2}

(1) Aix-Marseille Univ, IRD, APhM, MEPHI, IHU-Méditerranée Infection, Marseille,
France

(2) Laboratoire d'immunologie, Hôpital de la Conception, Marseille, France

* the authors equally contributed to the writing of the manuscript

Corresponding author contact: melenotteclea@gmail.com

Running title: Acute Q fever endocarditis a new clinical entity

Key words: Q fever, *Coxiella burnetii*, anticardiolipin antibodies, sex, gender, autoimmunity

References: 15

Word count of text: 1075

Word count of abstract: 165

Abstract

Objectives : Q fever is a zoonotic disease caused by *C. burnetii* which affects men more than women (sex ratio men/women: 2.2). Acute Q fever complications are associated with elevation of anticardiolipin (aCL) antibodies. Here, we investigate the sexual dimorphism of aCL antibodies during acute *C. burnetii* infection.

Methods : IgG aCL antibodies were evaluated at the time of Q fever serological diagnosis with enzyme-linked immunosorbent assay. Results were analysed according to sex.

Results : Among the 1,323 patients with Q fever tested for aCL, 1,013 had acute Q fever (692 men/321 women) and 310 had persistent focalized infection (226 men/84 women). In case of acute Q fever, men presented a significantly higher proportion of positive aCL antibodies (351/692, 50.7%) than women (113/321, 35.2%) ($p<0.05$). In addition, men had significantly higher aCL antibodies levels than women ($p<0.0001$).

Conclusions : We highlight a relationship between sex and markers of autoimmunity during Q fever. Further investigations are necessary to better understand the mechanisms of this sexual dimorphism.

Key words: Q fever, *Coxiella burnetii*, anticardiolipin antibodies, sex, gender, autoimmunity

Introduction

Viral and bacterial infections have been reported to be associated with autoimmune clinico-biological manifestations [1]. The elevation of anticardiolipin (aCL) antibodies has been described in cases of human immunodeficiency virus (HIV), hepatitis B virus, hepatitis C virus, malaria, syphilis, leprosy, leptospirosis and Q fever [2]. In case of Q fever, the elevation of aCL antibodies has recently been reported as predictive for acute Q fever complications, such as acute Q fever endocarditis, haemophagocytic syndrome, alithiasic cholecystitis and thrombosis [3,4]. Interestingly, an unbalanced sex ratio has been described during *C. burnetii* infection, the disease affecting men more frequently than women (sex ratio men/women: 2.2). Here, we investigated the sexual dimorphism of autoantibodies during *C. burnetii* infection with a specific focus on aCL antibodies.

Methods

Between 1991 and 2016, 2,434 patients with *C. burnetii* infection were followed in the French national reference centre for Q fever. Acute Q fever was diagnosed by the association of acute clinical symptoms (fever, hepatitis and/or pneumonia) with the following serologic criteria: IgG titers representing phase II (200) and IgM titers representing phase II (50) or seroconversion within 3 months of the primary symptoms [5]. Persistent *C. burnetii* infection was defined by the persistence of clinical symptoms for more than 3 months in addition to a positive microbiological criteria and to the identification of an infectious focus [5]. 1,323 patients were tested for IgG aCL antibodies assessed by enzyme-linked immunosorbent assay at first serological diagnosis. A cut-off threshold of 22 IgG anti-phospholipid-binding units (GPLU) was used to define positive samples. The study was approved by the local ethics committee of the IHU (Institut Hospitalo-Universitaire)-Méditerranée Infection under the registration number 12-016 [4].

Statistical analyses were performed using Prism 6.0 (Graphpad Software Inc.) and SPSS 22 Statistics Software. Multivariate models were adjusted for sex and age at baseline [4]. $P < 0.05$ was considered as significant.

Results

Among the 1,323 patients tested for aCL antibodies, 1,013 patients had acute Q fever and 310 had persistent *C. burnetii* focalized infection (Figure 1). Nine hundred and eighteen patients were men (sex ratio men/women: 2.26). The sexual dimorphism was more pronounced in cases of persistent focalized *C. burnetii* infection (sex ratio: 2.27) than in cases of acute Q fever (sex ratio: 2.16) ($p = 0.07$). Eighty four patients with acute Q fever evolved towards a persistent *C. burnetii* infection, 70 were men (83.3%), 14 were women (16.7%), and the sex ratio was 5.

The aCL antibodies levels were significantly higher in cases of acute Q fever (mean 88.2 ± 14 GPLU and median 18 GPLU) than in cases of persistent *C. burnetii* infection (mean 14.2 ± 2.4 GPLU and median 7 GPLU) (Figure 2A, $p < 0.0001$). In addition, the proportion of positive aCL antibodies were significantly higher in cases of acute Q fever (464/1,013, 45.8%) than in cases of persistent *C. burnetii* infection (32/310, 10.3%) ($p < 0.05$). In cases of acute Q fever, men had significantly higher aCL antibodies levels (mean 113.3 ± 20.3 GPLU and median 24 GPLU) than women (mean 50 ± 7 GPLU and median 11 GPLU) (Figure 2B, $p < 0.0001$) and men presented a significantly higher proportion of positive aCL antibodies (351/692, 50.7%) compared to women (113/321, 35.2%) ($p < 0.05$) for all age groups (Figure 2C). In men, higher levels of aCL antibodies were observed in almost all age groups (Figure 2D). Being a male was associated with a significant risk of positive aCL antibodies, regardless of age (OR=1.6; [95% CI, 1.3-3.2], $p < 0.001$) [4].

Acute Q fever hepatitis affected men more than women ($p=0.007$) and was associated with a significant increase in aCL in men ($p=0.047$). In case of acute Q fever, being male was a risk factor for acute Q fever to progress to persistent cardiovascular *C. burnetii* infection (OR=2.14; [95% CI, 1.17-3.91], $p=0.013$).

Discussion

Here, we report that aCL antibody levels are higher in men than in women with acute *C. burnetii* infection. This is distinct from the sexual dimorphism observed in the antiphospholipid syndrome (APS), in which women are more affected by the disease than men (sex ratio men/women: 1:5) [6]. In addition, the aCL antibodies sexual dimorphism observed in association with *C. burnetii* infection is different from that observed in association with HIV infection, in which aCL antibodies are described as significantly higher in women compared to men [7].

Our data support the possibility of a hormonal influence on the expression of autoantibodies in case of *C. burnetii* infection. It is known that women are described to be more prone to autoimmune diseases than men, and this increased susceptibility disappears after menopause [8]. A positive correlation has been shown between elevated aCL antibodies levels and high estradiol serum concentration in patients with systemic lupus erythematosus (SLE) and rheumatoid arthritis [9,10]. In addition, hyper-oestrogenic levels in premenopausal SLE women are associated with an increased risk of APS and cardiovascular manifestations [9]. In gonadectomized or intact non-autoimmune male and female C57BL/6 mice, oestrogen treatment induces the expression of aCL autoantibodies [11]. It has been hypothesized that oestrogens act as enhancers of cell proliferation and stimulate humoral B cell response [12].

In addition, the influence of testosterone should not be neglected. While oestrogen increases the autoantibody production, androgens decrease it [12]. Men with hypogonadism

are at risk of developing autoimmune diseases of which the Klinefelter syndrome is a glaring example. This syndrome has been associated with SLE, progressive systemic sclerosis, polymyositis, mixed connective tissue disease and with an increased frequency of aCL antibody positivity [13]. Testosterone deficiency may promote autoantibody formation, however, in cases of *C. burnetii* infection, testosterone rather appears to favour aCL autoantibodies production [4].

Other mechanisms such as chromosomal influence could be involved. In transgenic SJL mice, the XX sex chromosome was compared with XY, and showed greater susceptibility to autoimmune encephalomyelitis and SLE [14]. Women with Turner's syndrome were at significantly increased risk of autoimmune thyroid diseases and type 1 diabetes mellitus [15].

As limitations of this study, we have no data on patient's hormonal serum levels, we have not studied the chromosomal influence on autoantibody secretion and we did not consider the gender impact, i.e. gender-related social habits that could influence the results.

In conclusion, we highlight a relationship between sex and markers of autoimmunity during Q fever. We noted that acute Q fever induces a higher autoimmune response in men compared to women. However, the mechanisms of sexual dimorphism are multifactorial and although age was taken into account, other factors such as hormone levels and chromosomal impact should be included in future analyses.

128 **Transparency declaration**

129

130 **No funding source**

131 **No relationship with industry**

132 **No conflict of interest**

133

134 **Funding source:**

135 This work was supported by the French Government under the “investissement d’avenir”

136 (“investments for the future”) program managed by the Agence Nationale de la Recherche

137 (ANR, fr: National Agency for research), (reference: Méditerranée infection 10-IAHU-03).

138 This work was also supported by Région Provence-Alpes-Côte d’Azur and European funding

139 FEDER PRIMMI (Fonds Européen de Développement Régional - Plateformes de Recherche

140 et d'Innovation Mutualisées Méditerranée Infection).

141

142 References

- 143 [1] Konstantinov KN, Ulff-Møller CJ, Tzamaloukas AH. Infections and antineutrophil
144 cytoplasmic antibodies: Triggering mechanisms. *Autoimmun Rev* 2015;14:201–3.
145 doi:10.1016/j.autrev.2014.10.020.
- 146 [2] Sène D, Piette J-C, Cacoub P. Antiphospholipid antibodies, antiphospholipid syndrome
147 and infections. *Autoimmun Rev* 2008;7:272–7. doi:10.1016/j.autrev.2007.10.001.
- 148 [3] Million M, Bardin N, Bessis S, Nouiakh N, Douliery C, Edouard S, et al. Thrombosis
149 and antiphospholipid antibody syndrome during acute Q fever. *Medicine (Baltimore)*
150 2017;96. doi:10.1097/MD.00000000000007578.
- 151 [4] Melenotte C, Protopopescu C, Million M, Edouard S, Carrieri MP, Eldin C, et al.
152 Clinical features and complications of *Coxiella burnetii* infections from the French
153 National Reference Center for Q Fever. *JAMA Netw Open* 2018;1:e181580–e181580.
154 doi:10.1001/jamanetworkopen.2018.1580.
- 155 [5] Eldin C, Mélenotte C, Mediannikov O, Ghigo E, Million M, Edouard S, et al. From Q
156 fever to *Coxiella burnetii* infection: A paradigm change. *Clin Microbiol Rev*
157 2017;30:115–90. doi:10.1128/CMR.00045-16.
- 158 [6] Cervera R, Piette J-C, Font J, Khamashta MA, Shoenfeld Y, Camps MT, et al.
159 Antiphospholipid syndrome: Clinical and immunologic manifestations and patterns of
160 disease expression in a cohort of 1,000 patients. *Arthritis Rheum* 2002;46:1019–27.
161 doi:10.1002/art.10187.
- 162 [7] Abdollahi A, Morteza A. Serum concentrations of antiphospholipid and anticardiolipin
163 antibodies are higher in HIV-infected women. *Rheumatol Int* 2012;32:1927–32.
164 doi:10.1007/s00296-011-1859-1.
- 165 [8] Klein SL, Flanagan KL. Sex differences in immune responses. *Nat Rev Immunol*
166 2016;16:626–38. doi:10.1038/nri.2016.90.
- 167 [9] Kaliterna DM, Radić M, Ljutić D. Does estrogen stimulate the pathogenic sort of
168 anticardiolipin antibodies? *Isr Med Assoc J IMAJ* 2014;16:197–8.
- 169 [10] Seriola B, Cutolo M, Garnerio A, Accardo S. Relationships between serum 17 beta-
170 oestradiol and anticardiolipin antibody concentrations in female patients with
171 rheumatoid arthritis. *Rheumatol Oxf Engl* 1999;38:1159–61.
- 172 [11] Verthelyi D, Ansar Ahmed S. Characterization of Estrogen-Induced Autoantibodies to
173 Cardiolipin in Non-Autoimmune Mice. *J Autoimmun* 1997;10:115–25.
174 doi:10.1006/jaut.1996.0121.
- 175 [12] Straub RH. The complex role of estrogens in inflammation. *Endocr Rev* 2007;28:521–74.
176 doi:10.1210/er.2007-0001.
- 177 [13] Oktenli C, Yesilova Z, Kocar IH, Musabak U, Ozata M, Inal A, et al. Study of
178 Autoimmunity in Klinefelter’s Syndrome and Idiopathic Hypogonadotropic
179 Hypogonadism. *J Clin Immunol* 2002;22:137–43. doi:10.1023/A:1015467912592.
- 180 [14] Smith-Bouvier DL, Divekar AA, Sasidhar M, Du S, Tiwari-Woodruff SK, King JK, et al.
181 A role for sex chromosome complement in the female bias in autoimmune disease. *J Exp*
182 *Med* 2008;205:1099–108. doi:10.1084/jem.20070850.
- 183 [15] Jørgensen KT, Rostgaard K, Bache I, Biggar RJ, Nielsen NM, Tommerup N, et al.
184 Autoimmune diseases in women with Turner’s Syndrome. *Arthritis Rheum*
185 2010;62:658–66. doi:10.1002/art.27270.

187 **Figure 1.** Study Flowchart

188 **Figure 2.** (A) aCL antibodies levels in patients with acute Q fever and persistent *C. burnetii*
189 focalized infection. (B) aCL antibodies levels in acute Q fever according to the sex of patients.
190 (C) Percentage of patients with positive aCL antibodies in acute Q fever patients according to
191 the age of patients in men and women. (D) aCL antibodies levels in acute Q fever patients
192 according to the age of patients in men and women. The results are expressed as mean for
193 each age group. Mann-Whitney U test were used * $p < 0.05$, **** $p < 0.0001$

194