



HAL
open science

Experimental team encompasses wide range of competences

R Dumont, D Keeling, M Fitzgerald, S Sharapov, M Baruzzo, P Bonofiglio, C Challis, M Dreval, C Giroud, J Eriksson, et al.

► To cite this version:

R Dumont, D Keeling, M Fitzgerald, S Sharapov, M Baruzzo, et al.. Experimental team encompasses wide range of competences. FEC IAEA 2021 - the 28th IAEA Fusion Energy Conference, May 2021, Nice (E-Conference), France. cea-03244080

HAL Id: cea-03244080

<https://cea.hal.science/cea-03244080>

Submitted on 1 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Scenario preparation for the observation of alpha-driven instabilities and transport of alpha particles in JET DT plasmas

R. Dumont, D. Keeling, M. Fitzgerald, S. Sharapov, M. Baruzzo, P. Bonofiglio, C. Challis, M. Dreval, C. Giroud, J. Eriksson, J. Ferreira, N. Fil, J. Garcia, L. Giacomelli, V. Goloborodko, E. Joffrin, S. Hacquin, N. Hawkes, T. Johnson, Y. Kazakov, V. Kiptily, E. Lerche, M. Lennholm, J. Mailloux, F. Nabais, M. Nocente, L. Piron, M. Poradziński, P. Puglia, E. Solano, R. A. Tinguely, S. Vartanian, H. Weisen, and JET contributors

The logo for JET, consisting of the letters "JET" in a large, bold, blue sans-serif font.



This work has been carried out within the framework of the EUROfusion Consortium and has received funding from the Euratom research and training programme 2014-2018 under grant agreement No 633053. The views and opinions expressed herein do not necessarily reflect those of the European Commission.

Experimental team encompasses wide range of competences



R. Dumont¹, D. Keeling², M. Fitzgerald², S. Sharapov², M. Baruzzo³, P. Bonofiglio⁴, C. Challis², M. Dreval⁵, C. Giroud², J. Eriksson⁶, J. Ferreira⁷, N. Fil², J. Garcia¹, L. Giacomelli⁸, V. Goloborodko⁹, E. Joffrin¹, S. Hacquin¹, N. Hawkes², T. Johnson¹⁰, Y. Kazakov¹¹, V. Kiptily², E. Lerche¹¹, M. Lennholm², J. Mailloux², F. Nabais⁷, M. Nocente⁸, L. Piron³, M. Poradziński¹², P. Puglia¹³, E. Solano¹⁴, R. A. Tinguely¹⁵, S. Vartanian¹, H. Weisen¹³, and JET contributors*

¹CEA, IRFM, F-13108 Saint-Paul-lez-Durance, France

²CCFE, Culham Science Centre, Abingdon, OX14 3DB, UK

³Consorzio RFX, corso Stati Uniti 4, 35127 Padova, Italy

⁴Princeton Plasma Physics Laboratory, Princeton, New Jersey 08543, USA

⁵Institute of Plasma Physics, NSC KIPT, Kharkov, Ukraine

⁶Dept of Physics and Astronomy, Uppsala University, SE-75119 Uppsala, Sweden

⁷Instituto de Plasmas e Fusão Nuclear, IST, Universidade de Lisboa, Portugal

⁸University of Milano-Bicocca, piazza della Scienza 3, 20126 Milano, Italy

⁹Kyiv Institute for Nuclear Research, Prospekt Nauky 47, Kyiv 03680, Ukraine

¹⁰Fusion Plasma Physics, EES, KTH, SE-10044 Stockholm, Sweden

¹¹LPP-ERM/KMS, Ass. EUROFUSION-Belgian State, TEC partner, Brussels, Belgium

¹²Institute of Plasma Physics and Laser Microfusion, Hery 23, 01-497 Warsaw, Poland

¹³École Polytechnique Fédérale de Lausanne (EPFL), Swiss Plasma Center (SPC), CH-1015 Lausanne, Switzerland

¹⁴Laboratorio Nacional de Fusión, CIEMAT, Madrid, Spain


¹⁵MIT PSFC, 175 Albany Street, Cambridge, MA 02139, USA

*See the author list of 'Overview of JET results for optimising ITER operation' by J. Mailloux et al to be published in Nuclear Fusion Special issue: Overview and Summary Papers from the 28th Fusion Energy Conference (Nice, France, 10-15 May 2021)

JET is the only ongoing fusion device in which alpha effects can be studied experimentally



- TAEs (Toroidal Alfvén Eigenmodes) are instabilities excited by energetic ion radial pressure gradient, experience moderate damping by thermal plasma species
- α -driven TAEs may induce energetic ion transport / losses in burning plasmas (e.g. in ITER)
- TAEs observed in all plasmas with elevated q-profile ($q_0 > 1.5$) in JET DT plasmas (DTE1)
- Caveat: ICRH-accelerated ions always present, provided large TAE drive, dominating α -drive
- No conclusions regarding α -driven TAEs could be drawn from JET DTE1 data [Sharapov NF 1999], unlike from successful TFTR experiments [Budny NF 1992, Spong NF 1995, Nazikian PRL 1997]



Dedicated advanced scenario under development for the study of alpha-driven instabilities in JET




- α -driven TAE growth / damping rate is given by


$$\gamma \equiv \gamma_\alpha - \gamma_d = -Cq^2\beta_\alpha \left(1 - \frac{\omega_\alpha^*}{\omega}\right) F(v_\alpha/v_A) - \gamma_d$$

- Key parameters to maximise TAE drive (γ_α):
 - High α -pressure + large pressure gradient
 - Large P_{add} , T_i (increase fusion yield), T_e (increase slowing-down time)
 - Low n_e , low I_p (still large enough for alpha confinement)
 - Elevated q-profile

- Optimal conditions for TAE excitation not naturally fulfilled by parameters of baseline or hybrid steady-state scenarios in JET DT plasmas
 [Garcia, EX1-989] [Mailloux, OV1-1080]
 → dedicated development of an advanced scenario for JET-ILW




Observation of alpha-driven TAEs in JET requires afterglow phase in the pulse



JET DT pulse 41723 – CASTOR-K, MISHKA
 [Sharapov NF 1999]

- TAEs very stable during main heating phase
 - Beam + th. damping ~1-2%
 - Alpha drive ~0.1-0.2%
- α -driven TAEs may be observed only during afterglow phase





- No ICRH before afterglow period in DT plasmas
- ICRH used in D plasmas to develop scenario & probe TAE stability

ICRH used to reveal and tune ITBs at relatively moderate levels of NBI power




ITB triggered




- Pulse 94850: ITB triggered with NBI+ICRH - $R_{NT}=2.6 \times 10^{16}/s$ with $P_{tot}=(20+3.8)MW$
- No sign of shear reversal in most pulses. ITB triggered upon $q=2$ surface entering plasma
- Scenario development and ITB tuning done with $\sim 20MW$ NBI power + $\sim 4MW$ ICRH, then applied to NBI-only pulses at higher NBI power levels ($> 24MW$)

Afterglow successfully triggered by bespoke real-time control scheme





- Pulse 95973 set up to trigger afterglow when
 - Neutron rate large enough ($> 8 \times 10^{15}/s$)
 - dR_{NT}/dt low ($< 1 \times 10^{13}/s^2$)
- First successful demonstration, although just above threshold (limited NBI power)
- RF power applied to ensure safe pulse termination (requirement for DTE2)
- Neutron rate threshold can be used as dud detector (requirement for DTE2)

ELM control achieved in advanced scenario



- Pulse 95973: ELM-free/type-I ELMs period before peak performance
- In this example, radiation peaking induces power ramp-down just following afterglow
- ITB beneficial for performance → ELM control needed
- ELM pacing by D pellets (~1.4mm, ~30Hz-45Hz) effective
- Last sessions much less affected by ELM issues compared to previous sessions, despite larger NBI power
- H pellets successfully tested → usable during T campaign (although H minority concentration increase can make ICRH use difficult)


Core TAEs observed during RF-powered ITBs



- Core TAEs $n=3-7$ seen during D(H) ICRH at 51MHz, damped by thermal/NBI ions as P_{NBI} increases
- Broadband modes 60-120kHz often seen in ITB pulses after TAEs disappear under study [Fil 2021]
- Alfvén cascades observed only during pulse termination (shear reversal)



Record pulse, performance comparable to reference

C-wall pulse for alpha studies



- Pulse 96852: highest neutron rate obtained to date in JET-ILW with NBI only ($R_{NT}=2.55 \times 10^{16}/s$)
- Good ITB obtained with NBI starting at 45.0s (differs from previous sessions)
- Performance similar to reference JET C-wall pulse for alpha studies
- Afterglow automatically triggered on R_{NT} rollover
- No ELM-free/type I ELM period seen (pacing pellets used)

Extrapolation to DT confirms significant progress in scenario development



- Pulse 92054: former best NBI-only pulse for the study of alpha-driven instabilities [Dumont NF 2018]
- TRANSP DT simulation of 96852 very similar to CRONOS DT simulation of 92054 with extrapolated NBI power (31MW) [Garcia NF 2019]

Summary & prospects



- Main deliverable of development phase in D achieved: NBI-only pulses with good fusion performance in the presence of Internal Transport Barriers
- Core-localised TAEs in pulses with ICRH
- Pulses ready to be run in T and DT, include
 - ELM pacing by pellets
 - Real-time control-triggered afterglow
 - Gas injection with T-compatible modules
- Impossible to achieve 100% reproducibility in this scenario - included in strategy for DTE2 by allowing small changes in NBI switch-on time
- Data analysis ongoing, including fast ion transport/losses, MHD...
- Modelling effort ongoing
 - TRANSP modelling of best pulses, using refined equilibrium - extrapolations to DT
 - MHD stability analyses to evaluate damping mechanisms remaining during afterglow phase, and compare to alpha drive
- Experimental effort to couple TAE antenna power in these plasmas → probe stable modes [Tinguely, EX/P3-889]
- JET/TFTR comparisons included in DTE2 experimental plans