

Impact of leakage under divertor baffle on detachment onset in WEST with SOLEDGE3X modeling

H Yang, J Bucalossi, G Ciraolo, H Bufferand, N Fedorczak, P Tamain, G

Falchetto, N Rivals, J Gunn, Y Marandet, et al.

▶ To cite this version:

H Yang, J Bucalossi, G Ciraolo, H Bufferand, N Fedorczak, et al.. Impact of leakage under divertor baffle on detachment onset in WEST with SOLEDGE3X modeling. 47th EPS Conference on Plasma Physics, EPS, Jun 2021, Virtual conference, France. pp.P2.1025. cea-03300557

HAL Id: cea-03300557 https://cea.hal.science/cea-03300557

Submitted on 3 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Impact of leakage under divertor baffle on detachment onset in WEST with SOLEDGE3X modeling

H. Yang¹, J. Bucalossi¹, G. Ciraolo¹, H. Bufferand¹, N. Fedorczak¹, P. Tamain¹, G. Falchetto¹,

N. Rivals¹, J. Gunn¹, Y. Marandet², B. Pégourié¹, S. Vartanian¹, the WEST team^{*} ¹ CEA, IRFM, F-13108 Saint Paul-lez-Durance, France ² Aix Marseille University, CNRS, PIIM, Marseille, France ^{*} See http://west.cea.fr/WESTteam

The WEST (W-tungsten Environment in Steady-state Tokamak) is the transformation of the Tore Supra tokamak from a carbon limiter to a tungsten divertor configuration. In recent experiment campaign C5, the divertor pumping capability has been improved by the sealing of the space between the divertor baffle and the vacuum vessel. This modification leads to an interesting question about how the leakage below the outer baffle will influence the plasma in WEST. To investigate this question, we made simulations of three leak cases and one no leak case with the help of SOLEDGE3X-EIRENE transport code. Through the analysis of the simulation results, we work out the impact of leakage under the outer baffle and explain the physics behind some phenomena observed.

We investigated the density regimes analyzing how parameters like density, temperature, particle flux, and neutral pressure evolve as functions of gas puff rate and electron density at the outer midplane separatrix comparing among the cases with and without leak. The results show that the case without leak has better performance than the cases with leak in trapping the neutral particles and has higher neutral pressure near the baffle by more than 35%, which can lead to greater power dissipation in the divertor, thus lower the detachment threshold in upstream separatrix density by more than 10%. At the same time, the operational gas puffing range becomes broader by a factor from 1 to 5 in the case without leak with respect to the three cases with leak. Simulation results are confronted with experimental data from the configurations with and without leak under the baffle in order to get further insight on the neutral particle circulation in WEST.

Keywords: detachment, SOL, divertor baffle closure, leakage effect