



HAL
open science

The free energy of transfer in metal-extracting water-poor microemulsions as the crucial molecular driving force

Asmae El Maangar, Ange Maurice, Varun Rai, Johannes Theisen, Christophe Penisson, Fabien Olivier, Jean Duhamet, Thomas Zemb, Jean-Christophe Gabriel

► To cite this version:

Asmae El Maangar, Ange Maurice, Varun Rai, Johannes Theisen, Christophe Penisson, et al.. The free energy of transfer in metal-extracting water-poor microemulsions as the crucial molecular driving force. ECIS 2 2021 - 35th Conference of the European Colloid and Interface Society, ECIS, Sep 2021, Athens, Greece. cea-03338687v1

HAL Id: cea-03338687

<https://cea.hal.science/cea-03338687v1>

Submitted on 9 Sep 2021 (v1), last revised 7 Oct 2021 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The free energy of transfer in metal-extracting water-poor microemulsions as the crucial molecular driving force

Asmae El Maangar¹, Christophe Penisson¹, Ange Maurice², Jean-Christophe Gabriel^{2,3} and Thomas Zemb¹

¹ICSM. Univ. Montpellier. CEA. CNRS. ENSCM. Marcoule. France.

²SCARCE Laboratory, Energy Research Institute @ NTU (ERI@N), Nanyang Technology University, 637553, Singapore.

³Université Paris-Saclay, CEA, CNRS, NIMBE, LICSEN, 91191, Gif-sur-Yvette, France.

Presenting author e-mail: asmae.elmaangar@cea.fr

It is a matter of strategic independence for Europe to urgently find processes taking account of environmental and economic issues, when mining and recycling rare earth elements. Separation and recycling of rare earths from electronic waste is important for the success of present and future carbon-free technologies. Solvent extraction based on “water-poor” microemulsions is one of the first technologies allowing the take-off of circular economy. However, an optimized process on an industrial scale has not been established. One major reason is the lack of fundamental knowledge, therefore designing a cost-efficient, adaptive and predictive formulation is still out of scope of possibilities. Our objective is to develop the fundamental understanding involved in the process’ complex fluids (experimental and theoretical) concerning liquid-liquid extraction of REE and in order to render environmentally friendly processes economically competitive.

In order to gain more comprehensive view between quaternary phase diagrams (macroscopic view), extraction properties and the molecular driving forces in solvent extraction (colloidal view at nanoscale), downscaling the extraction process to microfluidics proves to be a useful approach allowing accurate and rapid adaptation to such composition changes [1]. we describe here the first automated microfluidic tool integrated with X-ray fluorescence microanalysis allowing to measure with enough precision the free energy of transfer of ions along the lines in quaternary phase diagrams. Our contribution towards a more complete understanding in this matter is the analysis and comparison of the phase behavior, extracting efficiency and selectivity of such systems as well as the correlation of these findings with the colloidal “jenaics” approach by identifying the molecular driving forces favoring or quenching the transfer.

Keywords: Complex fluids, Liquid-liquid extraction, Microfluidics, X-ray Fluorescence, on-line measurement, Rare earths, Thermodynamics, Kinetics.




Figure 1. Graphical free energy balance between supramolecular complexation, the chain reconfiguration term and the internal energy of the polar cores.