



Key role of phase dynamics & diamagnetic drive on Reynolds stress in fusion plasma turbulence

Y. Sarazin, Guilhem Dif-Pradalier, Ph. Ghendrih, X. Garbet, A. Berger, R. Bigué, Emilie Bourne, Virginie Grandgirard, Kevin Obrejan, R. Varennes, et al.

► To cite this version:

Y. Sarazin, Guilhem Dif-Pradalier, Ph. Ghendrih, X. Garbet, A. Berger, et al.. Key role of phase dynamics & diamagnetic drive on Reynolds stress in fusion plasma turbulence. AAPPS-DPP 2021 - 5th Asia Pacific Conference on Plasma Physics, Sep 2021, E-Conference, Japan. cea-03520862

HAL Id: cea-03520862

<https://cea.hal.science/cea-03520862>

Submitted on 11 Jan 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



European
Commission



Key role of phase dynamics & diamagnetic drive on Reynolds stress in fusion plasma turbulence

Y. Sarazin, G. Dif-Pradalier, X. Garbet, Ph. Ghendrih, A. Berger,
R. Bigué, E. Bourne, V. Grandgirard, K. Obrejan, R. Varennes
CEA, IRFM, 13108 Saint-Paul-lez-Durance, France



L. Vermare

LPP, CNRS, Ecole Polytechnique, Palaiseau, France

Ack: N. Fedorczak, Ö. Gürcan, P. Hennequin, C. Passeron
Festival de Théorie 2019, Chengdu Theory Festival 2018

AAPPS-DPP, 2021/09/29

- ▶ Turbulence rotates poloidally:
mean flow velocity $V_E \sim -E_r$
- ▶ Radial shear of V_E (i.e. E_r') \Rightarrow turbulent vortex decorrelation
 - Dominant mechanism for turbulence control
 - Transport barriers, optimized confinement



[Biglari-Diamond-Terry 1990;
Itoh-Itoh 1996; Terry 2000]

- ▶ Turbulence generates zonal & mean flows via Reynolds stress
- ▶ Issue addressed in this talk:
What dominant mechanisms govern Reynolds stress dynamics ?

- Start from GK equation, focusing on E×B advection:

$$\partial_t \bar{f} + \underbrace{\{J.\phi, \bar{f}\}}_{\text{Poisson brackets (account for } E \times B \text{ advection: } \mathbf{v}_E \cdot \nabla \bar{f} \text{)}} = RHS \quad \text{with gyro-average operator } J \approx 1 + \frac{\mu}{2} \nabla_{\perp}^2$$

Poisson brackets (account for E×B advection: $\mathbf{v}_E \cdot \nabla \bar{f}$)

- Account for quasi-neutrality (adiabatic electrons):

$$n_0 \Omega = \langle J. \bar{f} \rangle_v \approx \langle \bar{f} \rangle_v + \frac{1}{2} \nabla_{\perp}^2 p_{\perp} \quad \text{Potential vorticity} \quad \Omega = \phi - \langle \phi \rangle - \nabla_{\perp}^2 \phi$$

- Where it comes: transport of potential vorticity

$$\partial_t \Omega - \nabla_{\perp,i} \{ \phi + p_{\perp}, \nabla_{\perp,i} \phi \} = \langle J.RHS \rangle_v$$

[Parra-Catto PPCF 2009;
Abiteboul PoP 2011;
Ajay JPP 2020]

- Integrate & flux-surface average → equation on $\langle u_{E\theta} \rangle = \partial_r \langle \phi \rangle$

$$\frac{\partial \langle u_{E\theta} \rangle}{\partial t} + \frac{1}{r^2} \frac{\partial}{\partial r} [r^2 (\pi + \pi^*)] = rhs$$

[Diamond-Kim PoP 1991;
Smolyakov PoP 2000;
McDevitt PRL 2010]

Reynolds' stress

$$\pi = \langle \tilde{u}_{Er} \tilde{u}_{E\theta} \rangle$$

Diamagnetic Reynolds' stress

$$\pi^* = \langle \tilde{u}_r^* \tilde{u}_{E\theta} \rangle$$

with $\begin{cases} \tilde{u}_{Er} = -\frac{1}{r} \partial_{\theta} \tilde{\phi} \\ \tilde{u}_r^* = -\frac{1}{r} \partial_{\theta} \tilde{p}_{\perp} \end{cases}$

[Sarazin, Dif-Pradalier, Garbet et al., PPCF 63 (2021) 064007]


- ▶ Comparative weight & properties of
Reynolds stress $\pi = \langle \tilde{u}_{Er} \tilde{u}_{E\theta} \rangle$ vs.
diamagnetic Reynolds stress $\pi^* = \langle \tilde{u}_r^* \tilde{u}_{E\theta} \rangle$

- ▶ Dynamics of the **Reynolds stress**: specific role of the **phase**


Means: 1st principle gyrokinetic simulations (GYSELA)
reduced nonlinear model

► Flux-driven 5D GyroKinetics

- Kinetic description → gyro-centers distribution function (Boltzmann eq.)
- Self-consistency = Quasi-Neutrality (Poisson eq.)
- Adiabatic electrons in cases under study




- ▶ Analysis of a statistically saturated turbulent regime
- ▶ $\mathbf{E} \times \mathbf{B}$ turbulent flux exhibits bursts and avalanches
- ▶ Imprint of **avalanche-like patterns** on $\mathbf{E} \times \mathbf{B}$ mean flow
- ▶ Similar pattern on the total Reynolds' stress $\pi + \pi^*$




- Diamagnetic Reynolds stress about 2 times larger than E×B contribution

- Diamagnetic $\pi^* = \langle \tilde{u}_r^* \tilde{u}_{E\theta} \rangle$ and E×B $\pi = \langle \tilde{u}_{Er} \tilde{u}_{E\theta} \rangle$ contributions are **in phase** (minima & maxima coincide)



- Consequence:
Pressure is NOT simply advected by E×B flow ≠ passive scalar
⇒ possible impact on analysis of exp. data Cf. e.g. [Vermare PoP 2018]




► Let's assume $\partial_t p_{\perp} + \mathbf{u}_E \cdot \nabla p_{\perp} = 0$.

$$\text{Then } \hat{p}_{\perp k, \omega} = -\frac{\omega_p^*}{\omega} \hat{\phi}_{k, \omega}$$

($\omega_p^* = k_\theta \partial_r \langle p_{\perp} \rangle$ ion diamagn. freq.)

$$\text{So that } \pi^* = - \sum_{\omega, k} k_r k_\theta \frac{\omega_p^*}{\omega} |\hat{\phi}_{k, \omega}|^2$$

$$\text{Opposite to } \pi = \sum_{\omega, k} k_r k_\theta |\hat{\phi}_{k, \omega}|^2$$



► GYSELA simulations (ITG turbulence):

- $\omega \approx \omega_p^*$

► Let's assume $\partial_t p_{\perp} + \mathbf{u}_E \cdot \nabla p_{\perp} = 0$.

$$\text{Then } \hat{p}_{\perp k, \omega} = -\frac{\omega_p^*}{\omega} \hat{\phi}_{k, \omega}$$

($\omega_p^* = k_\theta \partial_r \langle p_{\perp} \rangle$ ion diamagn. freq.)

$$\text{So that } \pi^* = - \sum_{\omega, k} k_r k_\theta \frac{\omega_p^*}{\omega} |\hat{\phi}_{k, \omega}|^2$$


$$\text{Opposite to } \pi = \sum_{\omega, k} k_r k_\theta |\hat{\phi}_{k, \omega}|^2$$

► GYSELA simulations (ITG turbulence):

- $\omega \approx \omega_p^*$
- No clear relationship between $\hat{p}_{\perp k, \omega}$ and $\hat{\phi}_{k, \omega}$

► Linear GK relationship: resonances

$$\hat{p}_{\perp k, \omega} = - \left\langle \frac{\omega^* - \omega_d - k_{\parallel} v_{\parallel}}{\omega - \omega_d - k_{\parallel} v_{\parallel}} \mu B F_M \right\rangle_v$$



[Sarazin, Dif-Pradalier, Garbet et al., PPCF 63 (2021) 064007]

- ▶ Comparative weight & properties of
Reynolds stress $\pi = \langle \tilde{u}_{Er} \tilde{u}_{E\theta} \rangle$ vs.
diamagnetic Reynolds stress $\pi^* = \langle \tilde{u}_r^* \tilde{u}_{E\theta} \rangle$
- ▶ Dynamics of the **Reynolds stress**: specific role of the **phase**
Means: 1st principle gyrokinetic simulations (GYSELA)
reduced nonlinear model


► Magnetized plasma $\mathbf{B} \neq \text{Cst}$

Isotherm, cold ions, only resonant modes $k_{\parallel}=0$

$$\begin{cases} \partial_t n + \mathbf{v}_E \cdot \nabla n = S & \text{Density} \\ \partial_t \Omega + \mathbf{v}_E \cdot \nabla \Omega + g \partial_y n = 0 & \text{Vorticity} \end{cases}$$

$\mathbf{v}_E = \mathbf{z} \times \nabla \phi$

$\sim 2(\rho_s/R)$ effective gravity
(only retained compressibility term
→ governs interchange instability)



► Scale separation: mean $\langle \dots \rangle(x,t)$

& fluctuations $\tilde{\dots}(x,y,t)$

Mean poloidal flow $V_E = \partial_x \langle \phi \rangle$

$$n(x, y, t) = N(x, t) + \tilde{n}(x, y, t)$$

$$\Omega(x, y, t) = V'_E(x, t) + \nabla_{\perp}^2 \tilde{\phi}(x, y, t)$$

► Retaining single wave vector k_y

$$\tilde{n}(x, y, t) = n_k e^{ik_y y} + cc$$

$$\tilde{\phi}(x, y, t) = \phi_k e^{ik_y y} + cc$$

[Sarazin PoP 2000;
Bian PoP 2003;
McMillan PoP 2009]

► 1D nonlinear system (+ Dissipation)

[Sarazin PoP 2000;
 Bian PoP 2003;
 McMillan PoP 2009]

- Mean fields ($\in \mathbb{R}$)

$$\begin{cases} \partial_t N &= -\partial_x \Gamma + S \\ \partial_t V_E &= -\partial_x \Pi \end{cases}$$

Turbulent Flux $\Gamma = \langle \tilde{v}_{Ex} \tilde{n} \rangle = -\Im(2k_y n_k \phi_k^*)$

Reynolds stress $\Pi = \langle \tilde{v}_{Ex} \tilde{v}_{Ey} \rangle = -\Im(2k_y \phi_k^* \partial_x \phi_k)$

- Fluctuations ($\in \mathbb{C}$)

$$\begin{cases} \partial_t n_k &= -ik_y (V_E n_k + v_* \phi_k) \\ \partial_t \Omega_k &= -ik_y (V_E \Omega_k + g n_k - V_E'' \phi_k) \end{cases}$$

Potential vorticity

$$\Omega_k = (\partial_x^2 - k_y^2) \phi_k$$

Diamagnetic drift

$$v_* = -\partial_x N$$

► Linear instability = "interchange" ~ Rayleigh-Bénard

$$\partial_t n_k = -ik_y (V_E n_k + v_\star \phi_k)$$

Diamagnetic drift $v_\star = -\partial_x N$

$$\partial_t \Omega_k = -ik_y (V_E \Omega_k + g n_k - V_E'' \phi_k)$$

Potential vorticity $\Omega_k = (\partial_x^2 - k_y^2) \phi_k$

► Dispersion relation

$$\omega_\pm - k_y V_E = \frac{k_y V_E''}{2k_\perp^2} \pm \frac{k_y}{k_\perp} \left(\frac{V_E''^2}{4k_\perp^2} - gv_\star \right)^{1/2}$$

Doppler shift

Stabilization
⇒ Instability Threshold


Instability drive = curvature
× mean gradient
→ Linear growth rate γ

$$V_E = Cst \Rightarrow \gamma = \frac{k_y}{k_\perp} \sqrt{gv_\star}$$

- Discriminate **phase** and **amplitude** of fluctuations

$$\begin{cases} n_k = |n_k| e^{i\varphi_n} \\ \phi_k = |\phi_k| e^{i\varphi_\phi} \end{cases} \Rightarrow \begin{cases} \Gamma = -2k_y |n_k| |\phi_k| \sin \Delta\varphi_{n\phi} \\ \Pi = -2k_y |\phi_k|^2 \partial_x \varphi_\phi \end{cases}$$

Phase gradient $\partial_x \varphi_\phi$
 \equiv Phase shift of k_y waves






- Reynolds' stress Π can grow due to

Amplitude $|\phi_k|^2$
 Phase gradient $\partial_x \varphi_\phi \sim k_x$

Note: phase curvature already identified as a possible key player
"phase patterning can generate ZF w/o turbulence inhomogeneity": $\partial_x \Pi \sim |\phi_k|^2 \partial_x^2 \varphi_\phi$

[Z. Guo, P. Diamond PRL 2016;
 Xu NF 2020]

- **Zonal Flows** are excited above a threshold in source magnitude S_0
- Turbulence magnitude & effective transport are reduced at their onset




- **Zonal Flows** are excited above a threshold in source magnitude S_0
- Turbulence magnitude & effective transport are reduced at their onset
- ZF are driven by an instability of the phase gradient (reminiscent of the tilting instability)

[Drake PFB 1992; Guzdar PFB 1993;
Rosenbluth PoP 1994; Fedorczak PPCF 2013]

- Estimate of phase instability growth rate

$$\begin{aligned} \frac{\partial(\partial_x \varphi_\phi)}{\partial t} &= -k_y \frac{\partial V_E}{\partial x} \quad \leftrightarrow \quad \frac{dk_r}{dt} = -k_\theta V'_E \\ \frac{\partial V_E}{\partial t} &= 2k_y \frac{\partial}{\partial x} (|\phi_k|^2 \partial_x \varphi_\phi) \\ \Rightarrow \frac{\partial^2(\partial_x \varphi_\phi)}{\partial t^2} &\approx \boxed{-2k_y^2 \frac{\partial^2 |\phi_k|^2}{\partial x^2}} \partial_x \varphi_\phi \\ &\sim (\text{phase instability growth rate})^2 \end{aligned}$$




- Zonal Flows are excited above a threshold in source magnitude S_0
- Turbulence magnitude & effective transport are reduced at their onset
- ZF are driven by an instability of the phase gradient (reminiscent of the tilt instability)

[Drake PFB 1992
Rosenbluth PoP 1994]

- Estimate of phase instability growth rate

$$\begin{aligned} \frac{\partial(\partial_x \varphi_\phi)}{\partial t} &= - \frac{\partial V_E}{\partial x}, \quad \text{at } \partial_x \varphi_\phi = 0 \\ \frac{\partial V_E}{\partial t} &= 2k_y \frac{\partial \varphi_\phi}{\partial x} \Rightarrow \frac{\partial^2(\partial_x \varphi_\phi)}{\partial t^2} = -k_\theta V'_E \\ \Rightarrow \frac{\partial^2(\partial_x \varphi_\phi)}{\partial t^2} &\approx \boxed{-2k_y^2 \frac{\partial^2 |\phi_k|^2}{\partial x^2}} \partial_x \varphi_\phi \\ &\sim (\text{phase instability growth rate})^2 \end{aligned}$$

What about in tokamak plasma turbulence?



► Similar decomposition with **3D output GYSELA data** (ion turbulence)

$$\Pi = - \sum_{m>0,n} \frac{2m}{r} |\hat{\phi}_{mn}|^2 \partial_r \varphi_{mn}^\phi = \sum_{m>0,n} \Pi_{mn} \quad \begin{array}{l} \text{2D Fourier decomposition} \\ (m,n) = (\text{poloidal, toroidal}) \\ \text{mode numbers} \end{array}$$


$$= \mathcal{I}(r,t) \varphi'(r,t) \Delta(r,t)$$

Turbulence intensity

$$\sum_{m>0,n} \frac{2m}{r} |\hat{\phi}_{mn}|^2$$

$$\text{Phase gradient: } \varphi'(r,t) = - \sum_{m>0,n} W_{mn} \partial_r \varphi_{mn}^\phi$$

Mode weight $\log |W_{mn}|$ ($r/a \approx 0.72$)



$$W_{mn}(r) = \frac{\langle |\Pi_{mn}| \rangle_t}{\langle \Pi \rangle_t}$$

- Similar decomposition with **3D output GYSELA data** (ion turbulence)


$$\Pi = - \sum_{m>0,n} \frac{2m}{r} |\hat{\phi}_{mn}|^2 \partial_r \varphi_{mn}^\phi = \sum_{m>0,n} \Pi_{mn} \quad \begin{array}{l} \text{2D Fourier decomposition} \\ (m,n) = (\text{poloidal, toroidal}) \\ \text{mode numbers} \end{array}$$

$$= \mathcal{I}(r, t) \varphi'(r, t) \Delta(r, t)$$

Turbulence intensity

$$\text{Phase gradient: } \varphi'(r, t) = - \sum_{m>0,n} W_{mn} \partial_r \varphi_{mn}^\phi$$

- Reynolds stress Π well correlated with φ' , not with \mathcal{I}
 \Rightarrow phase dynamics is critical



► Zonal & mean flows excited by turbulence

- Reynolds stress $\pi = \langle \tilde{u}_{Er} \tilde{u}_{E\theta} \rangle \approx -\frac{1}{B^2 r} \left\langle \partial_\theta \tilde{\phi} \partial_r \tilde{\phi} \right\rangle$
- Diamagnetic Reynolds stress $\pi^\star = \langle \tilde{u}_r^\star \tilde{u}_{E\theta} \rangle \approx -\frac{1}{enB^2 r} \left\langle \partial_\theta \tilde{p}_\perp \partial_r \tilde{\phi} \right\rangle$

► Diamagnetic Reynolds stress in phase with & larger than E×B RS

- Pressure does not behave as a passive scalar
- Numerical evidence of its critical role in the build up of large scale flows at the plasma boundary → cf. talk Dif-Pradalier (this conf.)
- Experimental evidence? Role in triggering transport barriers?

► Reynolds stress dynamics governed by the phase gradient

- Growing evidence that phase is a key ingredient to RS drive
- Robustness w.r.t. turbulence properties (e.g. electron turbulence)?
- Role of phase instab. in triggering staircases / transport barriers?
- What mechanism governs the transport of phase, if any ?

Back up slides

- ▶ Drift velocities $\mathbf{u}_\perp^{(1)} = \mathbf{u}_E + \mathbf{u}_i^* = \frac{\mathbf{B} \times \nabla \phi}{B^2} + \frac{\mathbf{B} \times \nabla p_\perp}{enB^2}$
- $\mathbf{u}_\perp^{(2)} = \mathbf{u}_{pol} \approx -\frac{m_i}{eB^2} [\partial_t + (\mathbf{u}_E + \mathbf{u}_i^*) \cdot \nabla] \nabla_\perp \phi$

- ▶ Charge conservation $\nabla \cdot \mathbf{j} = 0$

$$\begin{aligned}\nabla \cdot \mathbf{j}_{pol} &\approx -\frac{nm_i}{B^2} (\partial_t + \mathbf{u}_E \cdot \nabla) \nabla_\perp^2 \phi - \frac{m_i}{eB^3} \nabla \cdot \left[\left(\frac{\mathbf{B} \times \nabla p_\perp}{B} \cdot \nabla \right) \nabla_\perp \phi \right] \\ &\approx -\frac{nm_i}{B^2} \left(\partial_t \nabla_\perp^2 \phi + \frac{1}{B} \{ \phi, \nabla_\perp^2 \phi \} \right) \\ &\quad - \frac{m_i}{eB^3} \{ \nabla_{\perp,i} p_\perp, \nabla_{\perp,i} \phi \} + \{ p_\perp, \nabla_\perp^2 \phi \}\end{aligned}$$

~ divergence of $\pi^* = \langle \tilde{u}_r^* \tilde{u}_{E\theta} \rangle$

~ divergence of $\pi = \langle \tilde{u}_{Er} \tilde{u}_{E\theta} \rangle$