



Alkyl formates as transfer hydroalkylation reagents -Hydrobenzylation of imines

Etienne Crochet, Lucile Anthore-Dalion, Thibault Cantat

► To cite this version:

Etienne Crochet, Lucile Anthore-Dalion, Thibault Cantat. Alkyl formates as transfer hydroalkylation reagents -Hydrobenzylation of imines. GreenCat 2022 - 4th International Green Catalysis Conference, Apr 2022, Rennes, France. . cea-03655072

HAL Id: cea-03655072

<https://cea.hal.science/cea-03655072>

Submitted on 29 Apr 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Alkyl formates as transfer hydroalkylation reagents – Hydrobenzylation of imines


Etienne Crochet, Lucile Anthore-Dalion, and Thibault Cantat

NIMBE, CEA, CNRS, Université Paris-Saclay, CEA Saclay, 91191 Gif-sur-Yvette, France





etienne.crochet@cea.fr

Hydrofunctionalization of unsaturated bonds, such as carbonyl, imines, alkenes or alkynes, provide a rapid and atom-economic access to functionalized molecules.¹ Among those processes, hydroalkylation, which formally adds a R-H alkane across a C=C, C=O or C=N bond, has been intensively studied. However, in these reactions a sacrificial hydride or hydrogen donor is required, usually a hydrosilane or a hydroborane.² Therefore, hydroalkylation by transfer was explored to increase the atom economy. In 2011, Oshima's group proposed a transfer from allenyl or homoallyl alcohols to aldehydes or imines under Cu-catalysis.³ Then, in 2013, Tang's group designed efficient and versatile hydroalkyl-donors based on Hantzsch esters or benzothiazoles.⁴ Yet, these two reactions did not improved the atom economy, although recyclable but under harsh conditions. Following previous work on transfer hydrosilylation,⁵ formate appeared as a new potential transfer reagent. We assumed that alkyl formates can perform transfer hydroalkylation reactions through transition-metal catalysis. This process would present a high atom economy and CO₂, as the only byproduct, can be recycle in formic acid, which can also be biosourced. As a proof of concept, we developed a novel catalytic reaction of transfer hydroalkylation from alkyl formates using benzyl formates and imines as model reagents.


Hydroalkylation



Transfer reagents

Oshima *et al.*, 2011Tang *et al.*, 2013

An example



- (1) *Hydrofunctionalization*; Ananikov, V. P., Tanaka, M., Eds.; Topics in Organometallic Chemistry; Springer Berlin Heidelberg: Berlin, Heidelberg, **2013**; Vol. 43.
- (2) Wang, X.-X.; Lu, X.; Li, Y.; Wang, J.-W.; Fu, Y. *Sci. China Chem.* **2020**, 63 (11), 1586–1600.
- (3) Sai, M.; Yorimitsu, H.; Oshima, K. *Angew. Chem. Int. Ed.* **2011**, 50 (14), 3294–3298.
- (4) Li, G.; Chen, R.; Wu, L.; Fu, Q.; Zhang, X.; Tang, Z. *Angew. Chem. Int. Ed.* **2013**, 52 (32), 8432–8436.
- (5) Chauvier, C.; Godou, T.; Cantat, T. *Chem. Commun.* **2017**, 53 (85), 11697–11700.