



HAL
open science

Synthesis and optical properties of rod-shaped graphene nanoparticles

Daniel Medina-Lopez, Thomas Liu, Christine Elias, Loïc Rondin,
Jean-sébastien Lauret, Stéphane Campidelli

► **To cite this version:**

Daniel Medina-Lopez, Thomas Liu, Christine Elias, Loïc Rondin, Jean-sébastien Lauret, et al.. Synthesis and optical properties of rod-shaped graphene nanoparticles. 241st ECS Meeting, May 2022, Vancouver, Canada. cea-03687145

HAL Id: cea-03687145

<https://cea.hal.science/cea-03687145>

Submitted on 3 Jun 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



VANCOUVER
BC • CANADA



241st ECS Meeting

May 29- June 2, 2022

Vancouver Convention Center

B05-0829 - (Invited) Synthesis and Optical Properties of Rod-Shaped Graphene Nanoparticles



Tuesday, 31 May 2022



20:20 - 20:40



Vancouver Convention Center - West Meeting Room 202

Abstract

The scientific community has given much interest to graphene in recent years because of its good mechanical, thermal and electrical properties. Nonetheless, applications in optics and optoelectronics can be limited due to the absence of a bandgap. In order to engineer a bandgap, we can reduce the size of graphene into graphene quantum dots (GQDs). GQDs synthesis is primarily described via top-down methods (lithography, hydrothermal and electrochemical approaches).[1] Though effective, these methods do not allow accurate control of the size, shape and edges of the GQDs. On the other hand, bottom-up graphene materials have developed exponentially for the last decade with the synthesis of highly controlled graphene nanoribbon structures and moderately soluble graphene quantum dots.[2,3] Our group demonstrated that bottom-up GQDs could act as single-photon emitters exhibiting high brightness and stability.[4]

In order to investigate the structure-property relationship, we designed a series of rod-shaped graphene nanoparticles that differ only by their length, keeping the same morphology, symmetry and edge states (see figure). These nanoparticles are fully soluble, which facilitate their purification and individualization in solution and give rise to well-defined absorption spectra. Here we report on the synthesis of these GQDs, and we will present results on advanced characterization of their optical properties.[5]

[1] Yang, J.-S., Martinez, D. A. & Chiang, W.-H. Synthesis, Characterization and Applications of Graphene Quantum Dots in *Recent Trends in Nanomaterials: Synthesis and Properties*, Khan, Z. H., Springer Singapore, (2017).

[2] Narita, A., Wang, X.-Y., Feng, X. & Müllen, K. New advances in nanographene chemistry. *Chem. Soc. Rev.* **44**, 6616–6643 (2015).

[3] Wang, X.-Y., Yao, X. & Müllen, K. Polycyclic aromatic hydrocarbons in the graphene era. *Sci. China Chem.* **62**, 1099–1144 (2019)

[4] Zhao, S. *et al.* Single photon emission from graphene quantum dots at room temperature. *Nat. Commun.* **9**, 3470 (2018).





Figure 1: Structures and absorption spectra of rod-shaped GQDs: LC₇₈-6t-Bu (yellow), LC₉₈-8t-Bu (red) and LC₁₁₄-10t-Bu (green).

Corresponding Author



Daniel Medina Lopez
CEA-Saclay

Authors



Thomas Liu
LUMIN ENS PARIS SACLAY



Christine Elias
LUMIN ENS PARIS SACLAY



Loïc Rondin
LUMIN ENS PARIS SACLAY



Js Lauret
ENS Paris Saclay



Stephane Campidelli
CEA Saclay

View Related

B05 - Nanocarbons

F. D'Souza (University of North Texas) and T. Torres (Autonoma University of Madrid)



Tuesday, 31 May 2022



19:00 - 21:20



Vancouver Convention Center - West Meeting Room 202

B05 - Fullerenes - Endohedral Fullerenes and Molecular Carbon

Carbon Nanostructures and Devices

Similar

(Invited) Influence of Vibrations on the Emission Properties of Single Graphene Quantum Dots

T. Liu (LUMIN ENS PARIS SACLAY), C. Tonnelé (MONS Univeristy), C. Elias, L. Rondin, B. Carles (LUMIN ENS PARIS SACLAY), D. Medina Lopez (CEA-Saclay), Y. Chassagneux (LPENS, ENS Paris), A. Narita (OIST), C. Voisin (Ecole Normale Supérieure - CNRS), S. Campidelli (CEA Saclay), D. Beljonne (MONS Univeristy), and J. Lauret (ENS Paris Saclay)

Top Down and Bottom Up Synthesized Graphene Quantum Dots As Nanothermometers For *In Vitro* Imaging

B. H. Lee, R. L. McKinney (Texas Christian University), M. T. Hasan (Tuskegee University), and A. V. Naumov (Texas Christian University)

(Invited) Synthesis and Characterizations of Highly Fluorescent Nanographene Molecules

A. Narita (Okinawa Institute of Science and Technology Graduate University, Max Planck Institute for Polymer Research)

Photoluminescence Brightening of Single-Walled Carbon Nanotubes Using Graphene Quantum Dots

S. H. Sajjadi, S. J. Wu, M. Reggente (École Polytechnique Fédérale de Lausanne), E. K. Goharshadi (Ferdowsi University Mashhad), N. Sharif (École Polytechnique Fédérale de Lausanne), H. Ahmadzadeh (Ferdowsi University Mashhad), and A. A. Boghossian (École Polytechnique Fédérale de Lausanne)

(Invited) CRISPR-Cas9-Based Cancer Therapy Delivery By Carbon Nanomaterials

A. V. Naumov (Texas Christian University)