



HAL
open science

Radiolysis on electrolytes of batteries: an efficient and quick screening for a fast selection of electrolytes

Yanis Levieux-Souid, Jean-Frédéric Martin, Philippe Moreau, Nathalie Herlin-Boime, Sophie Le Caër

► To cite this version:

Yanis Levieux-Souid, Jean-Frédéric Martin, Philippe Moreau, Nathalie Herlin-Boime, Sophie Le Caër. Radiolysis on electrolytes of batteries: an efficient and quick screening for a fast selection of electrolytes. 13ème Symposium France-Japon sur les Batteries au Lithium et Nouvelles Générations, Jul 2022, Provins, France. cea-03714032

HAL Id: cea-03714032

<https://cea.hal.science/cea-03714032>

Submitted on 5 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Radiolysis on electrolytes of batteries: an efficient and quick screening for a fast selection of electrolytes

Yanis Levieux-Souid,^a Jean-Frédéric Martin,^b Philippe Moreau,^c Nathalie Herlin-Boime^a and Sophie Le Caër^{a*}

^a NIMBE, UMR 3685 CEA, CNRS, Université Paris-Saclay, CEA Saclay F-91191 Gif-sur-Yvette Cedex, France.

^b CEA, LITEN, DEHT, STB, LM F-38054 Grenoble Cedex 9, France.

^c Institut des Matériaux Jean Rouxel, IMN, Université de Nantes, CNRS, 2 rue de la Houssinière, BP 32229 F-44322 Nantes Cedex 3, France

*email: sophie.le-caer@cea.fr

Recently we demonstrated that the highly reactive species generated by radiolysis (the chemical reactivity induced by the interaction between matter and ionizing radiation) of an electrolyte are the same as those generated during the cycling of lithium-ion batteries (LIBs) using similar solvents.^[1-4] Furthermore, radiolysis generated these species in measurable amounts at time scales significantly shorter than that of electrochemical cycling in a battery cell (minutes vs. days) and enabled proposing reaction mechanisms and measuring rate constants.^[1-3]

In order to provide safe and reliable lithium-ion batteries, the knowledge of aging phenomena is crucial. Among the different parts of the system prone to aging, the behavior of the electrolyte is a key parameter. The screening of different electrolytes is then of major interest. However, screening studies on electrolytes are time-consuming because they require numerous cycles in batteries; they are thus barely found. We demonstrate here that radiation chemistry is a valuable tool to screen very quickly, within a few hours, various electrolytes. Comparison with literature data providing classification of batteries by their performance efficiency showed that the H₂ production under electrolyte irradiation allowed obtaining the same stability order. It allowed a direct comparison with experiments performed on pouch-cells. Therefore, we evidenced that irradiation experiments on electrolytes alone, lasting a few hours without any manufacturing operations such as the fabrication of electrochemical cells, makes it possible to reproduce trends observed during battery cycling. This work attests to the interest of radiolysis studies for the effective, reliable and fast screening of electrolytes.

References

- [1] D. Ortiz et al., *Nature Comm.*, **2015**, 6, 6950.
- [2] D. Ortiz et al., *ChemSusChem*, **2015**, 8, 3605.
- [3] D. Ortiz et al., *J. Power Sources*, **2016**, 326, 285.
- [4] F. Varenne et al., *Sustainable Energy & Fuels*, **2018**, 2, 2100.