



**HAL**  
open science

## Isotopic effect on the formation kinetics of Pu(IV) intrinsic colloids

Manon Cot Auriol, Matthieu Viro, Thomas Dumas, Olivier Diat, Denis  
Menut, Philippe Moisy, Sergey I. Nikitenko

► **To cite this version:**

Manon Cot Auriol, Matthieu Viro, Thomas Dumas, Olivier Diat, Denis Menut, et al.. Isotopic effect on the formation kinetics of Pu(IV) intrinsic colloids. Plutonium Futures 2022, Sep 2022, Avignon, France. cea-03779851

**HAL Id: cea-03779851**

**<https://cea.hal.science/cea-03779851v1>**

Submitted on 6 Nov 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Isotopic effect on the formation kinetics of Pu(IV) intrinsic colloids

M. Cot-Auriol [1], M. Virost [1], T. Dumas [2], O. Diat [1], D. Menut [3],  
P. Moisy [2], S.I. Nikitenko [1]

[1] ICSM, Univ Montpellier, CEA, CNRS, ENSCM, Marcoule, France.

[2] CEA of Marcoule, BP17171, 30207 Bagnols-sur-Cèze, France.

[3] Synchrotron SOLEIL, MARS beamline, l'Orme des Merisiers, Saint Aubin BP 48, 91192 Gif-sur-Yvette, France.

Corresponding author: manon.cotauriol@cea.fr

Fundamental knowledge of the intrinsic colloids of plutonium is a major issue in predicting their behavior in the environment.[1,2] It has been demonstrated that these species are spherical nanoparticles measuring *ca.* 2 nm with a local structure similar to the one of bulk PuO<sub>2</sub>. [3,4] Recent studies have questioned the contribution of small Pu clusters in the formation mechanism of Pu(IV) intrinsic colloids.[5] The current work draws attention to this issue by investigating an unprecedented kinetic isotopic effect (KIE) observed during the preparation of Pu(IV) colloids in light (H<sub>2</sub>O) or heavy (D<sub>2</sub>O) water.

Plutonium colloids have been synthesized by the dilution of acidic Pu(IV) solutions in H<sub>2</sub>O and D<sub>2</sub>O. Time dependent measurements carried out with UV-Vis absorption spectroscopy demonstrated a much lower formation rate of the Pu(IV) colloids in D<sub>2</sub>O compared to H<sub>2</sub>O (Figure 1a). The observed KIE was attributed to the difference in zero-point energies of O-H and O-D bonds. This finding has confirmed that the formation of Pu(IV) colloids is limited by the rupture of O-H bonds during hydrolysis and polymerization of Pu(IV) species. In addition, formation of some unstable intermediate Pu(IV) species was observed during colloid formation. A thorough investigation of this intermediate using synchrotron SAXS and XAS techniques revealed the presence of an hexanuclear Pu(IV) cluster of quasi-spherical shape measuring *ca.* 0.8 nm and agreeing with the literature (Figure 1b). [6,7]




Figure 1: (a) UV-Vis absorption spectra acquired during the synthesis of 5 mM Pu(IV) hydrolytic colloids in H<sub>2</sub>O (red lines) and D<sub>2</sub>O (blue lines). (b)  $k^3$ -weighted Fourier transform (FT) of the intermediate species observed in H<sub>2</sub>O by UV-Vis absorption spectroscopy (dashed circle).[6]

- [1] C. Walther *et al.*, *Chem. Rev.* **113**, 995–1015 (2013).
- [2] A.Y. Romanchuk *et al.*, *Front. Chem.* **8**, 630 (2020).
- [3] E. Dalodière *et al.*, *Sci Rep* **7**, 43514 (2017).
- [4] C. Micheau *et al.*, *Environ. Sci.: Nano* **7**, 2252 (2020).
- [5] K.E. Knope *et al.*, *Inorg. Chem.* **52**, 6770-6772 (2013).
- [6] C. Tamain *et al.*, *Eur. J. Inorg. Chem.* **2016**, 3536-3540 (2016).
- [7] T. Dumas *et al.*, *J. Synchrotron Rad.* **29**, 30-36 (2022).