



HAL
open science

In situ 3D observations of a core–shell volume transition in an Ni₃Fe nanocrystal using Bragg coherent X-ray diffraction imaging

C. Chatelier, C. Atlan, M. Dupraz, Ni Li, E. Rabkin, S. Labat, J. Eymery, M.I. Richard

► To cite this version:

C. Chatelier, C. Atlan, M. Dupraz, Ni Li, E. Rabkin, et al.. In situ 3D observations of a core–shell volume transition in an Ni₃Fe nanocrystal using Bragg coherent X-ray diffraction imaging. *Acta Crystallographica Section A: Foundations and Advances* [2014-..], 2022, 78 (a2), pp.e766-e766. 10.1107/S2053273322090192 . cea-04066341

HAL Id: cea-04066341

<https://cea.hal.science/cea-04066341>

Submitted on 28 Aug 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

MS40-2-6 In situ 3D observations of a core-shell volume transition in an Ni₃Fe nanocrystal using Bragg coherent X-ray diffraction imaging

#MS40-2-6

C. Chatelier¹, C. Atlan¹, M. Dupraz¹, N. Li¹, E. Rabkin², S. Labat³, J. Eymery⁴, M.I. Richard¹

¹CEA Grenoble - ESRF - Grenoble (France), ²Technion-Israel Institute of Technology - Haifa (Israel), ³IM2NP - Aix Marseille Université - Marseille (France), ⁴CEA Grenoble - Grenoble (France)


Abstract

Coherent X-ray Diffraction Imaging (CDI) [1] allows for the *in situ* 3D mapping of the structural changes during phase transitions and chemical reactions, with a fair spatial resolution of the order of magnitude of 10 nm [2]. Using this technique in multi-Bragg conditions (Bragg CDI on multiple reflections), it was possible to follow the structural evolution (change in morphology, variation of the displacement and strain fields) as a function of temperature in a 300 nm Ni₃Fe nanocrystal. During this volume transition, which happens between 500°C and 600°C, a core-shell structure is progressively observed with two distinct lattice parameters (difference of 0.39%, corresponding to two different Ni:Fe ratios), leading to highly strained and coherent domains. A rounding of the particle is also observed, with a disappearing of the initial facets. Static molecular calculations (LAMMPS) of different systems, such as faceted and core-shell nanoparticles, complete this work and allow for the confirmation of a core-shell structure. This study shows the perfect ability of modern phase retrieval methods to reconstruct highly strained systems such as core-shell nanoparticles.

References

- [1] I. Robinson and R. Harder, Coherent X-Ray Diffraction Imaging of Strain at the Nanoscale, *Nat. Mater.* 8, 291 (2009).
- [2] S. Labat, M.-I. Richard, M. Dupraz, M. Gailhanou, G. Beutier, M. Verdier, F. Mastropietro, T. W. Cornelius, T. U. Schüllli, J. Eymery, and O. Thomas, Inversion Domain Boundaries in GaN Wires Revealed by Coherent Bragg Imaging, *ACS Nano* 9, 9210 (2015).

BCDI analysis (a) before and (b) after annealing



LAMMPS simulations: (a) faceted and (b) core-shell

