



A unified description of the atomic physics for electron Fokker-Planck calculations

Y Peysson, D Mazon, J Bielecki, D Dworak, K Król, A Jardin, M Scholz, J Walkoviak, J Decker

► To cite this version:

Y Peysson, D Mazon, J Bielecki, D Dworak, K Król, et al.. A unified description of the atomic physics for electron Fokker-Planck calculations. 21st Meeting on Atomic Processes in Plasmas, May 2023, Vienne, Austria. cea-04108114

HAL Id: cea-04108114

<https://cea.hal.science/cea-04108114v1>

Submitted on 26 May 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

DE LA RECHERCHE À L'INDUSTRIE



EPFL



www.cea.fr

A unified description of the atomic physics for electron Fokker-Planck calculations

***Y. Peysson^{1*}, D.Mazon¹
J. Bielecki², D. Dworak², K. Król², A. Jardin², M.
Scholz², J. Walkowiak², J. Decker³***

¹*CEA/IRFM, F-13108, Saint-Paul-lez-Durance, France*

²*Institute of Nuclear Physics, Polish Academy of Sciences, PL-
31342, Kraków, Poland*

³*EPFL, Swiss Plasma Center, CH-1015 Lausanne, Switzerland*

* yves.peysson@cea.fr

Growing interest for studying the impact of partially ionized high-Z atoms on fast electron dynamics in magnetized tokamak plasmas

- ***Passive sources → plasma-wall interactions***

- Tungsten-based plasma-facing components (planned to be used in the ITER tokamak reactor for reducing tritium retention)
- Metallic objects in contact with the scrape-off layer like : RF antennas (Copper, Iron,...)

- ***Active sources → massive gas injection***


- Mitigation of a runaway electron beam after a major disruption in the ITER tokamak reactor (Argon, Neon, ...)

- High-Z atoms have the property to be partially ionized from the core to the edge of the plasma.
- The distribution of the screened charge $Z_{0,s}$ (for an atom of species s) is therefore a function of the local plasma temperature.
- Fast electrons will interact differently with bound electrons depending upon their kinetic energy → **the partial screening effect**.
- Calculations that describe the fast electron dynamics in momentum and configuration spaces must incorporate this new effect *in the electron-ion collision Fokker-Planck operator, for all species present in the plasma and whatever their ionization state :*
 - Very wide range of plasma temperatures : 10 eV (*disruption*) to 10 keV (*ITER*)
 - Very wide range of electron kinetic energies : 10-100 eV to ~10 MeV (*runaway electrons*)

Distribution of ionization states in a Maxwellian plasma


Post-disruption

OPEN-ADAS, tungsten ($Z_s = 74$)




H. P. Summers, Technical report, ADAS (2004)


Impact of partially ionized atoms on magnetized tokamak plasma physics




Impact of partially ionized atoms on magnetized tokamak plasma physics




Impact of partially ionized atoms on magnetized tokamak plasma physics




Impact of partially ionized atoms on magnetized tokamak plasma physics




Impact of partially ionized atoms on plasma physics



Impact of partially ionized atoms on magnetized tokamak plasma physics



Impact of partially ionized atoms on magnetized tokamak plasma physics



- Partially ionized atoms are like in vacuum : $\langle r \rangle \ll n_e^{-1/3}$
- Partially ionized atoms are principally in the ground state : $\nu_{coll} \times \tau_{lifetime} \ll 1$

▪ Elastic scattering and screening effect

- e-i collisions dominated by small angle deflections → 1st Born approximation
- Screening effect : $Z_{0,s} \rightarrow Z_{0,s} - F(\mathbf{q})$ where $F(\mathbf{q})$ is the *atomic form factor* (\mathbf{q} is the recoil momentum)
- $F(\mathbf{q})$ is the Fourier transform of the *density of bound electrons* $\rho(r)$

$$F_{Z_{0,s}}(\mathbf{q}) \equiv \int_V \exp(-i\mathbf{q} \cdot \mathbf{r}/\hbar) \rho_{Z_{0,s}} d\mathbf{r}$$

▪ Inelastic scattering by atomic excitation

- Bethe's stopping-power for energy losses
- *Mean excitation energy* $\langle \hbar\omega_s \rangle$, function of the *density of bound electrons* $\rho(r)$

$$\langle \hbar\omega_{Z_{0,s}} \rangle = (1/Z_{0,s}) \sum_{ik} f_{ik} \ln(\hbar\omega_{ik})$$

U. Fano and J. W. Cooper, Rev. Mod. Phys., 40(3) (1968) 441-507



The density of bound electrons $\rho(r)$

- Incorporating the physics of partially ionized high-Z atoms in Fokker-Planck calculations requires to know *the density of bound electrons for each species and each ionization state*.
- Fast and accurate time-dependent Fokker-Planck calculations for wide parameter ranges, $T_e \in \{10-10^4\}$ eV and $E_c \in \{10^{-2}-10^3\}$ keV, requires analytical expressions for the partial screening in the collision operator → *simplified analytic atomic models*
- **Density of bound electrons :**
 - *Ab initio quantum relativistic calculations* : (DFT) Density Functional Theory or (MCDHF) Multi-Configuration Dirac-Hartree-Fock model
 - *Simplified analytic atomic models* : Thomas-Fermi or Yukawa models → **Problem : none of them able to describe consistently the density of bound electrons for all possible ionization states present in the plasma, depending on the tokamak operating regime.**

The density of bound electrons : ab initio codes


DFT : Density functional theory

MCDHF : Multi-Configuration Dirac-Hartree-Fock model




Simple atomic models for Fokker-Planck calculations in magnetized tokamak plasmas

Weakly ionized atoms
↓
Thomas-Fermi model



Strongly ionized atoms
↓
Yukawa model



P. Amore, et al., . Applied Math. and Comp., 232 (2014) 929–943

Generalization of the Molière's method initially developed for the Thomas-Fermi model

- **Atomic potential :** $4\pi\epsilon_0 r U_{Z_{0,s}}(r) = -Z_{0,s} - \sum_i A_{Z_{0,s,i}} \exp(-\lambda_{Z_{0,s,i}} r)$
- Bound electron density by solving the Poisson equation $\nabla^2 U_{Z_{0,s}} = -\rho_{Z_{0,s}}/\epsilon_0$:

$$\bar{\rho}_{Z_{0,s}}(\bar{r}) = \frac{Z_s - Z_{0,s}}{4\pi\bar{r}} \sum_i \bar{\lambda}_{Z_{0,s,i}}^2 \bar{A}_{Z_{0,s,i}} \exp(-\bar{\lambda}_{Z_{0,s,i}} \bar{r})$$

$$\bar{r} = r/a_0 \quad \bar{\rho}_{Z_{0,s}}(\bar{r}) = \rho_{Z_{0,s}}(\bar{r}) a_0^3 \quad \bar{\lambda}_{Z_{0,s,i}} \equiv \lambda_{Z_{0,s,i}} a_0 \quad \bar{A}_{Z_{0,s,i}} = A_{Z_{0,s,i}} / (Z_s - Z_{0,s})$$

- **Analytic atomic form factor :**

$$F_{Z_{0,s}}(\bar{q}) = (Z_s - Z_{0,s}) \sum_i \frac{\bar{A}_{Z_{0,s,i}}}{1 + (\bar{q} \bar{a}_{Z_{0,s,i}}/2)^2}$$

$$\bar{q} = q/(m_e c) \quad \bar{a}_{Z_{0,s,i}} \equiv 2\bar{\lambda}_{Z_{0,s,i}}^{-1}/\alpha$$

$$\sum_i \bar{A}_{Z_{0,s,i}} = 1 \quad \text{guarantees that} \quad F_{Z_{0,s}}(0) = Z_s - Z_{0,s} = N_s$$

Von Gert Moliere. Zeitschrift Naturforschung, 2a (1947) 133–145

- **Determination of $(\bar{A}_{Z_{0,s,i}}, \bar{\lambda}_{Z_{0,s,i}})$ is an ill-posed problem.** Standard non-linear regression techniques are useless.
- **Method of moments :** $\langle \bar{r}^l \rangle = \langle \bar{r}^l \rangle^{num}$ where $\langle \bar{r}^l \rangle^{num}$ is determined from ab initio quantum numerical calculations (DFT or MCDHF) → Uniqueness of the solution if it exists.
- Defining $\mathcal{R}_l \equiv \langle \bar{r}^l \rangle / (l + 1)!$ where

$$\mathcal{R}_l = \sum_i \bar{A}_{Z_{0,s,i}} \bar{\lambda}_{Z_{0,s,i}}^{-l}$$



$$\mathcal{R}_l^{num} = \frac{1}{(l + 1)!} \frac{4\pi}{(Z_s - Z_{0,s})} \int_0^\infty \bar{r}^{l+2} \bar{\rho}_{Z_{0,s}}^N(\bar{r}) d\bar{r}$$

- **2l equations to be solved, where l is the number of exponentials**
- **If no solution found $l \rightarrow l - 1$**

F. Salvat, et al., Phys. Rev. A, 36(2) (1987) 467–474



Density of bound electrons (up to 3 exponentials)

Neutral tungsten $Z_{0,s} = 0$




Density of bound electrons (up to 3 exponentials)

Partially ionized tungsten $Z_{0,s} = 42$




Atomic form factor (up to three exponentials)

Neutral tungsten $Z_{0,s} = 0$




Partially ionized tungsten $Z_{0,s} = 42$



$$F_{Z_{0,s}}(0) = Z_s - Z_{0,s} = N_s$$

Coefficient of determination R^2 for tungsten




Electron-ion Fokker-Planck collision operator :

$$\mathbf{C}_{e,Z_{0,s}}(f_e, f_{Z_{0,s}}) \simeq -\nabla_p (\mathbf{A}_{Z_{0,s}} f_e(t, \mathbf{x}, \mathbf{p})) + \nabla_p \nabla_p (\mathbb{D}_{Z_{0,s}} f_e(t, \mathbf{x}, \mathbf{p}))$$

Friction vector : $\mathbf{A}_{Z_{0,s}} = \int d^3 \mathbf{p}_s f_{Z_{0,s}}(t, \mathbf{x}, \mathbf{p}_s) \int d\Omega \frac{d\sigma_{e,Z_{0,s}}}{d\Omega} u_s \Delta \mathbf{p}$



Diffusion tensor : $\mathbb{D}_{Z_{0,s}} = \frac{1}{2} \int d^3 \mathbf{p}_s f_{Z_{0,s}}(t, \mathbf{x}, \mathbf{p}_s) \int d\Omega \frac{d\sigma_{e,Z_{0,s}}}{d\Omega} u_s \Delta \mathbf{p} \Delta \mathbf{p}^T$



Elastic scattering : Mott-Rutherford cross-section and partial screening effect

Coulomb collision (1st Born approximation) :

$$\frac{d\sigma_{e,Z_{0,s}}}{d\Omega} = Z_{0,s}^2 \frac{r_e^2}{4} \frac{(1-x^2)\bar{p}^2 + 1}{\bar{p}^4 x^4}$$

Elastic scattering : $|\Delta \mathbf{p}| = 2 |\mathbf{p}| \sin \frac{\theta}{2} = 2 |\mathbf{p}| x \longrightarrow \bar{q} = 2\bar{p} \sin (\theta/2)$

Partial screening effect : $Z_{0,s}^2 \rightarrow |Z_s - F_{Z_{0,s}}(\bar{q})|^2$

$$\int_{1/\Lambda_{e,Z_{0,s}}}^1 Z_{0,s}^2 \frac{dx}{x} \longrightarrow \int_{1/\Lambda_{e,Z_{0,s}}}^1 |Z_s - F_{Z_{0,s}}(\bar{q})|^2 \frac{dx}{x}$$

Coulomb logarithm	Screening function
$\int_{1/\Lambda_{e,Z_{0,s}}}^1 Z_s - F_{Z_{0,s}}(\bar{q}) ^2 \frac{dx}{x} \equiv$	$Z_{0,s}^2 \ln \Lambda_{e,Z_{0,s}}$ + $g_{Z_{0,s}}(\bar{p})$

L. Hesslow, et al., J. Plasma Phys., 84(6) (2018) 905840605

Collision operator as a divergence of a flux in momentum space (LUKE code)

$$\frac{df_e}{dt}|_{coll} \equiv \sum_s \sum_{Z_{0,s}} \mathbf{C}_{e,Z_{0,s}}(f_e, f_{Z_{0,s}}) + \mathbf{C}_{e,e}(f_e, f_e) \equiv \nabla_{\mathbf{p}} \cdot \mathbf{S}_{\mathbf{p}}^{coll}(f_e)$$

Coordinate system

(p, ξ, φ)

$$\begin{cases} S_p^{coll} = -D_{pp}^{coll} \frac{\partial f_e}{\partial p} + \frac{\sqrt{1 - \xi^2}}{p} D_{p\xi}^{coll} \frac{\partial f_e}{\partial \xi} + F_p^{coll} f_e \\ S_\xi^{coll} = -D_{\xi p}^{coll} \frac{\partial f_e}{\partial p} + \frac{\sqrt{1 - \xi^2}}{p} D_{\xi\xi}^{coll} \frac{\partial f_e}{\partial \xi} + F_\xi^{coll} f_e \end{cases}$$

$$D_{p\xi}^{coll} = D_{\xi p}^{coll} = F_\xi^{coll} = 0 \quad \text{by local axisymmetry}$$

Inelastic scattering

$$D_{pp}^{coll}(e-i) \ll D_{pp}^{coll}(e-e)$$

$$F_p^{coll}(e-i) \ll F_p^{coll}(e-e)$$

Elastic scattering

The transformation

$$Z_{0,s}^2 \ln \Lambda_{e,Z_{0,s}} \rightarrow Z_{0,s}^2 \ln \Lambda_{e,Z_{0,s}} + g_{Z_{0,s}}(\bar{p})$$

is applied to $D_{\xi\xi}^{coll}(e-i)$ only.

Screening function using the multi-Yukawa model

$$g_{Z_{0,s}}(\bar{p}) = Z_s (Z_s - Z_{0,s}) \sum_i \bar{A}_{0,s,i} \ln \left(1 + (\bar{p} \bar{a}_{Z_{0,s,i}})^2 \right)$$

$$- \frac{(Z_s - Z_{0,s})^2}{2} \sum_i \bar{A}_{0,s,i}^2 \left(\frac{(\bar{p} \bar{a}_{Z_{0,s,i}})^2}{1 + (\bar{p} \bar{a}_{Z_{0,s,i}})^2} + \ln \left(1 + (\bar{p} \bar{a}_{Z_{0,s,i}})^2 \right) \right)$$


$$+ 2 (Z_s - Z_{0,s})^2 \sum_i \sum_{j > i} \bar{A}_{0,s,i} \bar{A}_{0,s,j} \boxed{\tilde{g}_{Z_{0,s},2,i,j}(\bar{p})}$$

$$\begin{aligned} \tilde{g}_{Z_{0,s},2,i,j}(\bar{p}) = & - \frac{1}{2} \frac{a_{Z_{0,s,i}}^2 + a_{Z_{0,s,j}}^2}{a_{Z_{0,s,j}}^2 - a_{Z_{0,s,i}}^2} \ln \left| \frac{1 + \bar{p}^2 \bar{a}_{Z_{0,s,j}}^2}{1 + \bar{p}^2 \bar{a}_{Z_{0,s,i}}^2} \right| \\ & - \frac{1}{4} \ln \left| 1 + \bar{p}^2 \left(\bar{a}_{Z_{0,s,i}}^2 + \bar{a}_{Z_{0,s,j}}^2 \right) + \bar{p}^4 \bar{a}_{Z_{0,s,i}}^2 \bar{a}_{Z_{0,s,j}}^2 \right| \\ & + \frac{\bar{a}_{Z_{0,s,i}}^2 + \bar{a}_{Z_{0,s,j}}^2}{4 \left| \bar{a}_{Z_{0,s,i}}^2 - \bar{a}_{Z_{0,s,j}}^2 \right|} \\ & \times \left[\ln \left| \frac{2 \bar{a}_{Z_{0,s,i}}^2 \bar{a}_{Z_{0,s,j}}^2 \bar{p}^2 + \left(\bar{a}_{Z_{0,s,i}}^2 + \bar{a}_{Z_{0,s,j}}^2 \right)}{2 \bar{a}_{Z_{0,s,i}}^2 \bar{a}_{Z_{0,s,j}}^2 \bar{p}^2 + \left(\bar{a}_{Z_{0,s,i}}^2 + \bar{a}_{Z_{0,s,j}}^2 \right)} - \left| \bar{a}_{Z_{0,s,i}}^2 - \bar{a}_{Z_{0,s,j}}^2 \right| \right| \right. \\ & \quad \left. - \ln \left| \frac{\left(\bar{a}_{Z_{0,s,i}}^2 + \bar{a}_{Z_{0,s,j}}^2 \right) - \left| \bar{a}_{Z_{0,s,i}}^2 - \bar{a}_{Z_{0,s,j}}^2 \right|}{\left(\bar{a}_{Z_{0,s,i}}^2 + \bar{a}_{Z_{0,s,j}}^2 \right) + \left| \bar{a}_{Z_{0,s,i}}^2 - \bar{a}_{Z_{0,s,j}}^2 \right|} \right| \right] \end{aligned}$$


- Fully analytic expression
- Arbitrary number of exponentials

Screening function : comparison between multi-Yukawa model and numerical quantum calculations

Neutral tungsten $Z_{0,s}=0$



Partially ionized tungsten $Z_{0,s}=42$



Drag force by atomic excitation

- Bethe's stopping power formula where the energy E loss is :

$$-\frac{dE}{dx} \Big|_{Z_{0,s}} = 4\pi r_e^2 n_{Z_{0,s}} (Z_s - Z_{0,s}) \frac{m_e c^2}{\beta^2} [\ln B_{Z_{0,s}} - \beta^2]$$

$$B_{Z_{0,s}} = \frac{\sqrt{2}\gamma\beta\sqrt{\gamma-1}}{\langle \hbar\omega_{Z_{0,s}} \rangle / m_e c^2}$$

Atomic physics is described by
the mean excitation energy

- Since the energy loss over a distance Δx is equivalent of the work of an effective drag force,

$$F_p^{excitation}(p) = - \sum_s \sum_{Z_{0,s}} \frac{dE}{dx} \Big|_{Z_{0,s}}$$

- Incorporation in LUKE Fokker-Planck code : $F_p^{coll} = F_p^{e-i} + F_p^{e-e} + F_p^{excitation} + F_p^{ALD}$

$$S_p^{coll} = -D_{pp}^{coll} \frac{\partial f_e}{\partial p} + \frac{\sqrt{1-\xi^2}}{p} D_{p\xi}^{coll} \frac{\partial f_e}{\partial \xi} + F_p^{coll} f_e$$

H. Bethe and W. Heitler. Proceedings of the Royal Society of London, volume A146, (1934) 83–112

Mean excitation energy

- *Mean excitation energy* → $\langle \hbar\omega_{Z_{0,s}} \rangle = (1/Z_{0,s}) \sum_{ik} f_{ik} \ln (\hbar\omega_{ik})$

- f_{ik} : dipole oscillator strength of the transition
- ω_{ik} : dipole frequency for the transition between $|i\rangle$ and $|k\rangle$ quantum states

- **Exponential interpolation** : $\ln \langle \hbar\omega_{Z_{0,s}} \rangle \simeq \ln \left(\frac{I_H}{10} Z_s \right) \frac{Z_{0,s}}{Z_s - 1} + \ln (10Z_s)$
- **Local plasma approximation (LPA)** :

$$\begin{aligned} \ln \langle \hbar\omega_{Z_{0,s}} \rangle &= \sum_i \bar{\lambda}_{Z_{0,s},i}^2 \bar{A}_{Z_{0,s},i} \int_0^\infty \bar{r} \exp(-\bar{\lambda}_{Z_{0,s},i} \bar{r}) \\ &\times \ln \left(\sqrt{2\alpha^2 m_e c^2} \sqrt{\frac{Z_s - Z_{0,s}}{\bar{r}}} \sum_j \bar{\lambda}_{Z_{0,s},j}^2 \bar{A}_{Z_{0,s},j} \exp(-\bar{\lambda}_{Z_{0,s},j} \bar{r}) \right) d\bar{r} \end{aligned}$$

S. Rosendorff and H. G. Schlaile. Phys. Rev. A, 40(12) (1989) 6892–6896

J. Linhard and M. Scharff. Dan. Mat. Fys. Medd, 27(15) (1953)

W.K. Chu and D. Powers. Phys. Letters, 40A(1) (1972) 23–24, 1972

Mean excitation energy


- **Variational quantum description** : $\langle \hbar\omega_{Z_{0,s}} \rangle^2 = 2 \frac{K_0}{\langle \bar{r}^2 \rangle} \alpha^2 m_e c^2$

The *averaged kinetic energy of the cloud of bound electrons* K_0 is determined using the virial theorem (non-relativistic limit) : $2K_0 \simeq \langle U_{Z_{0,s}} \rangle / N_s$


$$\begin{aligned} \langle \hbar\omega_{Z_{0,s}} \rangle &= \frac{\alpha^2 m_e c^2 \sqrt{N_s}}{\sqrt{6 \sum_i \bar{\lambda}_{Z_{0,s,i}}^{-2} \bar{A}_{Z_{0,s,i}}}} \\ &\times \sqrt{\sum_i \bar{A}_{Z_{0,s,i}}^2 \frac{\bar{\lambda}_{Z_{0,s,i}}}{2} + \sum_i \sum_{j \neq i} \bar{A}_{Z_{0,s,i}} \bar{A}_{Z_{0,s,j}} \frac{\bar{\lambda}_{Z_{0,s,i}}^2}{\bar{\lambda}_{Z_{0,s,i}} + \bar{\lambda}_{Z_{0,s,j}}}} \end{aligned}$$

- Single exponential limit : $\langle \hbar\omega_{Z_{0,s}} \rangle = \alpha^2 m_e c^2 \sqrt{\bar{\lambda}_{Z_{0,s,1}}^3 N_s / 12}$
- Thomas-Fermi model for neutral atoms : $\bar{\lambda}_{Z_{0,s,1}} = 1.13 Z_s^{1/3}$
 $\langle \hbar\omega_{Z_{0,s}} \rangle = 9.44 Z_s \text{ eV} \longrightarrow \text{Bloch relation } \sim 10 Z_s \text{ eV is well retrieved}$

Mean excitation energy for argon



Mean excitation energy for tungsten



Conclusions

- A unified atomic model description allowing analytical formulations of the partial screening effect in Fokker-Planck calculations has been carried out, such that the impact of high-Z elements in hot tokamak plasmas on fast electron physics may be accurately described for any tokamak regimes.
- The atomic potential is expressed as a series of Yukawa potentials. *Extension of the Molière's method initially derived for the Thomas-Fermi model.* A general numerical procedure has been derived for the model calibration against ab initio quantum relativistic numerical calculations (DFT, MCDHF, ...).
- The robustness of the multi-Yukawa description has been evaluated for tungsten element as a function of the number of exponentials → best results : 3 exponentials
- Analytic forms of the elastic and inelastic in diffusion and friction terms of the Fokker-Planck collision operators with the multi-Yukawa atomic model have been derived.
- A robust analytical estimate of the *mean excitation energy* has been obtained from non-relativistic quantum calculations, giving a consistent dependence with the ion charge. Applicable to any element including tungsten.

- Case of three exponentials expandable to any number of exponentials :
 - $\bar{\lambda}$ should be **a real and positive solution** of the polynomial equation :

$$\bar{\lambda}^3 - X_1 \bar{\lambda}^2 + X_2 \bar{\lambda} - X_3 = 0$$



- Defining $\mathbf{X} = \begin{bmatrix} X_1 \\ X_2 \\ X_3 \end{bmatrix} = \begin{bmatrix} \bar{\lambda}_{Z_{0,s,1}} + \bar{\lambda}_{Z_{0,s,2}} + \bar{\lambda}_{Z_{0,s,3}} \\ \bar{\lambda}_{Z_{0,s,1}} \bar{\lambda}_{Z_{0,s,2}} + \bar{\lambda}_{Z_{0,s,2}} \bar{\lambda}_{Z_{0,s,3}} + \bar{\lambda}_{Z_{0,s,1}} \bar{\lambda}_{Z_{0,s,3}} \\ \bar{\lambda}_{Z_{0,s,1}} \bar{\lambda}_{Z_{0,s,2}} \bar{\lambda}_{Z_{0,s,3}} \end{bmatrix}$

the coefficients $\bar{A}_{Z_{0,s,i}}$ are given by the matrix relation : $\mathbf{A} = \mathbb{N}^{-1} \mathbb{M} \mathbf{X}$
where

$$\mathbf{A} = \begin{bmatrix} \bar{A}_{Z_{0,s,1}} \\ \bar{A}_{Z_{0,s,2}} \\ \bar{A}_{Z_{0,s,3}} \end{bmatrix} \quad \mathbb{M} = \begin{bmatrix} 1 & -\mathcal{R}_1^{num} & \mathcal{R}_2^{num} \\ \mathcal{R}_1^{num} & -\mathcal{R}_2^{num} & \mathcal{R}_3^{num} \\ \mathcal{R}_2^{num} & -\mathcal{R}_3^{num} & \mathcal{R}_4^{num} \end{bmatrix} \quad \mathbb{N} = \begin{bmatrix} \bar{\lambda}_{Z_{0,s,1}} & \bar{\lambda}_{Z_{0,s,2}} & \bar{\lambda}_{Z_{0,s,3}} \\ 1 & 1 & 1 \\ \bar{\lambda}_{Z_{0,s,1}}^{-1} & \bar{\lambda}_{Z_{0,s,2}}^{-1} & \bar{\lambda}_{Z_{0,s,3}}^{-1} \end{bmatrix}$$

- If no three real and positive $\bar{\lambda}$ solutions are found \rightarrow less number of exponentials

Collision terms in LUKE Fokker-Planck solver



Screening function from numerical quantum calculations

$$g_{Z_{0,s}}^{num}(\bar{p}) = (Z_s^2 - Z_{0,s}^2) (\ln(2\bar{p}/\alpha) + \gamma_{EM} - 1) + 2Z_s N_s \hat{I}_{1,Z_{0,s}} + N_s^2 \left(\frac{1}{2} - \hat{I}_{2,Z_{0,s}} \right)$$

$$\hat{I}_{1,Z_{0,s}} \equiv \frac{4\pi}{N_s} \int_0^\infty \bar{\rho}_{Z_{0,s}}^{num}(\bar{r}_1) \bar{r}_1^2 \ln \bar{r}_1 d\bar{r}_1$$



DFT

$$\hat{I}_{2,Z_{0,s}} = \frac{4\pi^2}{N_s^2} \int_0^\infty \bar{\rho}_{Z_{0,s}}^{num}(\bar{r}_1) \hat{J}_{2,Z_{0,s}}(\bar{r}_1) \bar{r}_1 d\bar{r}_1$$

$$\hat{J}_{2,Z_{0,s}}(\bar{r}_1) = \int_0^\infty \bar{\rho}_{Z_{0,s}}^{num}(\bar{r}_2) \bar{r}_2 d\bar{r}_2 \left((\bar{r}_1 + \bar{r}_2)^2 \ln(\bar{r}_1 + \bar{r}_2) - (\bar{r}_1 - \bar{r}_2)^2 \ln|\bar{r}_1 - \bar{r}_2| \right)$$

Figure 1

where γ_{EM} is the Euler-Mascheroni constant.

Calculation valid for $\bar{p} \gg 1$ only.

L. Hesslow, et al., J. Plasma Phys., 84(6) (2018) 905840605