



New DDEP recommendations for the radioactive decay scheme of ^{137}Cs

Sylvain Leblond

► To cite this version:

Sylvain Leblond. New DDEP recommendations for the radioactive decay scheme of ^{137}Cs . ICRM 2023 - 23rd International Conference on Radionuclide Metrology and its Applications, Mar 2023, Bucarest, Romania. cea-04256925



HAL Id: cea-04256925

<https://cea.hal.science/cea-04256925v1>

Submitted on 24 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



New DDEP recommendations for the radioactive decay scheme of ^{137}Cs

S. Leblond





Discovery of an iconic radionuclide

- In 1941 at Berkeley
 - Margaret Melhase and Glenn Seaborg
 - First chemical extraction of radioactive Caesium
 - Produced by thermal neutron induced fission on ^{235}U

- Since then...



(1941)



(1950)



Discovery of an iconic radionuclide

- In 1941 at Berkeley
 - Margaret Melhase and Glenn Seaborg
 - First chemical extraction of radioactive Caesium
 - Produced by thermal neutron induced fission on ^{235}U

- Since then...
 - More than 80 000 publications in a wide range of applications



(1941)




(1950)



Decay overview of ^{137}Cs

- A β^- decay to ^{137}Ba
 - Populates ground & excited states
 - Half-life around 30 years
 - Total Q-value close to 1176 keV


- Emissions
 - Two main β^- branches
 - Ground state ($E_{\max} \sim 1176 \text{ keV}$)
 - Second excited state ($E_{\max} \sim 514 \text{ keV}$)
 - Iconic 662 keV gamma





Decay overview of ^{137}Cs

- A β^- decay to ^{137}Ba
 - Populates ground & excited states
 - Half-life around 30 years
 - Total Q-value close to 1176 keV
- Emissions
 - Two main β^- branches
 - Ground state ($E_{\max} \sim 1176 \text{ keV}$)
 - Second excited state ($E_{\max} \sim 514 \text{ keV}$)
 - Iconic 662 keV gamma




Surely decay data are well known...

Decay data status

- Latest ^{137}Cs decay data evaluations
 - DDEP
 - R.G. Helmer and V.P. Chechev (2006)
 - ENSDF
 - E. Browne and J.K. Tuli (2007)

	DDEP	ENSDF
$T_{1/2}$ (y)	30.05 (8)	30.08 (9)
$I_\beta(11/2^-)$ (%)	94.36 (28)	94.7 (2)
$I_\beta(3/2^+)$ (%)	5.64 (28)	5.3 (2)
I_γ (%)	84.99 (20)	85.1 (2)

New publications
Can it be improved?





Half-life of ^{137}Cs : overview

- Abundant bibliography
 - More than 75 publications
 - From 1948 to 2016
 - Different experimental methods
 - Many languages

- How to select a meaningful dataset?

NATURE April 3, 1948 Vol. 161

DECAY HALF-LIFE OF Cs^{137}

M. P. Glazunov, A. I. Grivkova, B. A. Zaitsev, and V. A. Kiselev

Translated from Atomnaya Energiya, Vol. 10, No. 6, pp. 622-623, June 1961

Original article submitted January 9, 1961

R. NEWBERY
E. WILLSHAW

In U^{232} fission the groups have a similar shape to those for U^{235} but are one unit lower on the mass scale. The maxima of the fission-yield curves are approximately the same in all cases, being approximately 6×10^{-2} . A detailed account of the investigation is to be published.

W. E. GRUMMITT G. WILKINSON

Atomic Energy Project, National Research Council, Chalk River, Ontario, Dec. 6.

Fission Products of U^{233}

Experiments have been described^{1,2} which...

¹ Grummitt, W. E., and Wilkinson, G., *Nature*, 158, 163 (1946).
² Grummitt, W. E., Gieron, J., Wilkinson, G., and Yaffe, L., *Can. J. Research*, 25, B, 1947.

Hageman, F., Katsin, I. L., Stadler, M. H., Ghiorso, A., and Seaborg, G. T., *Phys. Rev.*, 72, 252 (1947).

English, A. C., Cranshaw, T. E., Demers, P., Harvey, J. A., Hincks, E. P., Jelley, J. V., and May, A. N., *Nature*, 158, 163 (1946).

technical note

Appl. Radiat. Isot. Vol. 43, No. 7, pp. 949-951, 1992
Int. J. Radiat. Appl. Instrum. Part A
© Pergamon Press Ltd 1992. Printed in Great Britain
0883-2889/92 \$5.00 + 0.00

A Determination of the Half-life of ^{137}Cs

J.-J. GOSTELY

Institut d'Electrochimie et de Radiochimie de L'Ecole, Polytechnique Fédérale de Lausanne, 1015 Lausanne, Switzerland

Applied Radiation and Isotopes

ERGAMON Applied Radiation and Isotopes 56 (2002) 125-130
www.elsevier.com/locate/apradiso

Half-life measurements at the National Institute of Standards and Technology

M.P. Unterweger*

LSEVIER Contents lists available at ScienceDirect
Applied Radiation and Isotopes journal homepage: www.elsevier.com/locate/apradiso

Determination of ^{137}Cs half-life with an ionization chamber
édéric Juget*, Youcef Nedjadi, Thierry Buchillier, François Bochud, Claude Bailat

CrossMark



Half-life of ^{137}Cs : overview

- Abundant bibliography
 - More than 75 publications
 - From 1948 to 2016
 - Different experimental methods
 - Many languages

- How to select a meaningful dataset?
 - DDEP guidelines
 - Discard measurements without uncertainty estimation
 - One measurement per author / laboratory

NATURE April 3, 1948 Vol. 161

DECAY HALF-LIFE OF Cs^{137}

M. P. Glazunov, A. I. Grivkova, B. A. Zaitsev, and V. A. Kiselev

Translated from Atomnaya Energiya, Vol. 10, No. 6, pp. 622-623, June 1961

Original article submitted January 9, 1961

R. NEWBERY
E. WILLSHAW

In U^{232} fission the groups have a similar shape to those for U^{235} but are one unit lower on the mass scale. The maxima of the fission-yield curves are approximately the same in all cases, being approximately 6×10^{-2} . A detailed account of the investigation is to be published.

W. E. GRUMMITT
G. WILKINSON

Atomic Energy Project,
National Research Council,
Chalk River, Ontario.
Dec. 6.

Fission Products of U^{233}

Experiments have been described^{1,2} which...

¹ Grummitt, W. E., and Wilkinson, G., *Nature*, 158, 163 (1946).
² Grummitt, W. E., Gieron, J., Wilkinson, G., and Yaffe, L., *Can. J. Research*, 25, B, 1947.

Hageman, F., Katsin, I. L., Stadler, M. H., Ghiorso, A., and Seaborg, G. T., *Phys. Rev.*, 72, 252 (1947).

English, A. C., Cranshaw, T. E., Demers, P., Harvey, J. A., Hincks, E. P., Jelley, J. V., and May, A. N., *Nature*, 158, 163 (1946).

technical note

Appl. Radiat. Isot. Vol. 43, No. 7, pp. 949-951, 1992
Int. J. Radiat. Appl. Instrum. Part A
© Pergamon Press Ltd 1992. Printed in Great Britain
0883-2889/92 \$5.00 + 0.00

A Determination of the Half-life of ^{137}Cs

J.-J. GOSTELY
Institut d'Electrochimie et de Radiochimie de L'Ecole,
Polytechnique Fédérale de Lausanne, 1015 Lausanne,
Switzerland

Applied Radiation and Isotopes

HALF-LIFE MEASUREMENTS AT THE PTB

K.F.Walz, K. Debertin and H. Schrader

Physikalisch-Technische Bundesanstalt (PTB), Braunschweig

ERGAMON Applied Radiation and Isotopes www.elsevier.com/locate/apradiso

Half-life measurements at the National Institute of Standards and Technology

M.P. Unterweger*

LSEVIER Applied Radiation and Isotopes journal homepage: www.elsevier.com/locate/apradiso

Contents lists available at ScienceDirect

Determination of ^{137}Cs half-life with an ionization chamber
édéric Juget*, Youcef Nedjadi, Thierry Buchillier, François Bochud, Claude Bailat


CrossMark

Half-life of ^{137}Cs : a first dataset to work with

- Highly discrepant
 - 23 references considered
 - $T_{1/2}$ varying from 26 to 33 years
 - Uncertainties from 0.011 to 3 years
 - Statistical test of the distribution:

$$\chi^2 \sim 11 \gg \chi^2_{crit} \sim 1.9$$

$$\chi^2 = (n - 1) \frac{\sigma_{ext}^2}{\sigma_{int}^2}$$




Half-life of ^{137}Cs : a first dataset to work with

- Highly discrepant
 - 23 references considered
 - $T_{1/2}$ varying from 26 to 33 years
 - Uncertainties from 0.011 to 3 years
 - Statistical test of the distribution:
- $\chi^2 \sim 11 \gg \chi^2_{crit} \sim 1.9$
- $$\chi^2 = (n - 1) \frac{\sigma_{ext}^2}{\sigma_{int}^2}$$
- Unweighted average $T_{1/2} = 30.09(8)$ a

Reminder:	DDEP	ENSDF
	$T_{1/2}$ (a)	$30.05(8)$
		$30.08(9)$

How to proceed further?





Cs-137 Half-life: final refinement

- Critical analysis of each publication
 - Evaluate possible experimental bias
 - Evaluate uncertainty budget assessment
- Support from previous works

M.J. Woods, *The Half-life of ^{137}Cs – A critical Review*, NIM A286 (1990)

Cs-137 Half-life: final refinement

- Critical analysis of each publication
 - Evaluate possible experimental bias
 - Evaluate uncertainty budget assessment


- Support from previous works

M.J. Woods, *The Half-life of ^{137}Cs – A critical Review*, NIM A286 (1990)

- Selected publications

- Dietz *et al.*, (1973)
- Martin *et al.*, (1990)
- Schrader *et al.*, (2010)
- Bellotti *et al.*, (2012)
- Juget *et al.*, (2016),
- Unterweger *et al.*, (2020)

GEC	$T_{1/2} = 30.174 \text{ (11) a}$
CRNL	$T_{1/2} = 30.029 \text{ (12) a}$
PTB	$T_{1/2} = 30.06 \text{ (4) a}$
INFN	$T_{1/2} = 29.96 \text{ (8) a}$
IRP	$T_{1/2} = 29.994 \text{ (29) a}$
NIST	$T_{1/2} = 29.88 \text{ (15) a}$



Cs-137 Half-life: final refinement


- Critical analysis of each publication
 - Evaluate possible experimental bias
 - Evaluate uncertainty budget assessment

- Support from previous works

M.J. Woods, *The Half-life of ^{137}Cs – A critical Review*, NIM A286 (1990)

- Selected publications

Dietz et al., (1973)	GEC	$T_{1/2} = 30.174 (11) \text{ a}$
■ Martin et al., (1990)	CRNL	$T_{1/2} = 30.029 (24) \text{ a}$
■ Schrader et al., (2010)	PTB	$T_{1/2} = 30.06 (4) \text{ a}$
■ Bellotti et al., (2012)	INFN	$T_{1/2} = 29.96 (8) \text{ a}$
■ Juget et al., (2016),	IRP	$T_{1/2} = 29.994 (29) \text{ a}$
■ Unterweger et al., (2020)	NIST	$T_{1/2} = 29.88 (15) \text{ a}$



Cs-137 Half-life: new DDEP recommendation

- A final dataset of 5 measurements
 - Consistent: $\chi^2 \sim 0.8 < \chi^2_{crit} \sim 3.3$
 - Unweighted mean $T_{1/2} = 29.984 (30) \text{ a}$
 - Weighted mean $T_{1/2} = 30.018 (16)_{\text{int}} (14)_{\text{ext}} \text{ a}$
 - Recommended uncertainty extended to match *Martin et al. (1990)*

$$T_{1/2} = 30.018 (22) \text{ a}$$

	DDEP	ENSDF	This work
$T_{1/2} (\text{a})$	30.05 (8)	30.08 (9)	30.018 (22)

Cs-137 Half-life: new DDEP recommendation

- A final dataset of 5 measurements
 - Consistent: $\chi^2 \sim 0.8 < \chi^2_{crit} \sim 3.3$
 - Unweighted mean $T_{1/2} = 29.984 (30) \text{ a}$
 - Weighted mean $T_{1/2} = 30.018 (16)_{\text{int}} (14)_{\text{ext}} \text{ a}$
 - Recommended uncertainty extended to match *Martin et al.* (1990)

$$T_{1/2} = 30.018 (22) \text{ a}$$

- Outlooks
 - Lower dependency to *Martin et al.*
 - 50% of the weight
- Need for additional measurements
 - Unpublished results ?
 - PTB (continuation of Schrader works)
 - NRC (IC from 1995 2009)

	DDEP	ENSDF	This work
$T_{1/2} (\text{a})$	30.05 (8)	30.08 (9)	30.018 (22)



Decay scheme of ^{137}Cs : overview


- A simple decay scheme

- Three main transitions

- $^{137}\text{Cs} \rightarrow ^{137}\text{Ba} (11/2^-)$ ~ 95%
 - $^{137}\text{Cs} \rightarrow ^{137}\text{Ba} (3/2^+)$ ~ 5%
 - $^{137}\text{Ba} (11/2^-) \rightarrow ^{137}\text{Ba} (3/2^+)$ ~ 85+10% ($\gamma + \text{EC}$)

- Three negligible transitions

- $^{137}\text{Cs} \rightarrow ^{137}\text{Ba} (1/2^+)$ ~ $6 \cdot 10^{-4}$ %
 - $^{137}\text{Ba} (11/2^-) \rightarrow ^{137}\text{Ba} (1/2^+)$ ~ $1 \cdot 10^{-5}$ %
 - $^{137}\text{Ba} (1/2^+) \rightarrow ^{137}\text{Ba} (3/2^+)$ ~ $6 \cdot 10^{-4}$ %



Decay scheme of ^{137}Cs : overview

- A simple decay scheme

- Three main transitions


- | | |
|--|-----------------------------------|
| <ul style="list-style-type: none"> ■ $^{137}\text{Cs} \rightarrow ^{137}\text{Ba (11/2^-)}$ ■ $^{137}\text{Cs} \rightarrow ^{137}\text{Ba (3/2^+)}$ ■ $^{137}\text{Ba (11/2^-)} \rightarrow ^{137}\text{Ba (3/2^+)}$ | ~ 95% |
| <ul style="list-style-type: none"> ■ $^{137}\text{Cs} \rightarrow ^{137}\text{Ba (3/2^+)}$ | ~ 5% |
| <ul style="list-style-type: none"> ■ $^{137}\text{Ba (11/2^-)} \rightarrow ^{137}\text{Ba (3/2^+)}$ | ~ 85+10% ($\gamma + \text{EC}$) |

- Three negligible transitions

- | | |
|---|-----------------------|
| <ul style="list-style-type: none"> ■ $^{137}\text{Cs} \rightarrow ^{137}\text{Ba (1/2^+)}$ ■ $^{137}\text{Ba (11/2^-)} \rightarrow ^{137}\text{Ba (1/2^+)}$ ■ $^{137}\text{Ba (1/2^+)} \rightarrow ^{137}\text{Ba (3/2^+)}$ | ~ $6 \cdot 10^{-4}$ % |
| <ul style="list-style-type: none"> ■ $^{137}\text{Ba (11/2^-)} \rightarrow ^{137}\text{Ba (1/2^+)}$ | ~ $1 \cdot 10^{-5}$ % |
| <ul style="list-style-type: none"> ■ $^{137}\text{Ba (1/2^+)} \rightarrow ^{137}\text{Ba (3/2^+)}$ | ~ $6 \cdot 10^{-4}$ % |


- Knowledge of one of the main transitions is enough

- Completeness of the decay scheme
 - Assuming knowledge of the α_T conversion coefficient (Brlcc)



Intensity of the 662 keV gamma emission


- A limited dataset
 - Only six publication (from 1965 to 1983)
 - Three remains after applying DDEP criteria
 - *Legrand et al.* (1973) CEA $I_{\gamma} = 85.3 (3) \%$
 - *Merritt et al.* (1978) CRNL $I_{\gamma} = 84.7 (7) \%$
 - *Behrens et al.* (1983) NPL $I_{\gamma} = 85.21 (7) \%$



Intensity of the 662 keV gamma emission


- A limited dataset
 - Only six publication (from 1965 to 1983)
 - Three remains after applying DDEP criteria
 - Legrand *et al.* (1973) CEA $I_{\gamma} = 85.3 (3) \%$
 - Merritt *et al.* (1978) CRNL $I_{\gamma} = 84.7 (7) \%$
 - Behrens *et al.* (1983) NPL ~~$I_{\gamma} = 85.21 (7) \%$~~

- Consistent dataset but an issue with *Behrens et al.*
 - Estimated using magnetic spectrometer
 - Usage of discrepant α_T conversion coefficient
 - Replaced by Goodier *et al.* (1975) $I_{\gamma} = 86 (1) \%$



Intensity of the 662 keV gamma emission


- A limited dataset
 - Only six publication (from 1965 to 1983)
 - Three remains after applying DDEP criteria
 - Legrand et al. (1973) CEA $I_\gamma = 85.3(3)\%$
 - Merritt et al. (1978) CRNL $I_\gamma = 84.7(7)\%$
 - Behrens et al. (1983) NPL ~~$I_\gamma = 85.21(7)\%$~~
- Consistent dataset but an issue with *Behrens et al.*
 - Estimated using magnetic spectrometer
 - Usage of discrepant α_T conversion coefficient
 - Replaced by Goodier et al. (1975) $I_\gamma = 86(1)\%$
- Weighted mean $I_\gamma = 85.26(30)\%$
 - Weight of Legrand et al. 78 %



Intensity of the beta to ground state transition


- Another limited dataset

- Only six publications (from 1957 to 1983)
- Four of them are discarded after analysis
- Only two measurements available
 - Hansen et al. (1969) CBNM $I_\beta = 5.4 (3) \%$
 - Behrens et al. (1983) NPL $I_\beta = 5.57 (7) \%$



Intensity of the beta to ground state transition


- Another limited dataset
 - Only six publications (from 1957 to 1983)
 - Four of them are discarded after analysis
 - Only two measurements available
 - *Hansen et al.* (1969) CBNM $I_\beta = 5.4 (3) \%$
 - *Behrens et al.* (1983) NPL $I_\beta = 5.57 (7) \%$
 - Question regarding the measurement of *Behrens et al.*
 - Inaccurate measurement of α_T conversion coefficient
 - Limitation of weight of *Behrens et al.* to 50 %
 - *Behrens et al.* (1983) NPL $I_\beta = 5.57 (30) \%$
 - Average $I_\beta = 5.48 (30) \%$



Decay scheme reconstruction

- Two main possibilities
 - From 662 keV gamma intensity
 - $I_\gamma = 85.26 \text{ (30) } \%$
 - From the beta decay to the ground state of ^{137}Ba
 - $I_\beta = 5.48 \text{ (30) } \%$ corresponds to $I_\gamma = 84.97 \text{ (29) } \%$
 - Final recommendation: weighted average of both

	DDEP	ENSDF	This work
$I_\beta(11/2^-) \text{ (%)}$	94.36 (28)	94.7 (2)	94.65 (30)
$I_\beta(3/2^+) \text{ (%)}$	5.64 (28)	5.3 (2)	5.35 (30)
$I_\gamma \text{ (%)}$	84.99 (20)	85.1 (2)	85.09 (29)




Conclusions & outlooks

$T_{1/2} = 30.018 \text{ (22) a}$

- A new evaluation of ^{137}Cs has been performed
- Recommendations are consistent with previous evaluations

	DDEP	ENSDF	This work
$T_{1/2} \text{ (a)}$	30.05 (8)	30.08 (9)	30.018 (22)
$I_\beta(11/2^-) \text{ (%)}$	94.36 (28)	94.7 (2)	94.65 (30)
$I_\beta(3/2^+) \text{ (%)}$	5.64 (28)	5.3 (2)	5.35 (30)
$I_\gamma \text{ (%)}$	84.99 (20)	84.99 (20)	85.09 (29)






Conclusions & outlooks

$$T_{1/2} = 30.018 \text{ (22) a}$$

- A new evaluation of ^{137}Cs has been performed
- Recommendations are consistent with previous evaluations

	DDEP	ENSDF	This work
$T_{1/2}$ (a)	30.05 (8)	30.08 (9)	30.018 (22)
$I_\beta(11/2^-)$ (%)	94.36 (28)	94.7 (2)	94.65 (30)
$I_\beta(3/2^+)$ (%)	5.64 (28)	5.3 (2)	5.35 (30)
I_γ (%)	84.99 (20)	84.99 (20)	85.09 (29)



Analysis of datasets has shown the need for additional measurements for both half-life and transition intensities



Acknowledgment

- DDEP collaboration
 - M.A. Kellett & X. Mugeot
 - My very patient reviewer: A. L. Nichols
- Previous evaluations
 - M.J. Woods (1990)
 - R.G. Helmer and V.P. Chechev (2006)
- *How Cesium-137 was discovered by an undergraduate student ?*
(Dennis D. Patton, The journal of nuclear Medicine 40 1999)