



HAL
open science

Analytical understanding of the effect of shaping on confinement in TEM regime References

X Garbet, P Donnel, Y Melka, Y Sarazin, V Grandgirard, K Obrejan, E Bourne, G Dif-Pradalier, Y Munsch

► **To cite this version:**

X Garbet, P Donnel, Y Melka, Y Sarazin, V Grandgirard, et al.. Analytical understanding of the effect of shaping on confinement in TEM regime References. FEC 2023 - 29th IAEA Fusion Energy Conference, Oct 2023, Londres, United Kingdom. . cea-04453448

HAL Id: cea-04453448

<https://cea.hal.science/cea-04453448>

Submitted on 12 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Analytical understanding of the effect of shaping on confinement in TEM regime

X, Garbet^{1,2}, P. Donnel¹, Y. Melka¹, Y. Sarazin¹, V. Grandgirard¹, K. Obrejan¹, E. Bourne³, G. Dif-Pradalier¹, Y. Munsch¹

¹ CEA, IRFM, Saint-Paul-lez-Durance, F-13108

² SPMS, NTU, 677371 Singapore

³ SCITAS, EPFL, CH-1015 Lausanne, Switzerland

Email address: xavier.garbet@cea.fr


Abstract

- Analytical dispersion relation for Trapped Electron Mode (TEM) instab. → solved numerically
- Retains shaping effects ⇒ evaluation of their impact on growth rate
- Beneficial role of negative triangularity results from finite orbit width effect and/or ballooning of the mode

Context

- Negative triangularity $\delta < 0 \Rightarrow$ lower turbulence \Rightarrow better confinement [1,2,3]
- Recovered in GyroKinetic simulations provided enough physical ingredients are included [4,5]
- Beneficial role of $\kappa < 0$ depends particularly on aspect ratio [6] & magnetic shearing [7]
- Extensive parametric exploration remains to be done – numerically expensive with GK codes
- TEM modes = key players to explain the difference between positive/negative κ
- This work: **derivation of an analytical prediction of the TEM growth rate.**
Inputs = magnetic topology (aspect ratio, elongation, triangularity, safety factor, magnetic shearing) density & temperature gradients.

Analytical model



Assumptions

- Adiabatic ions
- Collisionless
- Trapped electrons only (no passing)

Dispersion relation

$$\mathcal{L}(\omega) = 1 - f_t \left\langle \frac{\left(\frac{\omega_{dT}}{\omega_{dT}} \left(\mathcal{E} - \frac{3}{2} + \frac{1}{\eta_s} \right) - \omega \right)}{\left(\frac{\omega_{dT}}{\omega_{dT}} \bar{\Omega}_d(\lambda) \mathcal{E} - \omega - i0^+ \right)} \right\rangle_{\lambda, \mathcal{E}} = 0 \quad \eta = d \ln T_{eq} / d \ln N_{eq}$$

Average over energy $\mathcal{E} = \frac{E}{T}$

$$\langle \dots \rangle_{\mathcal{E}} = \frac{2}{\sqrt{\pi}} \int_0^{+\infty} d\mathcal{E} \mathcal{E}^{1/2} e^{-\mathcal{E}} \dots$$

Average over pitch-angle $\lambda = \mu B_{\min} / E$

$$\langle \dots \rangle_{\lambda} = \frac{1}{f_t} \int_{B_{\min}}^1 \frac{d\lambda}{\sqrt{2\bar{\Omega}_b(\lambda)}} \dots$$

Thermal magnetic drift frequency $\omega_{dT} = n \Omega_{dT} = \frac{nq}{r_s e_a B_{\Delta} R_{\Delta}} T$

Normalized bounce frequency $\bar{\Omega}_b(\lambda)$

Thermal diamagnetic frequency $\omega_{*T} = n \Omega_{*T}$ with $\Omega_{*T} = -\frac{T}{e_a} \frac{d \ln T}{d\psi}$

Trapped fraction: $f_t = \int_{\lambda_-}^{\lambda_+} \frac{d\lambda}{\sqrt{2\bar{\Omega}_b(\lambda)}}$

Shaping modifies bounce & precession frequencies

Considered magnetic equilibrium = "Culham" [10] → parametrized shaping (ϵ, κ, δ)
Expression bounce & precession frequencies following refs. [8,9]


$$\Omega_d = \frac{1}{e_a} \frac{\partial \mathcal{J}}{\partial H} \Big|_{\mu, \psi} = \Omega_{dT} \bar{\Omega}_d(\lambda) \mathcal{E} \quad \frac{1}{\Omega_b} = \frac{\partial \mathcal{J}}{\partial H} \Big|_{\mu, \psi} \quad \Omega_b = \bar{\Omega}_b(\lambda) \sqrt{\mathcal{E}} \frac{v_T}{2q_s R_{\max}}$$

$$\frac{\partial \mathcal{J}}{\partial \psi} \Big|_{\mu, H} = \frac{q_s}{r_s} \frac{1}{B_s R_s} \frac{1}{\pi} \int_{-\theta_b}^{\theta_b} d\theta \sqrt{g_s} \frac{B_s}{v_{\parallel}} (m_a v_{\parallel}^2 + \mu B_s) \left(\frac{r_s^2}{a_s^2} \frac{1}{r_s} \frac{\partial Z_s}{\partial \theta} - (s\theta - \alpha_m \bar{D}) \frac{1}{r_s} \frac{\partial R_s}{\partial \theta} \right)$$

Functions of (ϵ, κ, δ)

$$- \frac{\alpha_m q_s}{2q_s^2} \frac{1}{r_s} \frac{1}{B_s R_s} \frac{1}{\pi} \int_{-\theta_b}^{\theta_b} d\theta \sqrt{g} \frac{B_s}{v_{\parallel}} \mu B_s \quad \text{MHD pressure parameter} \quad \alpha_m = -2\mu_0 q_s r_s R_s \frac{1}{B_s} \frac{dp_s}{d\psi}$$

$$\frac{\partial \mathcal{J}}{\partial H} \Big|_{\mu, \psi} = \frac{1}{\pi} \int_{-\theta_b}^{\theta_b} d\theta \sqrt{g_s} \frac{B_s}{v_{\parallel}} \quad s=2, \alpha_m=0, \epsilon=0.3$$



Guideline for analysis: In the fluid limit (i.e. far from threshold): $\gamma \sim \sqrt{f_t \omega_{dT} \omega_{*T} \langle \bar{\Omega}_d \rangle_{\lambda}}$

Numerical results


Necessity to include finite orbit width and/or ballooning of the mode

Finite Orbit Width (FOW) ⇒ modified definition of pitch-angle average
⇒ new definition of trapped fraction, accordingly

$$\langle \dots \rangle_{\lambda} = \frac{1}{f_t} \int_{\lambda_-}^{\lambda_+} \frac{d\lambda}{\sqrt{2\bar{\Omega}_b(\lambda)}} J_0^2(C\theta_b) \dots \quad f_t = \int_{\lambda_-}^{\lambda_+} \frac{d\lambda}{\sqrt{2\bar{\Omega}_b(\lambda)}} J_0^2(C\theta_b(\lambda))$$

FOW ⇒ more weight on **deeply trapped particles** that are **stabilized by negative triangularity**
Consequence: **Negative triangularity becomes stabilizing when including FOW effects**


C	ϵ	κ	q	s	η	α_m
scan	0,3	1,75	2,0	2,0	1000	0,0



Similar effect expected from **mode ballooning** (puts more weight on stabilized deeply trapped particles)


Large aspect ratio is stabilizing due to a reduction of trapped fraction

C	ϵ	κ	q	s	η	α_m
1,0	scan	1,75	2,0	2,0	1000	0,0



Magnetic shear is stabilizing as expected

C	ϵ	κ	q	s	η	α_m
1,0	0,3	1,75	2,0	scan	1000	0,0



Complex pattern when mixing magnetic shear & triangularity observed in GENE [7] not recovered
Possible reasons: different input parameters and/or missing physics

Conclusion & Perspectives

- Analytical dispersion relation for TEM instability
→ enables fast estimate of growth rate as function of shaping parameters
- Stabilizing effect of negative triangularity recovered **ONLY IF** a physical mechanism puts more weight on deeply trapped electrons → FOW and/or ballooning
- More systematic parameter scan under way (PhD starting end of 2023)
- Quantitative comparison with GK simulations → upgrade of model (collisions, ions) if necessary
- If successful comparison model-simulations → derivation of quasi-linear estimate of transport

References

- Y. Camenen *et al.*, "Impact of plasma triangularity and collisionality on electron heat transport in tcv l-mode plasmas". Nuclear Fusion (2007)
- M. E. Austin *et al.*, "Achievement of reactor-relevant performance in negative triangularity shape in the diti-d tokamak". Phys. Rev. Lett. (2019).
- A. Marinoni *et al.*, "A brief history of negative triangularity tokamak plasmas". Reviews of modern plasma physics (2021)
- G. Merlo *et al.*, "Investigating profile stiffness and critical gradients in shaped tcv discharges using local gyrokinetic simulations of turbulent transport". Plasma Physics and Controlled Fusion (2015).
- G. Di Giannatale *et al.*, "Triangularity effects on global flux-driven gyrokinetic simulations", Journal of Physics: Conference Series (2022)
- A. Balestri *et al.*, "The aspect ratio dependence on confinement enhancement in negative triangularity plasmas", EPS 2023
- Gabriele Merlo *et al.*, "Interplay between magnetic shear and triangularity in ion temperature gradient and trapped electron mode dominated plasmas". Journal of Plasma Physics (2023).
- C. C. Hegna, "The effect of three-dimensional fields on bounce averaged particle drifts in a tokamak". Physics of Plasmas (2015).
- Zhangsheng Huang *et al.*, "The synergistic effects of three dimensional magnetic perturbations and finite beta on collisionless trapped electron mode in tokamak plasmas". Nuclear Fusion (2022).
- J. W. Connor *et al.*, Report No. CLM-M106. Culham Laboratory Report (1985)