



Time-dependent integrated modelling overview and DEMO aspects

C Bourdelle

► To cite this version:

C Bourdelle. Time-dependent integrated modelling overview and DEMO aspects. EFPW31 - 31st European Fusion Programme Workshop, Jan 2024, Prague, Czech Republic. cea-04465645


HAL Id: cea-04465645

<https://cea.hal.science/cea-04465645>

Submitted on 19 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Time-dependent integrated modelling Overview and DEMO aspects

C. Bourdelle

With input from: C. Angioni, N. Bonanomi, F. Casson, E. Fable, J. Garcia, L. Garzotti, M. Groth, F. Koechl, P. Mantica, P. Manas, P. Maget, M. Siccinio, P. Strand, ...



outline

- W physics needs and integrated modelling framework availability
- Validation of dynamical integrated modelling up to the separatrix
- Validation including pedestal and SOL, progresses
- Large scale validation perspectives
- DEMO predictions status, gaps and perspectives

Physics embedded in integrated modelling platform for time evolution of W and its radiation

[C. Angioni] to model W transport/radiation need:

- **neoclassical W transport:**
 - Proper background plasma profile prediction thanks to turbulence models for heat/**particle**/momentum
 - **asymmetry effects** due to T anisotropy and rotation
- **turbulent W transport,** especially if low neoclassical transport

This physics is in our integrated modelling platforms, including the self-consistent radiation feedback

	ASTRA	HFPS/JETTO	RAPTOR	ETS*
Background profiles up to ped top	TGLFsat2, QuaLiKiz	TGLFsat2, QuaLiKiz	QLKNN	TGLFsat2, QuaLiKiz
Ped profiles	IMEP (ad-hoc heat ped transport + MISHKA)	Europed, MISHKA stability, scalings	none	PENN
W neoclassical transport	NCLASS For asymmetry: NEO, FACIT	NCLASS For asymmetry: NEO, FACIT	none	NCLASS For asymmetry: NEO
W turbulent transport	TGLFsat2, QuaLiKiz	TGLFsat2, QuaLiKiz	none	TGLFsat2 EDWM
W radiation	STRAHL	SANCO	ADAS	AMNS based

* ETS not used yet with W transport/rad.



outline

- W physics needs and integrated modelling framework availability
- **Validation of dynamical integrated modelling up to the separatrix**
- Validation including pedestal and SOL, progresses
- Large scale validation perspectives
- DEMO predictions status, gaps and perspectives

Time-dependent flux driven integrated modelling over multiple confinement times: highly nonlinear coupling

- Level 1: j, T_i, T_e
 - Forgiving: transport driven by temp. gradients and is “stiff”
 - Always predict T_i and T_e profiles otherwise turbulence amplitude wrong
- Level 2: j, T_i, T_e, n_e
 - Non-trivial: particle transport not stiff (off-diagonal transport)
 - Depends sensitively on turb. spectra, collisions, kinetic resonances
- Level 3: $j, T_i, T_e, n_e, V_{tor}$
 - Challenging: momentum transport from symmetry breaking
 - Feedback potential for barrier formation (ExB shear)
- Level 4 : j, T_i, T_e, n_e, n_i multi-ion (isotopes, impurities), V_{tor}
 - Exciting territory, complex non-linear interplays
 - Heavy Impurity transport needs all L3 channels (sets neoclassical transport and poloidal asymmetries), and provides radiation feedback
- Level ++ : j, T_i, T_e, n_e, n_i multi-ion (isotopes, impurities), V_{tor} in burning plasmas
 - where $P_{alpha} > P_{aux}$ ($Q > 5$), background profiles impact P_{alpha} accounting for EP-MHD, Alpha redistribution and impact EP/ β on turbulent transport





Level 1. WEST L mode radiative collapse due to large density → LHCD off axis → T_i peaking reduction → increasing W inward neoclassical flux

RAPTOR-QLKNN-ADAS

T_e, T_i, j Predictive up to separatrix

Mixed with stand-alone W profile evolution
checked against NEO+QuaLiKiz predictions using
measured background profiles








Need both W peaking and decrease of the central LHCD absorption to capture the radiative collapse

[Ostuni NF2022]






Level 4. W-accumulation at JET due to NBI central particle source → density peaking → W neoclassical accumulation enhanced by rotation

Prediction, 1.5 s of plasma evolution ($5x\tau_E$).




State-of-the-art core plasma integrated modelling evolving j , T_e , T_i , n_D , n_{Be} , n_{Ni} , n_W , ω , rad., NBI
JETTO-SANCO-QuaLiKiz-NEO-PENCIL-FRANTIC-ad-hocETB successfully described **core W-accumulation due to NBI particle source** [Breton NF 2018]




Level 4. W-accumulation avoidance at JET: ICRH → enhanced outward turbulent particle transport → flater n_i core profile → reduced W neoclassical inward transport → delayed radiative collapse

Prediction: 1.5 s of plasma integrated modelling evolving j , T_e , T_i , n_D , n_{Be} , n_{Ni} , n_W , ω , rad., NBI, ICRH

JETTO-SANCO-QuaLiKiz-NEO-PENCIL-FRANTIC-ad-hocETB-PION




[Casson NF 2020]



Level 4 (w/o V_{tor}). W-accumulation avoidance at AUG: ECRH → enhanced outward turbulent particle transport → flater n_i core profile → reduced W neoclassical inward transport

AUG #32408






Prediction: plasma integrated modelling evolving j , T_e , T_i , n_D , n_B , n_W , rad., NBI.

Rot. CX, ECRH stand-alone ASTRA-QuaLiKiz-NEO-NBImodel-STRahl

As on JET, NBI particle source enhances W inward neoclassical convection

As ICRH on JET, ECRH on → T_e/T_i increase → turbulent particle diffusion → reducing W inward neoclassical flux





outline




- W physics needs and integrated modelling framework availability
- Validation of dynamical integrated modelling up to the separatrix
- **Validation including pedestal and SOL, progresses**
- Large scale validation perspectives
- DEMO predictions status, gaps and perspectives

Merging the state-of-the-art level 3-4 core modelling (NEO or FACIT with QuaLiKiz or TGLF) with EDGE2D-ERO W state-of-the-art

H. Kumpulainen EPS 2023, sub PPCF 2024

Vs previous e.g. COREDIV work
[Telesca JNM 2013] here **W neoclassical (NEO) + turbulent (QuaLiKiz) transport in core and pedestal**

NB: Background pedestal density and temperature impacting W penetration [cf Angioni's talk]: unkown dynamical particle source and heat+particle transport btw ELMs





outline

- W physics needs and integrated modelling framework availability
- Validation of dynamical integrated modelling up to the separatrix
- Validation including pedestal and SOL, progresses
- **Large scale validation perspectives**
- DEMO predictions status, gaps and perspectives

Large scale validation on all W EU tokamak possible now at large scale

Predictive int. modelling incl. W and radiation successfully used to prepare JET DTE2 [[Garcia NF2023](#)]

[TSVV11](#) made available the High Fidelity Pulse Simulator

Python-driven workflow based on **IMASified JINTRAC**

- NEO and FACIT available
- QuaLiKiz and TGLF available
- SANCO for self-consistent radiation
- EDGE2D

Coupled to experimental IMAS data from AUG, JET, TCV, WEST, **on the EUROfusion Gateway**

Adding IMEP could be done
More physics driven devt as needed ...

Large scale automated simulation launching tools operational (synergy project TSVV11-NL eScience center)
<https://github.com/duqtools/duqtools> (open-source)]


List of IMAS data

```
g2aho/aug/36982/2
g2aho/jet/75225/2
g2aho/jet/90350/2
g2aho/jet/92432/2
g2aho/jet/94875/1
g2aho/tcv/64958/2
g2aho/west/54568/1
g2aho/west/54728/1
g2aho/west/55181/1
g2aho/west/55525/1
...
```

Canonical UQ template

```
create:
  runs_dir: ./duqduq/{{ run.name }}
  template: ./path/to/template/
  template_data:
    user: {{ handle.user }}
    db: {{ handle.db }}
    shot: {{ handle.shot }}
    run: {{ handle.run }}
    sampler:
      ...
    dimensions:
      ...
  system: jetto-v220922
```

duqduq





outline

- W physics needs and integrated modelling framework availability
- Validation of dynamical integrated modelling up to the separatrix
- Validation including pedestal and SOL, progresses
- Large scale validation perspectives
- **DEMO predictions status, gaps and perspectives**



Level 4 (w/o V_{tor}). DEMO W interplay with plasma up to sep., impact of W content → radiation → P_{fus}

DEMO baseline 2019/2021 5.74 T 19.5 MA R=9m
A=3.1 5°MW ECRH


Prediction: 70 s (τ_E 4-6s) of integrated prediction j ,
 T_e , T_i , n_D , n_T , n_{He} , n_{Ar} , n_{Xe} , n_W , rad., ECRH,
continuous pellet model, fusion power self-consistently calculated and iterated

HFPS-NCLASS-TGLFsat2-SANCO

Pedestal: ITER-EPED scaling $P_{\text{ped}} = 0.54 B^{0.84} I_p \cdot n_{\text{ped}}$ such that pellet fueling feedback $n_{\text{ped}}/n_{Gr} = 95\%$ $n_{Gr} \propto I_p/a^2$

Sepatrix: ITER SOLPS scaling for n_{sep} and T_{sep}


W total content scanned




[Koechl Dec 2022 DCT]

GAP 1/2: Improving the core-pedestal-SOL predictive modeling towards opaque/pellet fueled plasmas

AUG, experiment
[Kallenbach PPCF2018]



ITER SOLPS
modelling
[Pitts NME2009]



- Shorter term (2024-2025 DEMO Central Team, PSD, transport): **Explore more intensively impact of uncertainties on separatrix/pedestal parameters on DEMO scenarios incl. core W behavior and pellet interplay**, using both HFPS and ASTRA-IMEP with SOLPS-ERO-DEMO informed boundary conditions.
- Longer term: **1st principle SOL + pedestal incl. W, transport and erosion**
 - **validation** on gas fueled present day experiments: Gyrokinetic or fluid turbulence + neutral models+ W, including SOL and confined plasmas: e.g. XGC1 on WEST [Dominski APS 2023]
 - **Predictions ITER/DEMO opaque/pellet fueled** challenging numerically, for now remains sensitive to initial conditions



Gap 2/2: Approaching Burning plasmas with a hierarchy of models Workplan in DEMO Central Team, transport aspects 2024-2025

- **Standard integrated modelling HFPS extended to Alpha impact on MHD stability hence on Alpha redistribution** (ATEP Ph. Lauber et al) And iterate, expect P_{fus} oscillations in time. could add W turbulent and neoclassical transport interplay > 2024
- **Flux driven Alpha heated global gyrokinetic modelling** of ITER Q=10 and DEMO core $\rho>0.6$ using GENE-Tango GPU accelerated – at limit of Today's computing capacity (A. Di Siena et al). ITER Q=10 1st modelled burning plasma on-going. could add W turbulent and neoclassical transport interplay > 2025
- **Theoretical understanding and Global gyrokinetic mapping** GENE/ORB5, with varied fraction of fast ions and $T_{fast}/T_{thermal}=10$ and 100 (3.5 MeV Alpha)

Effect of fast ions on turbulent transport, hence T_i , hence P_{fus}	HFPS/AST RA-TGLFsat2	GENE - Tango
Thermal ion dilution	yes	
Shafranov shift stabilization by increase of α_{MHD} by suprathermal pressure	Yes	
Electrostatic stabilization by resonant fast ions	Yes	yes
EM-stabilization (enhanced by fast-ions)	Yes	
Increased nonlinear stabilization	No	

More in J. Citrin's talk



Conclusion

- Validation up to pedestal top in H mode and up to separatrix in L mode
 - **Physics based Integrated modelling j, Ti, Te, multi-ion (isotopes, impurities), Vtor HFPS-JETTO/ASTRA allowed to understand JET, AUG and WEST radiative collapses.**
In AUG and JET, key deleterious role of density peaking due to NBI ‘erased’ by turbulent particle transport enhancement by electron heating ECRH or ICRH. On WEST, central LHCD needed on-axis to maintain temperature screening
 - Core integrated modelling set up was successfully used to predict JET DTE2 [[Garcia NF2023](#)]
 - HFPS connected to all EU tokamak IDS ready on Gateway with large scale UQ tools
- **Validation up to W source started using high fidelity neoclassical transport (NEO [[Kumpulainen EPS 2023, soon PPCF2024](#)]) but:**
 - Pedestal turbulent transport remains ad hoc for heat and particle
 - Neutral source model validation in pedestal difficult, in int. modelling feedback on line averaged density with ad-hoc neutral energy and poloidally sym. [e. g. [Luda NF2023](#)]
 - SOL turb. codes including all species, heat, radiation self-consistently just starting, for ex. XGC1 on WEST [[Dominski APS 2023](#)]



perspectives

- **Validation up to pedestal top in H mode and up to separatrix in L mode**
 - HFPS machine agnostic, operational with IDS data on the Gateway together with large scale validation tools [@TSVV11]. **Larger scale validation, not cherry picking, now possible**
 - requires data access (FAIR4fusion, JET data center status?),
 - soft and hardware support for Long Term Storage Facility and simulation database.
Incl. Validation against high β JET T/D/DT and JT60-SA N-NBI
 - Need strategy towards physics based pedestal + SOL transport and part. source TSVV11 + other TSVVs/ENRs. Need more well diagnosed core-ped-SOL including neutrals pulses ‘à la [[Horvarth PPF2023](#)]’ at JET, more to be expected at DIII-D Lyman-alpha diags
- **Physics based extrapolation towards opaque, pellet fueled devices + burning plasmas** remains challenging but 2024-2025 more intensive exploration of uncertain SOL/pedestal in HFPS-TGLFsat2-NEO-SANCO, including interplay with EP MHD and global GENE-TANGO modelling proceeding [TSVV10-TSVV11, DEMO CT / PSD]