



HAL
open science

Draft genome sequence of propane- and butane-oxidizing actinobacterium *Rhodococcus ruber* IEGM 231

Irena B Ivshina, Maria S Kuyukina, Anastasiya V Krivoruchko, Valérie Barbe, Cécile Fischer

► To cite this version:

Irena B Ivshina, Maria S Kuyukina, Anastasiya V Krivoruchko, Valérie Barbe, Cécile Fischer. Draft genome sequence of propane- and butane-oxidizing actinobacterium *Rhodococcus ruber* IEGM 231. *Genome Announcements*, 2014, 2 (6), pp.e01297-14. 10.1128/genomea.01297-14 . cea-04643767

HAL Id: cea-04643767

<https://cea.hal.science/cea-04643767>

Submitted on 10 Jul 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Draft Genome Sequence of Propane- and Butane-Oxidizing Actinobacterium *Rhodococcus ruber* IEGM 231

Irena B. Ivshina,^{a,b} Maria S. Kuyukina,^{a,b} Anastasiya V. Krivoruchko,^{a,b} Valérie Barbe,^c Cécile Fischer^d

Institute of Ecology and Genetics of Microorganisms, Ural Branch, Russian Academy of Sciences, Perm, Russia^a; Perm State University, Perm, Russia^b; CEA/DSV/IG/Genoscope/LF, Évry, France^c; Université Val d'Essonne-UMR8030 and CEA/DSV/IG/Laboratoire de Génomique et Biochimie du Métabolisme and CNRS, Évry, France^d

We report a draft genome sequence of *Rhodococcus ruber* IEGM 231, isolated from a water spring near an oil-extracting enterprise (Perm region, Russian Federation). This sequence provides important insights into the genetic mechanisms of propane and *n*-butane metabolism, organic sulfide and beta-sitosterol biotransformation, glycolipid biosurfactant production, and heavy metal resistance in actinobacteria.

Received 1 November 2014 Accepted 7 November 2014 Published 11 December 2014

Citation Ivshina IB, Kuyukina MS, Krivoruchko AV, Barbe V, Fischer C. 2014. Draft genome sequence of propane- and butane-oxidizing actinobacterium *Rhodococcus ruber* IEGM 231. *Genome Announc.* 2(6):e01297-14. doi:10.1128/genomeA.01297-14.

Copyright © 2014 Ivshina et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Anastasiya V. Krivoruchko, nast@iegm.ru.

Nocardioform actinobacteria of the species *Rhodococcus ruber* grow on the gaseous hydrocarbons ethane, propane, and *n*-butane as sole carbon and energy sources and can be used in prospecting for gas and oil deposits, as well as in other technologies (1–5). The *R. ruber* strain IEGM 231 was isolated from a water spring near an oil-extracting enterprise (Perm region, Russian Federation) and deposited in the IEGM Collection of Alkanotrophic Microorganisms (acronym IEGM, no. 768 WDCM, http://www.iegm.ru/iegmcol/strains/rhodoc/ruber/r_r_u_b_e_r_231.html) (6). The strain was shown in our laboratory to utilize propane, *n*-butane, and liquid alkanes (C₁₂–C₁₇); convert aryl alkyl sulfides into optically pure sulfoxides; transform beta-sitosterol into therapeutically prospective stigmast-4-en-3-one; produce nontoxic glycolipid biosurfactants; have a high resistance to heavy metals; and accumulate molybdenum and nickel.

The whole genome of *Rhodococcus ruber* IEGM 231 was sequenced using Illumina technology. A mate-paired library with an insert size of 6 kb was produced and sequenced with the MiSeq system (2 × 250 nt). An approximately 130-fold coverage was generated, and the data were assembled with the Velvet assembler (<https://www.ebi.ac.uk/~zerbino/velvet>). Gap filling was performed using GapCloser (<http://soap.genomics.org.cn/soapdenovo.html>) on scaffolds more than 2 kb in size. The annotation of coding sequences (CDS) and prediction of gene functions were performed using the MicroScope platform (<https://www.genoscope.cns.fr/agc/microscope/home/index.php>) (7), which integrated 115 contigs, resulting in 46 scaffolds from the assembly. The genome size of *R. ruber* IEGM 231 is estimated to be 6.01 Mb with a G+C content of 70.22%.

A total of 5,928 CDSs with an average length of 948 bp, 6 rRNAs, and 53 tRNAs were found in the *R. ruber* IEGM 231 genome. At least 73 CDSs coded for monooxygenases/hydroxylases, 22 CDSs coded for cytochromes P450, 45 CDSs coded for dioxygenases, 13 CDSs coded for peroxidases, and 285 CDSs coded for dehydrogenases. Among monooxygenases, genes coding for 2 pu-

tative propane monooxygenases *mmoABC* and *prmA*, 2 alkane 1-monooxygenases *alkB*, 9 flavin monooxygenases, and 1 cyclohexanone 1,2-monooxygenase were revealed, reinforcing the strong phenotypic abilities of the species toward hydrocarbons. Diversity of these sequences could also be evidence of the existence of several alkane and organic sulfide degradation systems in *R. ruber* IEGM 231 (8, 9). Two *choD* (coding for cholesterol oxidase) homologues and 1 gene homologous to 3-beta hydroxysteroid dehydrogenase/isomerase were present. They could account for steroid compound transformations, as already shown for other *R. ruber* strains (10, 11). The diversity of CDSs coding for glycolipid biosurfactant synthesis was presented by 1 malonyl CoA-ACP transacylase, 14 acyl-CoA synthetases, 1 fatty acid synthase I, 9 3-oxoacyl-ACP reductases, 1 cyclopropane mycolic acid synthase, 1 polyketide synthase, 1 meromycolate extension ACP, 3 mycolyltransferases, 1 maltooligosyl trehalose synthase, 1 maltooligosyl trehalose trehalohydrolase, and 1 trehalose synthase. Among heavy metal-associated CDSs, there were CDSs coding for heavy metal resistance proteins (5 CDSs), heavy metal transporters (8 CDSs), cation efflux enzymes (3 CDSs), metal binding proteins (3 CDSs), a mercuric reductase, and an alkylmercuryl lyase.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under accession numbers [CCSD01000001](https://www.ncbi.nlm.nih.gov/nuccore/CCSD01000001) to [CCSD01000115](https://www.ncbi.nlm.nih.gov/nuccore/CCSD01000115).

ACKNOWLEDGMENTS

We are grateful to the LABGeM team of UMR8030 (Évry, France) and the national infrastructure “France Génomique” for curation and annotation of the genome and for technical and bioinformatics support. The work was funded by a grant from the Russian Academy of Sciences Presidium Program “Leaving Nature: Current State and Development Problems” (no. 01201256869) and by a grant from the Russian Federation Ministry of Education and Science (State Assignment 6.1194.2014/K). The exchange of researchers was made possible by a grant from the University of Évry Val d'Essonne.

REFERENCES

- Xu K, Tang Y, Ren C, Zhao K, Sun Y. 2013. Diversity and abundance of *n*-alkane-degrading bacteria in the near-surface soils of a Chinese onshore oil and gas field. *Biogeosciences* 10:2041–2048. <http://dx.doi.org/10.5194/bg-10-2041-2013>.
- Philp JC, Kuyukina MS, Ivshina IB, Dunbar SA, Christofi N, Lang S, Wray V. 2002. Alkanotrophic *Rhodococcus ruber* as a biosurfactant producer. *Appl. Microbiol. Biotechnol.* 59:318–324. <http://dx.doi.org/10.1007/s00253-002-1018-4>.
- Kuyukina MS, Ivshina IB. 2010. Application of *Rhodococcus* in bioremediation of contaminated environments, p 231–262. In Alvarez HM (ed), *Biology of Rhodococcus*. Springer, Berlin.
- Ivshina IB, Kuyukina MS, Krivoruchko AV, Plekhov OA, Naimark OB, Podorozhko EA, Lozinsky VI. 2013. Biosurfactant-enhanced immobilization of hydrocarbon-oxidizing *Rhodococcus ruber* on sawdust. *Appl. Microbiol. Biotechnol.* 97:5315–5327. <http://dx.doi.org/10.1007/s00253-013-4869-y>.
- Grishko VV, Nogovitsina YM, Ivshina IB. 2014. Bacterial transformations of terpenoids. *Russ. Chem. Rev.* 83:323–342. <http://dx.doi.org/10.1070/RC2014v083n04ABEH004396>.
- Ivshina IB, Kuyukina MS. 2013. Turning Russian specialized microbial culture collections into resource centers for biotechnology. *Trends Biotechnol.* 31:609–611. <http://dx.doi.org/10.1016/j.tibtech.2013.08.002>.
- Vallenet D, Belda E, Calteau A, Cruveiller S, Engelen S, Lajus A, Le Fèvre F, Longin C, Mornico D, Roche D, Rouy Z, Salvignol G, Scarpelli C, Thil Smith AA, Weiman M, Médigue C. 2013. Microscope—an integrated microbial resource for the curation and comparative analysis of genomic and metabolic data. *Nucleic Acids Res.* 41:D636–D647. <http://dx.doi.org/10.1093/nar/gks1194>.
- Koch DJ, Chen MM, van Beilen JB, Arnold FH. 2009. *In vivo* evolution of butane oxidation by terminal alkane hydroxylases AlkB and CYP153A6. *Appl. Environ. Microbiol.* 75:337–344. <http://dx.doi.org/10.1128/AEM.01758-08>.
- Ceccoli RD, Bianchi DA, Rial DV. 2014. Flavoprotein monooxygenases for oxidative biocatalysis: recombinant expression in microbial hosts and applications. *Front Microbiol.* 5:00025. <http://dx.doi.org/10.3389/fmicb.2014.00025>.
- Fernández de las Heras L, Perera J, Navarro Llorens JM. 2014. Cholesterol to cholestenone oxidation by ChoG, the main extracellular cholesterol oxidase of *Rhodococcus ruber* strain Chol-4. *J. Steroid Biochem. Mol. Biol.* 139:33–44. <http://dx.doi.org/10.1016/j.jsbmb.2013.10.001>.
- Fernández de las Heras L, van der Geize R, Drzyzga O, Perera J, Navarro Llorens JM. 2012. Molecular characterization of three 3-ketosteroid- Δ^1 -dehydrogenase isoenzymes of *Rhodococcus ruber* strain Chol-4. *J. Steroid Biochem. Mol. Biol.* 132:271–281. <http://dx.doi.org/10.1016/j.jsbmb.2012.06.005>.