



HAXPES spectra and quantification of bulk Mo and MoSe₂ compound

Olivier Renault, Nicolas Gauthier, Eugénie Martinez, Kateryna Artyushkova

► To cite this version:

Olivier Renault, Nicolas Gauthier, Eugénie Martinez, Kateryna Artyushkova. HAXPES spectra and quantification of bulk Mo and MoSe₂ compound. ECASIA 2024 - European association on applications of surface and interface analysis, Jun 2024, Göteborg, Sweden. , 2024. cea-04658986

HAL Id: cea-04658986

<https://cea.hal.science/cea-04658986v1>

Submitted on 22 Jul 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

HAXPES spectra and quantification of bulk Mo and MoSe₂ compound

Olivier Renault¹, Nicolas Gauthier¹, Eugénie Martinez¹, Kateryna Artyushkova²

¹CEA-Leti, Univ. Grenoble Alpes, F-38000 Grenoble, France

²Physical Electronics, 18725 Lake Dr E, Chanhassen, MN 55317, USA

olivier.renault@cea.fr

The HAXPES spectra of bulk Mo and MoSe₂ compounds obtained with Cr K α_1 excitation ($h\nu=5414.8$ eV) are presented, with special attention paid to binding energy scale correction, quantification and repeatability aspects [1]. Cr K α HAXPES allows to ionize all Mo and Se core-levels of principal quantum numbers N=2, 3 and 4.

Empirically determined sensitivity factors (RSFs) were provided by the PHI Multipak software version 9.9.0.8, from the most recently released RSF database. The RSFs were derived from the pure-element relative sensitivity factor as defined in ISO 18118:2015 [2] which were measured on pure element samples using a Cr K α source. Using the determined pure element RSF's leads to quantification error of ~5% for the MoSe₂ compound.




Fig.1- Cr K α HAXPES survey spectrum of bulk MoSe₂ [1].

Acknowledgements

This work, carried out on the Platform for Nanocharacterisation (PFNC), was supported by the "Recherche Technologique de Base" and "France 2030 - ANR-22-PEEL-0014" programs of the French National Research Agency (ANR) "

References

- [1] P.-M. Deleuze, N. Gauthier, E. Martinez, K. Artyushkova and O. Renault, T. Maindron, *Surf. Sci. Spectra* 31, 014006 (2024).
- [2] International Organization for Standardization 2015, Surface chemical analysis - Auger electron spectroscopy and X-ray photoelectron spectroscopy - Guide to the use of experimentally determined relative sensitivity factors for the quantitative analysis of homogeneous materials, ISO 18118:2015