



Non-linear gyro-kinetic Ion Temperature Gradient (ITG) and Trapped Electron Modes (TEM) turbulence modelling in X-point geometry in negative (NT) and positive (PT) triangularity shapes

M Bécoulet, G T A Huijsmans, X Garbet, P Donnel, G Dif-Pradalier, A Marinoni, P Ulbl, S Coda, Y Camenen, C Chrystal, et al.

► To cite this version:

M Bécoulet, G T A Huijsmans, X Garbet, P Donnel, G Dif-Pradalier, et al.. Non-linear gyro-kinetic Ion Temperature Gradient (ITG) and Trapped Electron Modes (TEM) turbulence modelling in X-point geometry in negative (NT) and positive (PT) triangularity shapes. 50th EPS Conference on Plasma Physics, Jul 2024, Salamanque, Spain. . cea-04726648

HAL Id: cea-04726648

<https://cea.hal.science/cea-04726648v1>

Submitted on 8 Oct 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Non-linear gyro-kinetic Ion Temperature Gradient (ITG) and Trapped Electron Modes (TEM) turbulence modelling in X-point geometry in negative (NT) and positive (PT) triangularity shapes.

M.Bécoulet¹, G.T.A.Huijsmans¹, X.Garbet^{1,2}, P. Donnel¹, G. Dif-Pradalier¹, A. Marinoni³, P. Ulbl⁴, S. Coda⁵, Y. Camenen⁶, C. Chrystal⁷, M. J. Pueschel^{8,9}

¹*CEA, IRFM, 13108 Saint-Paul-Lez-Durance, France*

²*School of Physical and Mathematical Sciences, Nanyang Technological University, 637371 Singapore*

³*Jacobs School of Engineering, University of California San Diego, CA 92093-0403, USA*

⁴*Max-Planck-Institute for Plasma Physics, 85748 Garching, Germany*

⁵*EPFL-SPC, CH-1015 Lausanne, Switzerland*

⁶*CNRS, Aix-Marseille Univ., PIIM UMR7345, Marseille, France*

⁷*General Atomics, San Diego, CA, United States of America*

⁸*Eindhoven University of Technology, 5600 MB Eindhoven, Netherlands*

⁹*Dutch Institute for Fundamental Energy Research, 5612 AJ Eindhoven, Netherlands*



Acknowledgements: This work has been carried out within the framework of the EUROfusion Consortium, funded by the European Union via the Euratom Research and Training Programme (Grant Agreement No 101052200 — EUROfusion).

The Swiss contribution to this work has been funded by the Swiss State Secretariat for Education, Research and Innovation (SERI).

This material is based upon work supported by the U.S. Department of Energy, Office of Science, Office of Fusion Energy Sciences, using the DIII-D National Fusion Facility, a DOE Office of Science user facility under Award DE-FC02-04ER54698.

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favouring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union, the European Commission or SERI. Neither the European Union nor the European Commission nor SERI can be held responsible for them.

This work was carried out using the HPCs: CEA TGCC IRENE (France) and CINECA MARCONI (Italy)

Abstract.

The comparison of the non-linear gyro-kinetic particles code JOREK-GK with gyro-kinetic codes GS2 [A Marinoni et al *Plasma Phys Control Fus* 51(2009) 055016], GENE-X [P Ulbl et al 29th IAEA FEC 2023, London, UK] and GENE [F. Jenko, W. Dorlan et al *Phys of Plasmas* 7, 1904 (2000)] was done on a few selected negative/positive (NT/PT) triangularity TCV plasma parameters. It showed rather good agreement between codes in linear growth rates of ITG/TEM single modes in linear phase and clear beneficial effect of negative triangularity on core confinement compared to positive triangularity shape.

The global non-linear gyro-kinetic modelling using JOREK-GK code of the ITG/TEM saturated turbulence for realistic negative triangularity DIII-D pulses suggest gyro-Bohm ρ^* confinement scaling similar to H-modes. That could probably explain high confinement obtained experimentally for negative triangularity shaped plasmas. This scaling would be very favorable factor for reactor size machines with high confinement operation without harmful ELMs.

A larger parameter range should be explored in the future both experimentally and numerically to confirm the gyro-Bohm scaling for NT plasmas.



Kinetic particles in JOREK-GK code. Non-linear gyro-kinetic modelling of electrostatic turbulence ITG/TEMs in realistic tokamak X-point geometry.

[JOREK as MHD code: G Huysmans, NF 2007, PPCF 2009, M Hoelzl NF2021; JOREK-GK: G Huijsmans EPS 2023, M Becoulet EPS 2023, IAEA 2023]

- Particles are initialized on the equilibrium grid, plasma profiles and magnetic field calculated by JOREK as fluid MHD code => **X-point, SOL, divertor, walls, coils geometry can be included.**
- Non-linear full-f.
- Particles are advanced in time evolving electric field and static magnetic field –electrostatic turbulence.
- Ions: parallel and ExB motion of gyro-centers in gyro-averaged electric field.
- Electrons: adiabatic (ITG only) or kinetic electrons (ITG/TEM). Heavy electrons ($m_i/m_e=100$). Guiding centers.
- Projection and solution in weak form using the same basis functions as in fluid MHD. Two types of filters are used, hyper-diffusion in the poloidal plane and a Laplacian in the parallel direction.
- HPC: MARCONI(Italy), JFRS-1(Japan),IRENE-ROME (France): typical job N=5:5:40 toroidal harmonics, 5.10^8 particles (depending on the tokamak size) on 54 nodes, 48h wall time (~0.5ms to saturation), time step $dt \sim 5.10^{-8}$ s. Good (close to ideal) scaling for particles on HPC compared to fluid non-linear MHD.

Example:



COMPASS, TEM/ITGs with RMPs



Motivation (I):

Initial studies on TCV : EC heated L-mode ($\text{Te} > \text{Ti}$)=> electron heat transport decreases at sufficiently strong negative triangularity (NT). TEMs are stabilised at NT.



[Y Camenen NF 2007]




Motivation (II): Seems attractive for reactor!

- 1) DIII-D initial NT/PT experiments. $T_e = T_i$ [M Austin PRL2019, A Marinoni NF2021]: no pedestal, no ELMs, high core confinement ~as in H-mode, but without ELMs.
- 2) 2023 dedicated NT campaign on DIII-D diverted plasmas, $T_e = T_i$, low collisionality : high core confinement . No access to 2nd stability limit in NT=> no high pressure pedestal, ELMs.



[K E Thome et al PPCF to be published 2024.]





[A Nelson PRL2023]



Comparison global JOREK-GK with local flux tube GS2 code [A Marinoni et al PPCF 2009] for TCV L- modes in TEM regime ($T_e > T_i$), similar linear growth rates, smaller at NT compared to PT.






Comparison of global JOREK-GK with GENE (w/o X-point) [F. Jenko et al PPCF 7, 1904 (2000)] in linear TEM dominant regime ($T_e > T_i$) for TCV parameters: similar linear growth rates, smaller at NT compared to PT.






Comparison of JOREK-GK with GENE-X with the same profiles and equilibrium

GENE-X (band $\psi_n > 0.57$)

JOREK-GK (all plasma+ SOL $\psi_n > 0$)



Larger avalanches at PT triangularity => Larger perpendicular heat fluxes





Flux expansion NT (short leg) $f_x=3.5$, PT (long leg) $f_x=4.3$. Divertor // heat fluxes are mainly from electrons. From Eich's fit (NF 2013) for // electron heat flux: PT $\lambda_q = 4\text{mm}$; NT $\lambda_q = 3.5\text{mm}$.

$$\lambda_q(\text{mm}) = (0.7 \pm 0.2) \cdot B_{tor}^{-0.8 \pm 0.1} \cdot q_{95}^{1.05 \pm 0.2} \cdot P_{SOL}^{0.1 \pm 0.1} \cdot R_{geo}^{0 \pm 0.1}$$

$$q(s) = q_0 \cdot \exp\left(-\frac{s - s_0}{\lambda_q f_x}\right)$$





Eich's fit (NF2013)



JOREK-GK modelling for DIII-D pulses with X-point from 2023 NT campaign



EQDSK 193778,
low q95=2.8

EQDSK 194288,
high q95=4.67



ρ^* scaling in NT?

Low q95 # 193778 B=2T, q95=2.7 - this is a case Bscale=1. Keeping the same equilibrium and q-profile constructing new $B_t \Rightarrow B_{scale} * B_t$; $I_p \Rightarrow B_{scale} * I_p$; $n_e \Rightarrow n_e * B_{scale}^{4/3}$, $T_{tot} \Rightarrow T_{tot} * B_{scale}^{2/3}$, $P_{tot} \Rightarrow P_{tot} * B_{scale}^2$. As a result $\rho^* \Rightarrow \rho^* B_{scale}^{-2/3}$. Here: Bscale=0.5, Bscale=1, Bscale=1.5. Electron-ion collisions model: small angle random scattering [Lana Rekhviashvili, Zhixin Lu et al POP 30 (2023)]






High q95=4.7 (# 194288), B=2T => initial case Bscale=1.


B=2T (Bscale=1 #194288)



B=3T (Bscale=1.5)







scaled profiles DIII-D #194288




Running JOREK-GK for three cases (B=1T, 2T, 3T) up to established saturated quasi-stationary TEM/ITG turbulence.


Filters : perp 2e-11, parallel 2, particles 5e8, Npsi=150, Nth=500, mi/me=100, harmonics N=5:5:40





Scaled low q95(#193778) pulse. Time averaged in saturation (0.38-0.58ms) heat conductivity. Confinement improves with decreasing ρ^* .





Better confinement at low q95 (#193778) than at high q95 (# 194288) at $B_t=2\text{T}$ for both => $\sim I_p$ scaling is valid for NT plasmas.



Why confinement increases with decreasing ρ^* ? Mean poloidal flow V ExB (N=0) increases with Bt for central plasma, ExB ‘well’ at the edge is larger for B=1T.





The normalized to Bohm $\chi_B = 1/16T_{eV}/B_T$ time and line (OMP) averaged heat conductivities as a function of ρ^* ($\psi_n=0.5$) for all scaled cases.
Gyro-Bohm scaling for NT.




X-point + SOL geometry is important!

Similar to #193778 plasma size/shape/parametres but w/o X-point in NT. Mirror flipped equilibrium for PT w/o X-point. BUT: because of zero potential perturbation at the boundary w/o X point (boundary conditions) edge turbulence was significantly reduced, no spreading to open field lines as with X-point. BUT : experimentally more turbulence toward the edge,




Preliminary: Bohm-like ρ^* scaling for PT , but w/o X-point here (similar heat conductivity for Bt=1T and Bt=2T). Should be done with X-point (future work).
 For NT with X-point: gyro-Bohm.




NT #193778 with X



PT w/o X



averaged: 0.38-0.58ms, $\rho^*(\psi_n = 0.5)$



Conclusions

1. Particles(gyro-kinetic ions, guiding center electrons) in global non-linear JOREK-GK code in realistic X-point geometry. Modelling of electrostatic turbulence : gyro-kinetic ions+ adiabatic (ITGs only) or kinetic electrons (TEM/ITGs) can be modelled.
2. Comparison of JOREK-GK with GENE, GS2 for TCV ($T_e > T_i$) in linear TEM regime : similar linear growth rates for single modes, which are usually larger in PT compared to NT.
3. Divertor heat fluxes in divertor for TCV cases: PT $\lambda q = 4\text{ mm}$; NT: $\lambda q = 3.5 \text{ mm}$.
4. Modelling of realistic DIII-D pulses at low q_{95} ,high q_{95} . Gyro-Bohm ρ^* confinement scaling in NT. Bohm ρ^* scaling in PT (preliminary, since done w/o X-point).
5. Future: A larger parameter range should be explored both experimentally and numerically to confirm the gyro-Bohm scaling for NT plasmas. Collisions model to improve (now only electron-ion). Decrease electron mass (now heavy electrons $m_i/m_e=100$)