



**HAL**  
open science

## Synthesis and Characterization of Polynuclear Actinide(IV) Species

Maeva Munoz, Christelle Tamain, Matthieu Viro, Dominique Guillaumont, Vallérie Vallet

► **To cite this version:**

Maeva Munoz, Christelle Tamain, Matthieu Viro, Dominique Guillaumont, Vallérie Vallet. Synthesis and Characterization of Polynuclear Actinide(IV) Species. Atalante 2024, Sep 2024, Avignon, France. cea-04736411

**HAL Id: cea-04736411**

**<https://cea.hal.science/cea-04736411v1>**

Submitted on 14 Oct 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Synthesis and Characterization of Polynuclear Actinide(IV) Species

Maeva Munoz<sup>1</sup>, Christelle Tamain<sup>1</sup>, Matthieu Viro<sup>2</sup>,  
Dominique Guillaumont<sup>1</sup>, Valérie Vallet<sup>3</sup>

<sup>1</sup> CEA, DES, ISEC, DMRC, Univ Montpellier, Marcoule, France. <sup>2</sup> ICSM, Univ Montpellier, CEA, CNRS, ENSCM, Marcoule, France. <sup>3</sup> Laboratoire PhLAM, UMR CNRS 8523, Université des Sciences et Technologies de Lille, Villeneuve d'Ascq, France

The actinide chemistry is largely influenced by the formation of polynuclear species that have attracted valuable interest over the last decades. These species have been proven to play a potential role in the management of waste, fuel reprocessing and even in biological media. The polynuclear species of tetravalent actinides (i.e., actinides clusters) discussed here materialize as molecules containing multiple metal centers connected by oxo and/or hydroxo groups. This forms the core of the cluster, often described by its nuclearity,  $N$ , which represent the number of actinides in it. These entities are formed through condensation reactions (referred to as olation and oxolation reactions) in the presence of water and are surface-stabilized by organic or inorganic ligands.<sup>[1]</sup>

In the solid state, various polynuclear species of actinides(IV) have been identified from XRD crystal structures, with various nuclearities ranging from 2 to 38. However, the hexamer consisting of 6 plutonium atoms ( $\text{Pu}_6$ ) is frequently observed.<sup>[1]</sup> This nuclearity has been studied and stabilized several times in the form of the  $[\text{Pu}_6\text{O}_4(\text{OH})_4]^{12+}$  core. Takao *et al.* investigated the formation of hexanuclear clusters of neptunium, uranium, and thorium in solution, in the presence of carboxylate ligands.<sup>[2]</sup> Our group have also studied the formation of plutonium hexameric clusters under various conditions. The results revealed similarities in UV-Vis spectra associated with this nuclearity.<sup>[1] [3, 4]</sup> However, studies in aqueous solutions are still inadequate, and it remains difficult to identify these polynuclear species in solution.

Additionally, recent studies suggest that these species could play a role in the formation mechanism of  $\text{PuO}_2$  colloidal nanoparticles.<sup>[5]</sup> Indeed, some hexanuclear Pu(IV) clusters have been identified as reactive intermediates in this process. Colloid formation is an irreversible process that may pose challenges in the context of used nuclear fuel reprocessing. Understanding the formation of polynuclear actinide species is thus crucial for comprehending and mastering the formation of colloids in solution.

The first objective of this study is to develop reliable spectroscopic databases, in order to help the identification of polynuclear species in solution. This work begins with the synthesis of plutonium(IV) single crystals characterized by X-ray diffraction (SCXRD), which specific spectra are collected with UV-visible and vibrational spectroscopies (IR and Raman). This approach will simplify the identification of these species in solution, enhance the understanding of the mechanisms involved in the formation of polynuclear species, as well as their evolution towards the formation of colloids. Preliminary results will be presented in the poster.

- [1] Knope, K. E.; Soderholm, L. Solution and Solid-State Structural Chemistry of Actinide Hydrates and Their Hydrolysis and Condensation Products. *Chem. Rev.*, **2013**, *113* (2), 944–994. <https://doi.org/10.1021/cr300212f>.
- [2] Takao, K.; Takao, S.; Scheinost, A. C.; Bernhard, G.; Hennig, C. Formation of Soluble Hexanuclear Neptunium(IV) Nanoclusters in Aqueous Solution: Growth Termination of Actinide(IV) Hydrous Oxides by Carboxylates. *Inorg. Chem.*, **2012**, *51* (3), 1336–1344. <https://doi.org/10.1021/ic201482n>.
- [3] Tamain, C.; Dumas, T.; Guillaumont, D.; Hennig, C.; Guilbaud, P. First Evidence of a Water-Soluble Plutonium(IV) Hexanuclear Cluster. *Eur. J. Inorg. Chem.*, **2016**, *2016* (22), 3536–3540. <https://doi.org/10.1002/ejic.201600656>.
- [4] Chupin, G.; Tamain, C.; Dumas, T.; Solari, P. L.; Moisy, P.; Guillaumont, D. Characterization of a Hexanuclear Plutonium(IV) Nanostructure in an Acetate Solution via Visible–Near Infrared Absorption Spectroscopy, Extended X-Ray Absorption Fine Structure Spectroscopy, and Density Functional Theory. *Inorg. Chem.*, **2022**, *61* (12), 4806–4817. <https://doi.org/10.1021/acs.inorgchem.1c02876>.
- [5] Cot-Auriol, M.; Viro, M.; Dumas, T.; Diat, O.; Menut, D.; Moisy, P.; Nikitenko, S. I. First Observation of  $[\text{Pu}_6(\text{OH})_4\text{O}_4]^{12+}$  Cluster during the Hydrolytic Formation of  $\text{PuO}_2$  Nanoparticles Using H/D Kinetic Isotope Effect. *Chem. Commun.*, **2022**, *58* (94), 13147–13150. <https://doi.org/10.1039/D2CC04990B>.